


Session Objectives

- Identify and describe the components of a history for a common gastrointestinal complaint.
- Describe the components and sequence of the abdominal physical exam.
- Isolate and describe normal abdominal exam findings.
- Define, describe, and identify the different types of bowel sounds and their clinical significance
- Correlate abdominal exam findings with likely etiologies



Case Presentation

- 44-year-old female with complaint of constipation which has been worsening over the past 3 to 4 months.
- What questions should you ask this patient?
- Any questions pertaining to osteopathic principles and practice?
- During your history taking consider the patient's discomfort in reporting certain complaints


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Definitions - Constipation

- Physicians:
 - Fewer than 3 bowel movements per week
- Patients:
 - Hard stools
 - Feeling of incomplete evacuation
 - Excessive straining
 - Sense of anorectal blockage during defecation
 - Need for manual maneuvers to defecate



Constipation - HPI

| HPI Components | What should you ask patient? |
|----------------------------|--|
| Onset | When did it start? Older patient new-onset → cancer |
| Palliation/ Provocation | What makes it better/worse, What have they tried? |
| Quality | Lumpy or hard stools, incomplete evacuation, excessive straining. Which of these is most distressing to the patient? |
| Severity | Frequency, intervals between bowel movements Prolonged straining, digital evacuation → Pelvic floor dysfunction Need laxative to produce loose stools, how often and what dosage Sensation of incomplete evacuation |
| Timing | Constant, intermittently occurring How many bowel movements per week? |

Source: Jonathan Gotfried, etal. Constipation. Merck Manual Professional, Updated April 2025
<https://www.merckmanuals.com/professional/gastrointestinal-disorders/symptoms-of-gastrointestinal-disorders/constipation?query=constipation>

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Constipation - HPI

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine

| HPI Components | What should you ask patient? |
|---------------------|--|
| Associated symptoms | Fever, nausea, abdominal pain Alternates with diarrhea → Irritable bowel syndrome Paradoxical diarrhea in elderly → fecal impaction Bloating and cramping → Irritable bowel syndrome Hematochezia → Colon cancer, diverticulosis, inflammatory bowel disease Weight loss ≥ 10 lbs → colon cancer |
| Other | Pain on DRE → Anal fissure, hemorrhoids Leakage of stool on DRE → Fecal impaction, rectal prolapse Medications → opioids, calcium channel blockers, iron supplements PMHX: Parkinson's, hypothyroidism, dementia |


Source: Jonathan Gotfried, etal. Constipation. Merck Manual Professional, Updated April 2025
<https://www.merckmanuals.com/professional/gastrointestinal-disorders/symptoms-of-gastrointestinal-disorders/constipation?query=constipation>



Anatomy



Examination Techniques



Abdominal Exam - Preparation

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Good Light
- Warm Room, hands, instruments
- Relaxed Patient:
 - Pillow or folded sheets under head and knees.
 - Patients arms at his/her sides
- Exposure – position drapes appropriately
- Empty Bladder
- Examine Area of Pain **Last**



Inspection



Inspect the Abdomen

- With the patient lying supine, the abdomen should be:
 - Completely exposed
 - Revealing all 4 quadrants
- Tell your patient that you are inspecting their abdomen.

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Inspect Abdominal Wall

- Contour
 - Symmetry, Masses
- Skin
 - Scars, striae
 - Dilated Veins, Discoloration
 - Rash or lesions
- Umbilicus
- Motion
- Pulsations

**Remember
to observe
from the
right side!**

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Inspect Abdominal Wall Contour

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



- Distention
 - Generalized
 - Localized
- Protuberant
- Scaphoid
- Flat



Inspect Abdomen for Scars

- Describe or diagram
- Include in the description:
 - Location
 - Size & Shape
 - Type (surgical/injury)
 - Healing stage
- Correlate with patient's history

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine




Inspect Abdomen for Dilated Veins

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- A few small visible veins may be normal.
- Abnormal dilation seen in:
 - Portal hypertension
 - Emaciation
 - IVC obstruction



Patient with caput medusa and ascites



Inspect Abdomen for Jaundice

- Pre-hepatic
 - Red blood cells rupture faster than the liver can conjugate bilirubin
- Hepatic
 - Liver's inability to conjugate or excrete bilirubin
- Post-hepatic
 - Flow of bile into the intestine is blocked

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Inspect Abdominal Umbilicus

- Observe:
 - Contour
 - Location
 - Inflammation
 - Bulges



NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine




Inspect Abdomen for Pulsations and Peristalsis

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Abdominal aortic aneurysm may be visible as a pulsatile mass
- Visible peristalsis is usually the result of intestinal obstruction



Visible peristalsis video:

<http://www.nejm.org/doi/full/10.1056/NEJMicm0910079>



Auscultation



Auscultation of the Abdomen

- Auscultate the abdomen **BEFORE** percussing or palpation as these maneuvers may alter frequency of bowel sounds.
- Place the diaphragm of your stethoscope gently on the abdomen
- Listen for bowel sounds: Note their frequency and character
- Listen for vascular bruits
- Auscultate the abdomen in **ALL 4 quadrants**

Auscultate
BEFORE
Palpation or
Percussion




Auscultation of Abdomen for Other Bowel Sounds

- **Borborygmi:** Prolonged gurgles of hyperperistalsis; stomach "growling" or "rumbling".
- **High-pitched tinkling sounds:** Suggestive of dilated bowel due to the presence of air and fluid under tension
- **Rushes of high-pitched** sounds that coincide with abdominal cramps: Suggestive of intestinal obstruction

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Percussion




Percuss and Measure the Liver: Mid-clavicular Line

- Percuss the liver in a downward or upward direction.
- Start at an area below the umbilicus (midclavicular line) and percuss up from abdominal tympany until dullness is appreciated (lower border of the liver)
- Next, starting at the nipple line percuss from lung resonance down toward liver dullness (upper border of the liver)
- When dealing with a patient with large breasts, ask her or him to displace the breast as necessary.


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Percuss the Spleen

- Percuss the spleen on the anterior lateral, not directly on the front of the abdomen.
- Percuss the lowest interspace in the left anterior axillary line (tympanitic).
- Ask the patient to take a deep breath, and percuss again
- Should stay tympanitic
- If becomes dull upon inspiration suggests splenic enlargement




Palpation




Palpate the Abdomen

- First, palpate the abdomen **lightly**.
 - To indentify tenderness, muscular resistance and superficial organs and masses.
- Then palpate **deeply** using finger pads, not finger tips.
 - To delineate masses (pregnant uterus, abdominal aortic aneurysm, or a distended bladder).
- **Note:** If patient complaint is “**abdominal pain**” make sure you examine the painful area **last**.




Palpate the Liver: *Bi-manual Compression*

- The left hand is placed under the right posterior ribs, and the right hand palpates the liver deeply



NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Palpate the Spleen

- Brace the posterior ribs with the left hand and palpate with the right hand
- Ask the patient to take a deep breath as the spleen is palpated



Palpation of Kidney

- Bimanual technique. Place one hand on patients back parallel to the 12th rib.
- Lift patients back.
- Place the other hand below the costal margin lateral and parallel to the rectus abdominus muscle.
- Ask patient to take a deep breath.
- At peak of inspiration attempt to capture patients kidney between your hands.

Bates' Guide to the Physical Examination and History Taking; 13th Ed. – Chapter 19

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine




Examiner stands on same side of kidney being palpated.



Palpation of Aortic Pulsations

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Summary Slide

- Points:
 - **Consider** patient's discomfort with discussing their problems
 - **Observe** from the patient from the right side of the patient
 - **Explain** to the patient what you are doing
 - **Observe** the patient for signs of pain, don't rely solely on verbal cues



Lecture Feedback Form: Link Below

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine



<https://comresearchdata.nyit.edu/redcap/surveys/?s=HRCY448FWYXREL4R>

