

Class of 2029

Physiology of Neurons & Synapses

- Raddy L. Ramos, PhD
- Assoc. Prof, Biomedical Sciences
- Email with questions or to set up a meeting: rramos02@nyit.edu



Do.
Make.
Heal.
Innovate.
Reinvent the Future.




Session objectives

- Students will be able to describe how different ionic conductances create the resting membrane potential of neurons.
- Students will be able to describe how different ionic conductances generate an action potential as well as the different phases of the action potential
- Students will be able to describe the anatomy and physiology of chemical and electric synapses.
- Students will be able to describe the major steps that characterize synaptic transmission beginning with an action potential in the presynaptic neuron to depolarization or hyperpolarization in the postsynaptic neuron
- Students will be able to describe several examples of the clinical relevance of ion channels and synaptic transmission.




Neuronal resting membrane potential (rVm)

- Neuronal membrane changes, electric activity, and synaptic connections creates behavior!
- rVm of neuron is “hyperpolarized” (more negative) relative to the extracellular side.
- rVm is $\sim -65\text{mV}$ in many central neurons but can vary by $\pm 15\text{mV}$.
- Membrane potential physiology also important to understand cardiovascular function.



Resting membrane potential of neurons

- Hyperpolarized rVm is established by...
- 1] Na/K pump ->> 3Na⁺ out / 2K⁺ in
- 2] membrane channels selectively permeable to certain ions “leak” K⁺ and “leak” Na⁺ channels.
- 40X more “leak” K than “leak” Na
- 3] unequal concentration of charged ions present in the intra vs. extracellular space determine direction and extent of ion flow.



Resting membrane potential simulation:

<https://neuromembrane.ualberta.ca/>

RESTING POTENTIAL Simulation

LEAK CHANNELS

Relative Permeability

Na⁺ : K⁺

0 : 0

Net Conductance (mS)

3

+

Na⁺/K⁺ PUMP

+

MEMBRANE SETTINGS

Initial Voltage (mV)

0.0

Specific Capacitance

1 (μF/cm²)

CONCENTRATION SETTINGS

[Na⁺] Inside (mM)

50

[Na⁺] Outside (mM)

50

[K⁺] Inside (mM)

50

[K⁺] Outside (mM)

50

SIMULATION SETTINGS

Total Time (min)

30

Outside cell

Inside cell

CREATE SIMULATION



Leak K^+ channels contribute the most but not entirely to the rVm (-65mV)

- If the K^+ channels contributed the sole ionic conductance at rest then rVm would be equal to E_K (-80mV)
- The rVm is more depolarized than E_K
- Leak Na^+ channels also contribute to rVm.
- The relative permeability of $K^+ : Na^+$ is approx 40:1.
- Thus rVm is closest to E_{K+} compared to other ions

Outside	Inside	Ratio Outside:Inside	E_{ion} (at 37°C)
$[K^+]_o = 5 \text{ mM}$	$[K^+]_i = 100 \text{ mM}$	1:20	-80 mV
$[Na^+]_o = 150 \text{ mM}$	$[Na^+]_i = 15 \text{ mM}$	10:1	62 mV
$[Ca^{2+}]_o = 2 \text{ mM}$	$[Ca^{2+}]_i = 0.0002 \text{ mM}$	10,000:1	123 mV
$[Cl^-]_o = 150 \text{ mM}$	$[Cl^-]_i = 13 \text{ mM}$	11.5:1	-65 mV



Conclusions (Part 1):

- Neurons exhibit a negative resting membrane potential due to...
- Na/K pump
- Membrane channels selectively permeable to certain ions especially leak K^+ (minimal leak Na^+ channels)
- 40:1 permeability ratio between K^+ vs Na^+ leak channels
- Unequal concentration of charged ions present in the intra vs. extracellular space determine direction and extent of ion flow and membrane potential change.



Action Potentials (APs)



APs are the information carrying electrical signal of neurons about the external world, motor commands, emotions, etc.

Neurons have a threshold for AP generation – depolarization to $\sim -40\text{mV}$ then... bang!


What determines the AP threshold? – the “voltage-gating” of voltage-gated Na^+ channels (activation voltage $\sim -40\text{mV}$)

APs are generated by the activation of voltage-gated Na^+ and K^+ channels.

Travel down the axon and cause neurotransmitter release which is used to communicate across neuronal connections (synaptic transmission).



<https://neuromembrane.ualberta.ca/>



?

ACTION POTENTIAL

Simulation

?

⌂

⬇

⬆

📄

⌵

⌵ LEAK CHANNELS

+

Relative Permeability

Na⁺

1

K⁺

40

Net Conductance (mS)

3

↶

⌵ NA⁺ CHANNEL

+

Conductance (mS)

120

Reverse Potential

0.0 mV

☒ Inactivation Gate

☐ TTX Neurotoxin

↶

⌵ K⁺ CHANNEL

+

Conductance (mS)

30

Reverse Potential



0.0 mV

⌵ MEMBRANE SETTINGS

⌵ CONCENTRATION SETTINGS

⌵ EXTERNAL CURRENT


Voltage-gated sodium channel inactivation



Membrane potential : Rest - depolarized - repolarize - rest

Channel inactivation is produced by a separate part of the protein – “known as the inactivation gate”
Genetic mutations that affect the “inactivation gate” result in channels being open for too long – **EPILEPSY!**
Drugs have been developed to modify channel inactivation – speed up or slow down inactivation.

4 Phases of the AP



- 1. Rising phase = initial rapid **depolarization** – opening of **voltage-gated** Na^+ channels
- 2. Overshoot = portion of the AP that is depolarized > 0 mV – opening, closing, inactivation of Na^+ channels and opening of K^+ channels.
- 3. Falling phase = rapid **hyperpolarization** – opening of **voltage gated** K^+ channels
- 4. Undershoot = portion of AP that is hyperpolarized past resting V_m toward (E_{K^+} , -80 mV) - closing of K^+ channels.

Clinical correlates: Mechanisms of neurotoxins



Tetrodotoxin (TTX)

- potent sodium channel blocker
- found in symbiotic bacteria in Pufferfish eg. genus *Lagocephalus*
- eaten in small amounts causes tingling and numbness of the mouth.
- eaten in large amounts can cause limb weakness, respiratory failure and cardiac arrest
- symptom onset varies from immediately to ~15mins




Saxitoxin

- potent sodium channel blocker
- produced by dinoflagellates found in shellfish
- causes "paralytic shellfish poisoning"
- similar symptoms as in TTX poisoning





α -dendrotoxin

- potent potassium channel blocker (Kv1)
- found in venom of the green mamba *Dendroaspis angusticeps*
- prolongs action potentials affecting neuro and cardiovascular function



Mechanism of drug action: Lidocaine


- local anesthetic used for a variety of applications
- Sodium channel blocker
- can be applied topically as a cream
- can be injected directly into nerve, tissue, CSF





Targeting VG Na channels for the treatment of pain (ex. after surgery).

Block action potentials in peripheral neurons (DRGs) carrying pain info to brain.

DRGs express voltage gated Na⁺ channel (Nav1.8) that is not found in central neurons.



VX-548 now called *suzetrigine* (Journavx)



<https://www.journavx.com/how-journavx-works>

VG Sodium channel gene mutations in PNS cause pain syndromes


- Mutations can change channel function **increasing** activity – too many action potentials, abnormal pain signals sent to CNS

1] Inherited Primary Erythromelalgia (Nav1.8) rare syndrome of intermittently **red, hot, painful extremities**. Usually affects the lower extremities (predominantly the feet) but may also involve the upper extremities (predominantly hands) and rarely involves the face. Symptoms present in the first two decades of life.

2] Familial Episodic Pain Syndrome (Nav1.7) also rare and characterized by intense, recurrent pain attacks, which often subside with age. Autonomic symptoms include excessive sweating, palpitations, and breathing difficulties. Pain can be in more proximal areas such as shoulders, chest, knees.

- Mutations can change channel function **reducing** activity – too few action potentials, reduced pain signals sent to CNS

1] Congenital Pain Insensitivity/ Congenital Analgesia (Nav1.7) condition characterized by the (near) complete absence of pain perception typically associated with noxious stimuli. Patients have painless injuries beginning in infancy but otherwise normal sensory modalities.





Conclusions (Part 2):

- The neuronal membrane contains voltage-gated Na^+ and K^+ channels that contribute to the generation of the action potential
- The action potential has distinct phases which are produced by the sequential opening and closing of Na^+ and K^+ channels
- Action potential generation can be modified by ion channel blockers such as TTX and lidocaine and Journavx.
- Gene mutations affecting voltage gated Na^+ channels produce pain syndromes

Neurons are connected to one another at the synapse



- the site where neurons functionally communicate with one another

Chemical synapses & neurotransmitter release




Glutamate synapses depolarize the postsynaptic neuron


*** Presynaptic **voltage-gated** **Ca²⁺ channels** cause neurotransmitter release at the terminal.





- AP induced release of Glutamate into cleft
- Ionotropic glutamate receptors (AMPA) have a Na⁺ pore. Glu binding opens channel and Na⁺ rushes into the cell (some K⁺ out of the cell).
- Depolarizes postsynaptic neuron which might contribute to AP generation in postsynaptic neuron.



Excitatory postsynaptic potentials- EPSPs



GABA synapses hyperpolarize the postsynaptic neuron



- AP induced release of GABA
- Ionotropic GABA_A receptors have a Cl^- pore.
- Cl^- rushes into the cell due to concentration gradient
- Make it harder for the postsynaptic neuron to fire APs


Boldog et. al 2018



Neurons are connected to one another at the synapse

- the site where neurons functionally communicate with one another

Electrical synapses:
Gap junctions/Connexons



Direct ion & small molecule flow from one neuron to the other. Usually bidirectional.

*Certain connexins expressed specific cell types.
~30 disease due to connexin gene mutations*








Figure 8.1

Gap junctions/Electrical synapses: no neurotransmitter used




Connors & Long 2004

- Gap junctions allow direct ion & small molecule flow from one neuron to the other which can depolarize or hyperpolarize the postsynaptic neuron
- Amplitude attenuation of currents
- Usually bidirectional
- Fastest type of neural communication <1ms
- Can depolarize neurons



Patellar-tendon reflex:

Great example of how neuronal connections and synaptic physiology creates behavior! Sensation & movement response




Sensory pathway:

Stretch-sensitive channels on 1a axons depolarize neuron and open voltage gated Na^+ channels causing AP.

Motor pathway:



1a neuron synapses on motor neuron and releases Glu. Motor neuron fires an AP and releases Ach at quad muscle causing contraction.



Synaptic Transmission & Behavior:



Neuromuscular synapse/junction

- synapse formed by motor neuron with a muscle fiber.



Neuromuscular synapse/junction

- synapse formed by motor neuron with a muscle fiber.



Muscle contraction & behavior!

Synaptic transmission

Presynaptic

- *Resting membrane potential*
- *Action potential generation*
- *Neurotransmitter synthesis*
- *Vesicle packaging of neurotransmitter*
- *Axonal calcium channels*
- *Reuptake of transmitter*

Synaptic cleft

- *Neurotransmitter breakdown*


Postsynaptic

- *Postsynaptic transmitter receptors*
Ionotropic/metabotropic

Synaptic mechanisms of disease

Lambert-Eaton Syndrome:


Auto-immune Disease of VGCC at the NMJ




- Autoimmune disease characterized by the production of antibodies against voltage gated calcium channels (VGCC) present in NMJ
- Symptoms include muscle weakness throughout body especially limbs & face and autonomic symptoms.
- Weak muscles (weakness is often relieved for a short time after exercise or exertion)
- Trouble walking, climbing stairs or getting up from a chair, Tingling sensation in the hands or feet
- Eyelid drooping, Fatigue, Dry mouth or dry eyes
- Trouble speaking and swallowing
- Trouble breathing, Bladder and bowel changes
- Erectile dysfunction, Decreased sweating
- Weight loss
- Some pharmacological immunological treatments available to modulate VGCC function

Myasthenia Gravis:

Disease of AchR at the NMJ



 AChR autoantibodies

- Autoimmune disease characterized by the production of antibodies against AchR
- Symptoms include muscle weakness throughout body especially limbs and face.
- Some pharmacological and immunological treatments available

New drug on the market: Vyvgart (FDA 2021)



<https://vyvgarthcp.com/gmg/about/moa>



Conclusions (Part 3):

- Neurons are connect to one another via chemical and electrical synapses
- The postsynaptic effect of a given neurotransmitter will depend on the type of transmitter itself as well as the type of receptor found on the postsynaptic neuron (ionotropic vs. metabotropic)
- All of the players (neurons, channels, myelin, transmitters, etc) involved in synaptic transmission are vulnerable to perturbation in disease and are drug-able targets for therapy.

Lecture Feedback Form:

<https://comresearchdata.nyit.edu/redcap/surveys/?s=HRCY448FWYXREL4R>

