

Anatomy: Orbit and Extraocular Muscles

Jonathan Geisler, Ph.D.
Department of Anatomy
jgeisler@nyit.edu

Session Objectives

- Infer what structures can be affected by blowout fractures of the orbit.
- Differentiate actions and functions of extraocular muscles.
- Describe how extraocular muscles work in pairs to induce or avoid torsion.
- Describe how the extraocular muscles work in pairs to move the eye in primary position.
- Predict lesions in nerves of the orbit based on tests of extraocular muscles.

Part 1: Orbit Boundaries and Contents

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine




Axial MRI of Head

- Orbit is full of adipose tissue!

NEW YORK INSTITUTE OF TECHNOLOGY


College of Osteopathic
Medicine



From COA, Moore and Dalley (2006)

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine




From Netter's Atlas of Anatomy

Trauma in the Orbit

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



From COA, Moore, Dalley, Agur, 2014


- When under pressure orbit contents can break into surrounding spaces.
- Called blowout fractures

Levator palpebrae superioris


- Origin: lesser wing of sphenoid
- Insertion: superior tarsus and superficial fascia of eyelid
- Contains smooth muscle fibers (superior tarsal or Müller's muscle)

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



From COA, Moore, Dalley, Agur, 2014





From Netter's Atlas of Anatomy

Recti Muscles

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




- Superior, inferior, medial, and lateral
- Originate from common tendinous ring
- Insert into sclera immediately posterior to cornea

From COA, Moore and Dalley, Agur, 2005

Superior oblique

- Originates from lesser wing of sphenoid
- Passes through *trochlea*
- Inserts into sclera deep to superior rectus




Oblique Muscles

Inferior oblique

- Originates from medial orbital wall
- Passes posterolaterally beneath eyeball
- Inserts into sclera deep to lateral rectus


NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



From COA, Moore, Dalley, Agur, 2014

Innervation




From COA, Moore, Dalley, Agur, 2014

- $SO_4LR_6AO_3$
- Superior Oblique:
Trochlear n. (CNIV)
- Lateral Rectus: Abducens
n. (CNVI)
- All Others: Oculomotor n.
(CNIII)

- Superior Tarsal (Muller's) muscle: postganglionic sympathetics from superior cervical ganglion

Part 2: Simple Movements of the Eye


Actions on Eyeball




Actions vs. functions of extraocular muscles

- *Action* is what it would do if acted in isolation
- A muscle's *function* is what it *actually* contributes to a particular motion
- Sometimes these are the same – example: Levator palpebrae superioris

*Let's start with the
simple cases ...*







Elevation of the Abducted Eye

**lateral
rectus**


Elevation of the Abducted Eye




Elevation of the Abducted Eye




Depression of the Abducted Eye




Depression of the Abducted Eye







Depression of the
Adducted Eye



Depression of the
Adducted Eye




**Depression of the
Adducted Eye**



Elevation of the
Adducted Eye

**medial
rectus**



**inferior
oblique**

**Elevation of the
Adducted Eye**

To Summarize...

- Superior rectus is a *strong elevator* of the *already abducted* eyeball.
- Inferior rectus is a *strong depressor* of the *already abducted* eyeball.
- Inferior oblique is a *strong elevator* of the *already adducted* eye
- Superior oblique is a *strong depressor* of the *already adducted* eye

Part 3: Complex Movements of the Eye


Textbook Actions of Extraocular Muscles

NEW YORK INSTITUTE
OF TECHNOLOGY


College of Osteopathic
Medicine

Agur & Dalley 2005

TABLE 8.5. MUSCLES OF ORBIT



- Eye is in
PRIMARY POSITION




Torsion Often Accompanies Actions of Extraocular Muscles

Axes of Rotation of Extraocular Muscles

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine



Axes of Rotation of Extraocular Muscles

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine



Elevation Without Torsion

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine




Elevation (right eye)

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Elevation (left eye)




**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Depression Without Torsion

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine



Intorsion Without Elevation or Depression

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine



Extorsion Without Elevation or Depression

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine



Extorsion Without Elevation or Depression

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Part 4: Nerve lesions affecting extraocular muscles

Testing Extraocular Muscles and the Nerves that Innervate Them

**NEW YORK INSTITUTE
OF TECHNOLOGY**



College of Osteopathic
Medicine



- “H” shaped pattern is used to test the extraocular muscles
- “Forcing” function = action

Patient presents
with....

“Down and out”



Patient presents
with....

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Bielschowsky's sign



From www.eyeatlas.com

Summary Slide

- Contents of the orbit are not compressible. Thus pressure requires defects in its base or walls.
- Action and Function often mean very different things for extraocular muscles.
- Torsion occurs when the optic axis is at an angle to axis of the orbit.
- Torsion is helpful when tilting the head.
- Action and Function are usually described relative to primary position. Obviously the eye can be looking elsewhere.
- Extraocular muscle function is tested by having the patient NOT in primary position.

Lecture Feedback Form:

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

<https://comresearchdata.nyit.edu/redcap/surveys/?s=HRCY448FWYXREL4R>