Session Objectives

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

1. List and differentiate bacterial infections of the respiratory system and explain the diseases they cause.
2. Describe the unique microbiologic features and any associated virulence factors for each pathogen
3. Explain any relevant symptoms and diagnostic tests for each pathogen.
4. Describe any treatment or prevention strategy for each pathogen.

Outline


NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Healthcare-Associated and Ventilator Associated Pneumonia
  - *Pseudomonas aeruginosa (also structural lung disease with CF)*
  - *Acinetobacter baumannii*
  - *Escherichia coli (and other enteric bacteria)*
  - *Serratia marcescens*
  - *Methicillin-resistant staphylococcus aureus*
- Aspiration Pneumonia
  - *Actinomyces israelli*
- Immunocompromised Hosts/Structural Lung Disease
  - *Nocardia asteroides*
  - *Mycobacterium tuberculosis*
  - *Burkholderia cepacia (cystic fibrosis)*
- Zoonotic/Occupational Exposures
  - *Coxiella burnetii*
  - *Bacillus anthracis*
  - *Francisella tularensis*
  - *Yersinia pestis*

Pseudomonas aeruginosa


NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- **Characteristics:**
  - Gram –ve bacillus, aerobic, motile
  - Oxidase & Catalase +ve
  - Blue-green pigment (pyocyanin)
  - Grape-like odor
  - Likes moist environments
- Virulence
  - Encapsulated
  - LPS Endotoxin (fever, shock)
  - Exotoxin A (inactivates EF-2)
- Associated with Cystic Fibrosis patients



PSEUDOMONAS is associated with:
Pneumonia
Sepsis
Ecthyma gangrenosum
UTI
Diabetes
Osteomyelitis
Mucoid polysaccharide capsule
Otitis externa
Nosocomial infx
Addiction (IV drug use)
Skin infx (hot tub folliculitis)

Nocardia asteroides




- Characteristics:
  - Gram +ve; Branching filament
  - Aerobic; non-motile
  - Catalase +ve
  - Weakly acid fast (mycolic acid in cell wall)
  - Found in soil; transmitted via inhalation
- Virulence:
  - "**cord factor**" (CF): facilitates intracellular survival
- Causes pulmonary infections in **immunocompromised** that can disseminate to other organs (ex: brain, skin)
- Tx: sulfonamides

Mycobacterium tuberculosis

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Characteristics:
  - Rod shaped, aerobic
  - Does not Gram stain well
  - Auramine-Rhodamine stain can replace acid-fast
  - Cultures on Lowenstein-Jensen medium
  - Facultative intracellular
- Virulence:
  - cord factor: activates macrophages (promoting granuloma formation) and induces release of TNF-α
  - Sulfatides: inhibit phagolysosomal fusion




A

Source: S. Riedel, J.A. Hobden, S. Miller, S.A. Morse, T.A. Mietzner, B. Detrick, T.G. Mitchell, J.A. Sakanari, P. Hotez, R. Mejia Jawetz, Melnick, & Adelberg's Medical Microbiology, 28e
Copyright © McGraw-Hill Education. All rights reserved.

Mycobacterium tuberculosis

NEW YORK INSTITUTE


- Tuberculosis Pathogenesis
  - Inhaled mycobacteria → engulfed by alveolar macrophages and replicate
    - More lymphocyte and macrophage recruitment → CASEATING GRANULOMA
    - GHON FOCUS + HILAR LYMPHADENOPATHY = GHON COMPLEX (primary TB)
  - About 90% of people with Primary TB → <u>Latent TB</u> (Dormant; not infectious)
  - Less than 10% may progress → <u>Primary progressive TB</u>
  - If spreads to bloodstream → <u>Miliary or Extrapulmonary TB</u>



Mycobacterium tuberculosis




- Tuberculosis Diagnosis
  - Tuberculin Skin Test
    - Measure induration 48-72 hours after injection
    - + PPD: active or latent TB; BCG vaccine (false positive)
  - Interferon Gamma Release Assay
    - Good for those who are likely to have false PPD results
  - Sputum sample and culture
  - imaging with chest X-ray
    - If negative → latent TB
    - If necrosis of upper lobes → Active TB (primary progressive or reactivated TB)
    - If “millet seeds” diffusely across lungs → Miliary TB



Coxiella burnetii

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Characteristics:
  - Considered gram –ve but doesn't Gram stain well
  - Obligate intracellular: grows/replicates inside host cells
  - "*Rickettsia*-like"; No vector
  - Is a bacteria but has atypical spore-like structure



- <u>Q-fever:</u>
  - Inhaled as aerosol from cattle/sheep amniotic fluid/urin
  - S/Sx: headache, cough, flulike sx, atypical pneumonia
  - May lead to endocarditis, pneumonitis, and hepatitis
  - Tx: doxycycline

Bacillus anthracis

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Anthrax:
  - Infection through spore inhalation, ingested, or injured skin from infected animals
  - Cutaneous Anthrax: painless, necrotic papule surrounded by vesicles→ black eschar
  - **Pulmonary Anthrax:** inhalation of spores (from animals; also, a bioweapon) → flulike sx that rapidly progress to fever, pulmonary hemorrhage, mediastinitis and shock
    - Rapidly fatal
    - AKA Wool-sorter's disease
    - Tx: ciprofloxacin



Yersinia pestis

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- **Bubonic Plague:**
  - 40-100% mortality
  - s/sx:
    - Sudden onset of high fever, chills, weakness, rapid progression to septic shock
    - Intense inflammation develops in enlarged lymph nodes→ buboes
    - Can stay in lymph nodes, but often reaches bloodstream → meningitis, pneumonia, septic shock, or multi-organ failure
  - **Primary pneumonic plague:**
    - inhale infective droplets from coughing patient
  - Tx: streptomycin



Practice Question #1

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- **Q:** An 80-year-old female with PMHx of T2DM and dementia is brought to the hospital due to three weeks of persistent fever, pleuritic chest pain, and cough despite treatment with levofloxacin for suspected pneumonia. Currently her vitals are T:100.8 F, BP 130/78; HR: 88; RR: 18; O2: 96% RA. On physical exam, the patient has decreased breath sounds of the right lower lung field. Chest CT is ordered and reveals right lower lobe infiltrate, right pleural effusion, and right-sided hilar lymphadenopathy. What is the most likely cause of this patient's condition?

  A. Mycobacterium tuberculosis

  B. Streptococcus pneumoniae

  C. Escherichia coli

  D. Staphylococcus aureus

  E. Mycoplasma pneumoniae

Practice Question #2

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Q: A 44-year-old wool worker develops fever, malaise, and rapidly progressive shortness of breath. Chest X-ray reveals a widened mediastinum. Blood culture grows a large, Gram-positive rod. Which of the following virulence factors explains the pathogenicity of this organism?

  A. Protein A that binds Fc region of IgG
  B. Polysaccharide capsule resisting opsonization
  C. Poly-D-glutamate capsule inhibiting phagocytosis
  D. Lipooligosaccharide endotoxin activating macrophages
  E. Heat-labile toxin increasing cAMP

Practice Question #3

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Q: A 42-year-old farmer presents to the hospital with two days of fever, severe headache, and cough. The patient states he started feeling symptoms a day after he had to assist with delivering a sheep. Gram stain only shows inflammatory cells. Which of the following organisms is the most likely cause of this patient's illness?

  A. Francisella tularensis
  B. Yersinia pestis
  C. Coxiella burnetii
  D. Bacillus anthracis
  E. Pasteurella multocida

Summary Slide

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Healthcare-Associated and Ventilator Associated Pneumonia
  - *Pseudomonas aeruginosa:* Gram –ve rod; catalase +ve, oxidase +ve; aerobic; motile; grape odor; moist environments; blue-green pigment on agar; seen in CF pts
  - *Acinetobacter baumannii:* Gram -ve; aerobic; catalase +ve; high resistance
  - *Escherichia coli:* Gram –ve rod; indole +; mannitol fermenter; K-capsule virulence causes pneumonia
  - *Serratia marcescens:* Gram -ve; catalase +ve; slow fermenter on MacConkey; red pigment
  - *Methicillin-resistant staphylococcus aureus:* Gram +ve cocci in clusters; β-hemolytic; catalase +ve; coagulase +ve; resistance through altered PBP's due to *mecA* gene
- Aspiration Pneumonia
  - *Actinomyces israelli:* Gram +ve branching filaments; anaerobe; yellow sulfur granules; associated with dental caries/extractions and if aspirated → pneumonia

Summary Slide

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Immunocompromised Hosts/Structural Lung Disease
  - *Nocardia asteroides:* Gram +ve branching filament; aerobic; non-motile; catalase +ve; weakly acid fast; cord factor virulence; inhaled soil; pulm infx seen in immunocompromised

  - *Mycobacterium tuberculosis:* aerobic rod; does not Gram stain well; acid fast; Lowenstein-Jensen medium; facultative intracellular in alveolar macrophages; virulence through cord factor and sulfatides; caseating granulomas; ghon focus + hilar adenopathy = ghon complex in primary TB; miliary TB; Tuberculin skin test; IFN gamma; RIPE tx

  - *Burkholderia cepacia:* Gram –ve rod; aerobic; catalase +ve; oxidase +ve; pneumonia in pt with CF

Summary Slide

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Zoonotic/Occupational Exposures
  - *Coxiella burnetii:* doesn't gram stain well; obligate intracellular; Q-fever
  - *Bacillus anthracis:* Aerobic Gram +ve spore forming; poly D-glutamate capsule; Lethal toxin; edema toxin; Pulmonary anthrax
  - *Francisella tularensis:* Gram –ve coccobacillus; facultative intracellular; ticks and wild rabbits; Tularemia
  - *Yersinia pestis:* Gram –ve bacillus; nonmotile; facultative intracellular anaerobe; safety pin-like appearance; The Plague