


Session Objectives

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

- Identify **somatosensory receptors** and their encoding.
- Describe the transmission of **touch and proprioception** to the cerebral cortex.
- Describe the transmission of **pain and temperature** information to the cerebral cortex.
- Understand the relation between somatosensory lesions and clinical symptoms.



See **detailed objectives** in CPG.

PLEASE READ THE Scholar Rx Brick!

Session Objectives Mapped to COMLEX Blueprint

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

- **Dimension 2**
 - **Clinical presentation 4.** Patient presentations related to the nervous system and mental health
 - 4.11 Sensory disturbances and pain
 - tactile disturbances, including sensory loss, numbness, vibration/temperature/proprioception loss, tingling, and paresthesia
 - pain, chronic nonmalignant
 - pain, neuropathic, nociceptive, mixed, sympathetic
 - nerve-, muscle-, and pain-related syndromes, including complex regional pain syndrome, post-herpetic neuralgia, meralgia paresthetica

Session Objectives Mapped to USMLE Blueprint

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

V. Nervous System & Special Senses

A. Normal Processes

2. Organ structure and function

iv. Sensory systems

a. general sensory modalities, including sharp, dull, temperature, vibratory, and proprioception

B. Abnormal Processes: Health and Health Maintenance, Screening, Diagnosis, Management, Risks, Prognosis

5. Cranial and peripheral nerve disorders


iii. Neurologic pain syndromes: complex regional pain syndrome (reflex sympathetic dystrophy, causalgia); fibromyalgia; postherpetic neuralgia; phantom limb pain/syndrome; thalamic pain syndrome; trigeminal neuralgia

Classification of peripheral fibers

Myelinated

Unmyelinated

Based on conduction velocity




Based on axonal diameters
(no need to remember)



Reasonable Simplification

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine




Conduction Speed

A strong electrical stimulus of the skin will evoke a compound action-potential in a nerve fiber.

Its complex waveform reflects the summed response of individual axons having different diameters or myelination

Note the delayed response by C fibers



**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Diversity of receptors for somatosensation

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



1 Rod
3 Cones



1 Hair cell

5 touch
2 proprioception
1 force, 1 joint, 1 hair
> 2 temperature
> 2 pain/itch/ache
visceral
(plus more not mentioned &
being discovered)

Categories of receptors for somatosensation

Based on the **source** of stimuli

- **Interoceptors:**


- Internal stimuli within the body, e.g., baroreceptors, pH sensors

- **Exteroceptors**

- External stimuli outside the body, e.g., skin touch, heat sensors

- **Proprioceptors**

- Position changes of the body parts, e.g., receptors at joints



Based on the **types** of stimuli

- **Chemoreceptors**

- Smell, taste, pH, etc.

- **Photoreceptors**

- Visual signals

- **Thermoreceptors**

- Temperature sensors

- **Mechanoreceptors**

- Detecting physical deformation


- **Nociceptors** (Latin “*nocere*” = to hurt)

- Noxious stimuli: mechanical, thermal, chemical, etc.

Skin/cutis is richly embedded with touch sensors

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine




These are “exteroceptors”, also “mechanoreceptors” (type of receptors).


Source: Kandell and Schwartz, Principles of Neuroscience

Touch receptors have different sensitivity


Receptors




Meissner's
corpuscle




Merkel
cells




Pacinian
corpuscle



Ruffini
endings



Small RF



Small RF



Large RF



Large RF


Fast Adapting

Slow Adapting

Fast Adapting

Slow Adapting

Neural
spike train



Stimulus




Touch

Vibration, pressure

Touch sensitivity varies across body surface

- Reflects **density** of receptors
- Packed highest where there is most important physical interaction: **hands** and **face**



Use two point-discrimination to assess: is it “one” or “two”?



Golgi tendon organs sense force at the tendon

NEW YORK INSTITUTE
TECHNOLOGY

College of Osteopathic
Medicine



Free nerve endings wind through collagen

Pulling force at the junction (from external load or active muscle)
bends the free nerve endings, opens the ion channels,
and depolarizes the membrane

Joint receptors are sensitive to capsule strain

NEW YORK INSTITUTE
OF TECHNOLOGY


College of Osteopathic
Medicine

Free nerve endings embedded
within the joint capsule

Insensitive to limb position within
its normal range of motion

Detects limb position at
extremes and/or inappropriate
directions



For example, they are recruited
during valgus/varus motion of
the knee



Descending modulation of pain

NEW YORK INSTITUTE
TECHNOLOGY

College of Osteopathic
Medicine



Fill in the blank

Summary of sensory receptors

Receptor type	Characteristics	Fiber type	Conduc tion
Touch (mechanoreceptors)			
Pain (nociceptors)			
Proprioception (proprioceptors)			

Pause
here...

Summary of sensory receptors

Receptor type	Characteristics	Fiber type	Conduction
Touch (mechanoreceptor) Meissner's corpuscles Merkel cells Pacinian corpuscles Ruffini endings	Light touch, fast adapting, small RF Static touch, slow adapting, small RF Vibration & pressure, fast adapting, large RF Pressure, slow adapting, large RF	A β , large diameter, myelinated	Fast
Pain (nociceptors) Free nerve endings Noxious stimuli (thermal, chemical, etc.)	Sharp pain Dull pain	A δ , small diameter, myelinated C, small diameter, unmyelinated	Fast Slow
Proprioception Free nerve endings Joint receptors Golgi tendon organ	Extreme limb position or directions Pulling force at the junction of nerve endings and tendon collagen fibers	A α , large diameter, myelinated	Fast

Anatomical Organization of the Nervous System

- 2° Afferents for pain and temperature cross at the level they enter the spinal cord


- 2° Afferents for touch and proprioception cross at the level of the brainstem

Damage at X

- **Ipsilateral** loss of touch and proprioception
- **Contralateral** loss of pain and temperature

Damage at Y

- **Contralateral** loss of both sensory modalities



**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Dorsal Column-Medial Lemniscus Pathway (DCML)

4° - Primary
somatosensor
y
cortex

3° - Ventral posterior lateral
(VPL) thalamic nucleus

Conveys "conscious proprioception"
and "discriminative touch"
Above & below T₆

Decussates at medulla

2° - Nucleus gracilis

2° - Nucleus cuneatus



1° - Dorsal root
ganglion

1° - Dorsal root
ganglion


1° = 1st order in series

2° = 2nd order in series

3° = 3rd order in series



Internal capsule




Medial lemniscus

Dorsal columns

+ Fasciculus Cuneatus
(upper half of body)

Fasciculus Gracilis
(lower half of body)



Somatosensation in the head processed by three nuclei

Principal/main/pontine trigeminal nucleus (pons)


- analogous to **dorsal column nuclei** for DCML
- receives **touch and proprioceptive** information from face, cornea, tongue, ear, oronasal mucous membranes, dura, oropharynx, and laryngopharynx

Spinal trigeminal nucleus (medulla)


- analogous to **nucleus proprius** for STT
- receives **nociceptive** signals and **temperature** info from the same areas as the above

Mesencephalic trigeminal nucleus (midbrain)


- Analogous to dorsal root ganglia (**DRG**), where the cell bodies of pseudounipolar neurons (**1st order** neurons) for the **proprioception** of jaw muscles and periodontal regions are located
- Receives proprioceptive information from the mandible and projects to the **trigeminal motor nucleus** to mediate **jaw jerk reflex**



Pontine/principal trigeminal pathway




- Conveys “conscious proprioception” and “discriminative touch” of the head
- analogous to DCML



YORK INSTITUTE
CHNOLOGY

e of Osteopathic
ne

Sensory homunculus of the somatosensory cortex



Penfield and Rasmussen (1950)
The cerebral cortex of man

Primary somatosensory cortex is the **entire post-central gyrus including medial wall.**

Classically, this is the first receiving station within cortex for somatosensory information. Unimodal (no visual or auditory processing which occurs in more posterior parietal cortex).

“Somatotopy” – nearby portions of the body are represented by nearby portions of cortex.

Area of cortex for a given portion of the body/face reflects the density of sensory innervation.

The **leg** is most **medial**, then trunk, **hand**, **face**, and then oral/pharynx.



What kind of deficits would be presented when the lesions are at these levels?

NEW YORK INSTITUTE
OF TECHNOLOGY



Touch/Proprioception

Pain/Temperature


Forebrain





Pons





Medulla



Spinal
cord



Primary
afferent




of Osteopathic
ne

Pause
here...


What kind of deficits at different levels?

Touch/Proprioception

Pain/Temperature




Contralateral
Face & Body




Contralateral
Face & Body

Forebrain

Pons




Ipsi-/Contra-lateral face;
Contralateral body



Ipsi-/Contra-lateral face;
Contralateral body

Medulla




Contralateral body




Ipsi-/Contra-lateral face;
Contralateral body

Spinal
cord




Ipsilateral body




Contralateral body

Primary
afferent



Ipsilateral dermatome




Ipsilateral dermatome

Find the lesion

At what level?

Which side?





**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine


Pause
here...

Brown-Sequard syndrome

At what level?
Which side?



NEW YORK INSTITUTE
OF TECHNOLOGY



All lesions at the lower body – damage to the lumbar cord:

1. loss of all sensation **at the level** of the lesion at the left side – **ipsilateral dorsal horn** lesion
2. loss of proprioception, vibration, discriminative touch, and tactile sense **below the level** of the lesion at the left side – **ipsilateral DCML** lesion
3. loss of pain, temp, and crude touch **below the level** of the lesion at the right side – **contralateral spinothalamic tract** lesion
4. LMN signs **at the level** of the lesion at the left side – **ipsilateral ventral horn** lesion
5. UMN signs **below the level** of lesion at the left side – **ipsilateral corticospinal tract** lesion