

Session Objectives

- 1) Understand the origin and destination of descending cerebral motor tracts
- 2) Identify the organization and destination of descending brainstem tracts
- 3) Describe the organization and capabilities of the corticospinal tract
- 4) Describe the organization and capabilities of the corticobulbar tract
- 5) Understand the difference between upper and lower motor neurons and their associated dysfunctions as well as brainstem damage

Nolte's The Human Brain: An introduction to its functional anatomy

Chapter 18 Overview of motor systems



"Motor Control Systems Involve Both Hierarchical and Parallel Connections" and afterwards.

<http://arktos.nyit.edu/login?url=https://www-clinicalkey-com.arktos.nyit.edu/#!/content/book/3-s2.0-B9780323653985000187>

Lower motor neurons versus Upper motor neurons

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Nerves and ganglia are in PNS.
Tracts and nuclei are in CNS.

Lower motor neurons (also termed “motoneurons”) innervate skeletal muscle.

Cells bodies in spinal cord or cranial nerve nuclei.
Their tracts rarely decussate.
Their nerves never decussate.

Upper motor neurons have cell bodies at the cerebral cortex and brainstem. These project to lower motor neurons of the brainstem and spinal cord, not to muscles.



Variety of projection patterns: some decussate, some don't, and some project bilaterally.

The pyramidal system = cerebral cortical projections to the **spinal cord** and **brainstem**

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine


Confusing designation since only the **corticospinal tract** (for the body and limbs) forms the medullary pyramids, not the **corticobulbar tracts** (for the cranial nerve nuclei).



Converging inputs to the pyramidal system

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




These systems do not act on lower motor neurons, but on upper motor neurons of the cerebral cortex and brainstem.

That's why the basal ganglia and cerebellum are called “extra-pyramidal”.

“Extrapyramidal symptoms” are dysfunctions of the basal ganglia (typically) and cerebellum.

Cerebral cortex (e.g. pyramidal system) is linked to ipsilateral basal ganglia and contralateral cerebellum.

Corticospinal tract



NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine

Corticospinal fibers converge and pass downward through the posterior limb of the **internal capsule***, continue as the **crus cerebri**, disperse in the ventral **pons**, regroup in the medulla and come to the ventral surface as the pyramid where they mostly decussate at the lowermost portion and descend the spinal cord to their relevant level.

Damage to anywhere above decussation leads to motor deficits on the contralateral side of the body.


Damage to the lateral spinal cord leads to motor deficits on the same side.

Lesion of corticospinal tract (upper motor neurons)

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Augmented stretch reflexes



Healthy

Clinical


Thillman et al (1991) The mechanism of spastic muscle hypertonus.
Variation in reflex gain over the time course of spasticity.

Abnormal “synergies”



Involuntary elbow and wrist flexion especially during arm abduction

Altered walking



“Circumduction gait” - compensates for poor weight-bearing and ability to lift leg


<https://www.youtube.com/watch?v=y160w4sAQNw>

Spasticity from stroke or spinal cord damage can take days to weeks to develop.
The immediate motor symptom is weakness and flaccidity called “spinal shock”.

Lesion of corticospinal tract (upper motor neurons)

Normal

Babinski sign



Normally present for infants < 1 year


Also post-seizure states via
diminished/altered pyramidal control

Lesion of ventral horn (lower motor neuron) or its efferent nerve

Flaccid paralysis (as opposed to spastic paralysis)


- Diminished voluntary strength
- Diminished reflexes & muscle tone
- Fasciculations (spontaneous muscle twitches)
- Muscle atrophy (takes time)

Example shows atrophy of left deltoid due to damage of axillary nerve.



Eichinger (2016) Outcomes following closed axillary nerve injury

Corticobulbar tract




More ventral origin than corticospinal tract.

Projects to cranial nerve nuclei in the brain stem.

Cerebral cortex = “upper motor neurons”

Cranial nerve nuclei = “lower motor neurons”

Corticobulbar tract



Bilaterally innervates cranial nuclei controlling jaw, tongue, larynx, pharynx, and scm/upper trapezius.

EXCEPT: *Fibers projecting to **facial nucleus** show distinctive pattern:*


Contralateral nucleus receives innervation from whole corticobulbar tract (upper and lower face).

Ipsilateral facial nucleus only receives innervation for controlling upper face.

No redundancy for lower face. No compensation for loss.

Input to all other nuclei is bilateral though hypoglossal has more from contralateral cerebral cortex, i.e. partial compensation with tongue control.

Lesioning corticobulbar tract (upper motor neuron)



CN 7/Facial nerve

Remember forehead /eyebrow cortical motor neurons project **bilaterally** to facial motor nuclei, but cheek cortical motor neurons only project **contralaterally** to facial motor nuclei

Patient can raise their eyebrows and close eyelids, but cannot voluntarily move their lower 1/2 of face

Motor issues from a unilateral lesion of the corticobulbar track can also include the tongue -> deviation away from the lesioned side of cerebral cortex.

Lesioning corticobulbar tract (upper motor neuron)

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Remember forehead /eyebrow cortical motor neurons project **bilaterally** to facial motor nuclei, but cheek cortical motor neurons only project **contralaterally** to facial motor nuclei


Patient can raise their eyebrows and close eyelids, but cannot voluntarily move their lower 1/2 of face

Motor issues from a unilateral lesion of the corticobulbar track can also include the tongue -> deviation away from the lesioned side of cerebral cortex.

Brainstem tracts: Rubrospinal tract

**NEW YORK INSTITUTE
OF TECHNOLOGY**



College of Osteopathic
Medicine



- In midbrain
- Contralateral upper limbs
- Very small tract in humans, but large in monkeys (typical animal model)
- Red nucleus is very large in humans and has strong linkage to cerebellum

Brainstem tracts: Reticulospinal tract

MLR
(mesecephalic
locomotor
region)



- Stimulation leads to forward location Musienko (2012) Spinal and supraspinal control of the direction of stepping during locomotion



Asymmetrical tonic reflex – elbow flexion when head turns away, other arm extends. Present in infants, re-expressed with cortical stroke

Startle reflex - protective shoulder shrug and elbow flexion elicited by loud unexpected sound



Brainstem tracts: Vestibulospinal tract

- Medial vestibulospinal tract controls head and neck
- Lateral vestibulospinal tract has strong extensor activation for limb and axial muscles, e.g. activation will extend ipsilateral legs
- “Anti-gravity” function



“Locked-In” syndrome




- Pontine lesion interrupts corticospinal & corticobulbar tracts
- Lose all voluntary control, except some eye movements and blinks
- Remain conscious and aware, normal sleep-cycle but “locked in” their body.
- Difficult to determine missed cases, misdiagnosed as persistent vegetative state

Improved technology to allow faster interactions

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine



Summary

- The corticospinal tract arises from the precentral gyrus (mostly M1) and rostral adjacent cortex (PMC and SMA) and projects to the contralateral spinal cord.
- The corticobulbar tract arises from the same areas and projects bilaterally to the cranial nerve nuclei, excepting the facial nucleus.
- Upper motor neuron damage can be expressed as weakness and spastic paralysis whereas lower motor neuron damage is expressed flaccid paralysis.
- Different descending brainstem tracts have different projection patterns.
- Damage above and below the red nucleus leads to decorticate and decerebrate rigidity, respectively.