



Brachial Plexus: Lesions and Neuropathies

Principles and Practices of Osteopathic Medicine I

Matthew Mihlbachler, Ph.D.
Associate Professor, Anatomy
mmihlbac@nyit.edu

Part 1: introduction, dermatomes, myotomes,
brachial plexus review

Session Objectives


- Understand spinal nerve injuries (radiculopathies) and their relationships with myotomes and dermatomes
- Review the basic architecture of the brachial plexus, roots to branches
- Identify upper limb neurological deficiencies by specific muscle and sensory deficits

Things to Remember

- Brachial plexus, distribution of peripheral nerves, muscles attachments and actions
- This is not a complete survey of brachial plexus neuropathies or symptoms
- Variations in the brachial plexus may cause variation in signs and symptoms associated with injury
- A “nerve” is just a bundle of axons, just because a specific nerve is damaged doesn’t mean all the axons are non-functional


Sensory and motor components of spinal nerve

Dorsal and ventral rami of spinal nerve contain both sensory and motor fibers from dorsal root and ventral root, respectively.




Joint movements tested for intact innervation of myotome:

- C5: Abduction of arm at glenohumeral joint (axillary and suprascapular nn.)
- C6: Flexion of forearm at elbow (musculocutaneous n.)
- C7: Extension of forearm at elbow (radial n.)
- C8: Flexion of fingers (median and ulnar nn.)
- T1: Adduction and abduction of fingers (ulnar n.)




Dermatomes

- C5: Lateral arm and lateral forearm
 - C6: Thumb and lateral forearm
 - C7: Middle or index finger
 - C8: Little finger
 - T1: Medial elbow
- Black dots represent areas where dermatome overlap is minimal.




College of
Osteopathic
Medicine



(Gilroy et al.) Atlas of Anatomy 2nd ed., Fig. 24.11C

Overview of the brachial plexus.



Musculocutaneous nerve:

C5, C6, C7


Anterior compartment of arm

All: coracobrachialis

biceps brachii

brachialis

Musculocutaneous nerve



Ulnar nerve:

C8, T1

Anterior compartment of forearm

flexor carpi ulnaris

ulnar half of flexor digitorum

profundus

Anterior compartment of hand


all except: muscles of the

Thenar eminence and radial

2 lumbricals

Musculocutaneous nerve

Ulnar nerve



Median nerve:

C6, C7, C8, T1

Anterior compartment of forearm

all except: flexor carpi ulnaris
and ulnar two digitations of
flexor digitorum

Anterior compartment of hand

Muscles of the thenar eminence

Opponens pollicis

Abductor pollicis brevis


Flexor pollicis brevis

Radial two lumbricals


Musculocutaneous nerve

Median nerve

Ulnar nerve





Axillary nerve:
C5, C6
teres minor, deltoid





Radial nerve:
C5, C6, C7, C8, T1
Posterior compartments
All: arm
All: forearm

Brachial plexus







Branches

Cords

Divisions

Trunks

Roots

C5

C6

C7

C8

T
1


Branches

Cords

Divisions

Trunks

Roots




Branches

Cords

Divisions

Trunks

Roots



Branches

Cords

Divisions

Trunks

Roots

C5


C6

C7

C8

T

1




Branches

Cords

Divisions

Trunks

Roots




Branches

Cords

Divisions

Trunks

Roots



Branches

Cords

Divisions

Trunks

Roots

C5


C6

C7

C8

T

1



Branches

Cords

Divisions

Trunks

Roots

C5


C6

C7

C8

T

1



Branches

Cords

Divisions

Trunks

Roots

C5


C6

C7

C8

T

1



Branches

Cords

Divisions

Trunks

Roots

C5


C6

C7

C8

T

1




Branches

Cords

Divisions

Trunks

Roots




Branches

Cords


Divisions

Trunks


Roots



Lesion of C5 root



Lesion of C5 root




Complete loss of function:

dorsal scapular nerve
subclavian nerve

Partial loss of function:

long thoracic nerve
suprascapular nerve
lateral pectoral nerve
subscapular nerves
musculocutaneous nerve
axillary nerve
radial nerve

Lesion of inferior trunk




Complete loss of function:

medial pectoral nerve
 medial cutaneous nerve of arm
 medial cutaneous nerve of forearm
 ulnar nerve

Partial loss of function:

thoracodorsal nerve
 median nerve
 radial nerve

Lesion of ulnar nerve



Complete loss of function:

ulnar nerve

Partial loss of function:



Brachial Plexus: Lesions and Neuropathies

Principles and Practices of Osteopathic Medicine I

2020

Matthew Mihlbachler, Ph.D.

Associate Professor, Anatomy

mmihlbac@nyit.edu


Part 2: Lesions of brachial plexus trunks – Erb-Duchenne palsy and Klumpke's palsy


Injuries to the superior trunk of the brachial plexus – Erb-Duchenne palsy

Wide separation between head and shoulder



Excessive traction on neck during delivery





Erb-Duchenne palsy: Lesion of the superior root

Injuries to the superior trunk (C5-C6) of the brachial plexus (Erb-Duchenne Palsy):


loss of supination of forearm from biceps brachii m.
due to damage to C5-C6 of musculocutaneous n.

Musculocutaneous n.
(C5, C6, C7)

Musculocutaneous n.
innervating anterior arm mm.

Biceps brachii m.:
major supinator of forearm

Radial tuberosity



Forearm in pronation due to
loss of major supinator of forearm


Pronated position

Pronator teres m.



Supinator m.


Pronator quadratus m.

Thumb will be
facing medially
in pronation



Injuries to the superior trunk (C5-C6) of the brachial plexus (Erb-Duchenne Palsy):
Causes loss of musculocutaneous n. function.
Loss of flexion of forearm from anterior arm mm. and sensory deficits to lateral side of forearm






Erb-Duchenne palsy: Lesion of the superior root


Injuries to the superior trunk (C5-C6) of the brachial plexus (Erb-Duchenne Palsy):

Loss of abduction of arm by deltoid m. and supraspinatus m. due to damage to C5-C6 of axillary n. and suprascapular n.

Supraspinatus m.:
1. Loss of abduction of arm
2. Innervated by suprascapular n. (C5-C6)




Deltoid m.:
1. Loss of abduction of arm
2. Innervated by axillary n. (C5-C6)




Injuries to the superior trunk (C5-C6) of the brachial plexus (Erb-Duchenne Palsy):

Loss of lateral rotation of arm by infraspinatus m. and teres minor m. due to damage to C5-C6 of suprascapular n. and axillary n.

Arm is medially rotated due to loss of lateral rotators of arm



Infraspinatus m.:
1. Loss of innervation by suprascapular n. (C5-C6)
2. Loss of lateral rotation of arm




Teres minor m.:
1. Loss of innervation by axillary n. (C5-C6)
2. Loss of lateral rotation of arm

Subscapularis m.:
1. Innervated by subscapular nn. from posterior cord (C5-C6)
2. Arm still medially rotated from pectoralis major m.


Injuries to the superior trunk (C5-C6) of the brachial plexus (Erb-Duchenne Palsy):

Clinical signs of damage to upper trunk:


1. Loss of lateral rotation of arm
2. Loss of flexion of forearm and supination impaired
3. Pronated forearm, medially rotated arm, and extended forearm due to unopposed action of pronators, pectoralis major m., and triceps brachii m., respectively, result in "waiter's tip position".
4. Sensory deficits on lateral aspect of shoulder, arm, forearm, and hand.



Damage to inferior trunk of brachial plexus (C8-T1): Klumpke's Palsy




Abduction of arm during a fall
when grabbing an object



Abduction of arm during delivery







Klumpke's palsy: Lesion of the inferior root

Damage to inferior trunk of brachial plexus (C8-T1): Klumpke's Palsy

Intrinsic hand muscles innervated by C8-T1




Lumbrical actions: Flex MP joint, extend PIP and DIP joints,




Lumbricals: functional



Lumbricals: non-functional




Flexor digitorum profundus



Flexor digitorum superficialis

Damage to inferior trunk of brachial plexus (C8-T1): Klumpke's Palsy


Results in "claw hand" due to loss of action of lumbrical mm. and interossei.



1. Flexion at PIP joints in digits 2-3 occurs due to unopposed action of flexor digitorum superficialis and due to loss of lumbrical and interossei mm. Flexion of DIP joints depends on functionality of lateral side of flexor digitorum profundus m. (C8)

2. PIP of digits 4-5 flexed but DIP not flexed because of unopposed action of FDS but lack of action from the medial side of FDP

Hyperextension of MCP joints occurs due to unopposed action of extensor digitorum m. due to loss of lumbrical mm.



Over time, areas between metacarpals hollows out due to atrophy of lumbrical and interossei mm.

Schreuders et al. (2007), *Physikalische Medizin Rehabilitationsmedizin Kurortmedizin*, 17, 20-27, Fig. 1


Damage to inferior trunk of brachial plexus (C8-T1): Klumpke's Palsy



Yang et al. (2016), *Clinics (Sao Paulo)*, 71(4), 193-198, fig. 2

Damage to inferior trunk of brachial plexus (C8-T1): Klumpke's Palsy


Loss of cutaneous innervation of medial arm and medial forearm



Damage to inferior trunk of brachial plexus (C8-T1): Klumpke's Palsy

Clinical signs of Klumpke's Palsy:

- 1. Loss of all lumbrical and interosseous mm. results in loss of abduction and adduction of fingers; hyperextension of MCP joints due to unopposed action of extensor digitorum m.
- 2. Loss of all lumbrical and interosseous mm. results in flexion of PIP joints in digits 2-5 due to unopposed action of intact flexor digitorum superficialis m. Flexion of DIP joints in digits 1-2 depends on functionality of the lateral side of flexor digitorum profundus mm. and severity of injury.
- 3. Total claw hand results from loss of all intrinsic hand muscles innervated by C8-T1 fibers from ulnar and median nn.
- 4. Radial deviation in wrist flexion from loss of action of flexor carpi ulnaris m.
- 5. Loss of cutaneous innervation of medial arm and medial forearm and medial hand





Brachial Plexus: Lesions and Neuropathies

Principles and Practices of Osteopathic Medicine I


2020

Matthew Mihlbachler, Ph.D.

Associate Professor, Anatomy

mmihlbac@nyit.edu

Part 3: Lesions and neuropathies of long thoracic, axillary, and radial nerves




Lesion of the long thoracic nerve

Damage to long thoracic n. (C5-C7):

Radical mastectomy or knife wound to lateral chest

- 1. Winged scapula (medial border of scapula protrudes posteriorly)
- 2. Inability to abduct arm above horizontal plane due to loss of rotation of scapula





C5-C7 ventral rami


Long thoracic n.

Serratus anterior m.



Medial border of scapula thrust posteriorly when patient pushes hand against wall







Lesion of the axillary nerve

Damage to Axillary n. (C5-C6)

Occurs in quadrangular space from fracture of surgical neck of humerus or inferior dislocation of glenohumeral joint.


- 1. Loss of abduction of arm from deltoid m.
(loss of lateral rotation of arm from teres minor m., but compensated for by infraspinatus m.)
- 2. Loss of cutaneous sensation to proximal lateral arm




Damage to Axillary n. (C5-C6)

Occurs in quadrangular space from fracture of surgical neck of humerus or inferior dislocation of glenohumeral joint.

- 1. Severely reduced ability to abduct arm from deltoid m.
(loss of lateral rotation or arm from teres minor m., but compensated for by infraspinatus m.)
- 2. Loss of cutaneous sensation to upper lateral arm







Neuropathy of the radial nerve




Damage to radial n. in axilla (C5-C8, T1):

Due to poorly fitted crutches or draping arm over back of chair (Saturday night palsy)
Innervation to extensors of posterior arm and forearm
Cutaneous innervation to posterior arm, and forearm, dorsolateral hand, proximal part of dorsal side of lateral 3.5 digits




Damage to radial n. in axilla (C5-C8, T1):

1. Loss of extension at elbow from inaction of triceps brachii m.
2. Loss of extension of wrist from inaction of extensors of hand (posterior forearm) results in unopposed action of long flexors causing wrist-drop
3. Loss of sensation from posterior arm, posterior forearm, dorsolateral hand, and dorsum of proximal part of lateral 3.5 digits




Wrist-drop:
Loss of extension in hand




Fracture of mid-shaft of humerus

Can damage radial n. in
radial groove



Fracture

Damage to radial n. in radial groove due to fracture of mid-shaft of humerus



Innervation to triceps brachii m. not damaged since most motor branches original proximal to radial groove


Posterior brachial cutaneous n.: NOT damaged as it branches from the radial n. proximal to radial groove

Radial n. lesion in radial groove. Contains motor fibers in deep branch to forearm extensors and sensory fibers in superficial branch to dorsolateral hand and lateral 3.5 digits


Posterior antebrachial cutaneous n.: NOT damaged as it branches from the radial n. proximal to radial groove

Damage to radial n. in radial groove (C5-C8, T1):

1. Loss of extension of wrist from inaction of extensors of hand (deep branch of radial n.) results in wrist-drop.
2. Triceps are not paralyzed as they are innervated in axilla (no loss of extension of elbow)
3. Loss of sensation from dorsolateral hand, and dorsum of proximal part of lateral 3.5 digits (superficial branch of radial n.)
4. Posterior arm and posterior forearm are unaffected as these sensory branches come off radial n. proximal to radial groove.



Wrist-drop:
Loss of extension in hand



Loss of sensation from
dorsolateral hand and
proximal part of dorsum
of lateral 3.5 digits only



Brachial Plexus: Lesions and Neuropathies

Principles and Practices of Osteopathic Medicine I


2020

Matthew Mihlbachler, Ph.D.

Associate Professor, Anatomy

mmihlbac@nyit.edu


Part 4: Lesions and neuropathies of median and ulnar nerves



Median nerve injury

Supracondylar fracture:

Can damage median nerve or brachial artery by triceps brachii m. pulling distal tissues posteriorly.
Can be a very serious injury in children.



Loss of motor functions of median n. damage at elbow

Loss of motor innervation by median n. to most flexors and all pronators of anterior forearm

Loss of motor innervation by anterior interosseous n. to flexor digitorum profundus m. (lateral part), flexor pollicis longus m., and pronator quadratus m.

Loss of motor innervation to thenar eminence mm. from recurrent branch of median n. Thenar mm. will atrophy, loss of opposition of thumb. Loss of action of lateral two lumbrical mm. from digital branches.

Flexor pollicis longus m.

Pronator quadratus m.

Median n.

Pronator teres m.

Flexor carpi radialis

Palmaris longus

Flexor digitorum superficialis m.

Flexor digitorum profundus m. (lateral half)

Median n.

Result of damage to median n. at elbow when patient attempts to make a fist

Ulnar n. intact:
Flexion of and DIP joints
(and by extension PIP
joints) of digits 4 and 5
still possible due to intact
medial half of flexor
digitorum profundus m.

Loss of innervation to lateral two lumbricals,
lateral half of flexor digitorum profundus m.,
and all of flexor digitorum superficialis m.
prevents flexion of digits 2 and 3

Loss of innervation to
flexor pollicis longus m.
and flexor pollicis brevis
m. prevent flexion
of thumb.
Loss of opposition of
thumb.




“Hand of benediction” Or ape hand
(a) Median nerve injury

Loss of sensory functions of median n. in hand from damage at elbow


Sensory innervation:

- 1. Lateral 3.5 digits (palmar sides) and dorsum of their nail beds
- 2. Lateral 2/3rd of palm of hand



- Loss of opposition of thumb
- Severe impairment of flexion of hand
 - Less impairment in digits 4-5 (ulnar n. intact)
 - Digits 2-3 have diminished ability to flex and extend
 - “Ape hand” “Hand of Benediction” presentation (be careful with these terms...they are used inconsistently)
- Loss of pronation
- Loss of sensation from lateral 2/3 of palm of hand, palmar side of lateral 3 ½ digits, and nail beds of lateral 3 ½ digits

Damage to median n. at wrist:
Laceration by knife or broken glass by accident
or attempted suicide



Loss of digital branches to lateral two lumbrical mm. and sensory loss to palmar side of lateral 3.5 digits and their nail beds and distal lateral 2/3rd of palm

Loss of palmar cutaneous branch of median n. to proximal lateral 2/3rd of palm

Median n.

Lateral two lumbrical mm.

Flexor pollicis brevis

Abductor pollicis brevis

Loss of recurrent branch of median n.to thenar eminence mm.

Summary of consequences of damage to median n. at wrist:

Loss of motor innervation:


1. All three thenar eminence mm. lose innervation by recurrent branch of median n.
 - Loss of opposition and atrophy of thumb muscles (thenar eminence goes flat)
2. Lateral two lumbrical mm.
 - Much less significant deficits since interosseous mm. are innervated by ulnar n. (both flex MCP and extend PIP and DIP joints of digits 2 and 3).

Loss of sensory innervation same as damage to elbow:

1. Lateral 3.5 digits (palmar side) and dorsum of their nail beds
2. Lateral 2/3rd of palm of hand




"Hand of benediction" Or ape hand
(a) Median nerve injury



Ulnar nerve injury

Neuropathy of ulnar n. at elbow


“Cubital tunnel syndrome”



Lesion of ulnar n. at elbow:

-Loss of action of all interosseous mm. and medial two lumbrical mm.

Loss of adduction-abduction
by interosseous mm:
Patient can not grip a piece of
paper between fingers



Lesion of ulnar n. at elbow:


- Results partial claw hand at rest due to loss of action of medial two lumbrical mm. and all interosseous mm.
- Impaired flexion of MCP joints of digits 4-5, joints are extended
- PIP joints of digits 4-5 flexed because of unopposed action of flexor digitorum superficialis mm.
- DIP joints of digits 4-5 not flexed because of loss of action of the lateral side of flexor digitorum profundus m.
- "Ulnar Paradox", even though the lesion is higher, and thus "more severe", the "claw effect" is less than a lower lesion (see following slides)

Hyperextension of MCP joints of digits 4-5 occurs due to unopposed action of extensor digitorum m. due to loss of interossei mm. and medial two lumbrical mm.

Flexion of PIP joints

DIP joints not flexed

Digits 2-3 not affected




Lesion of ulnar n. at elbow:

Loss of cutaneous innervation to medial 1/3rd of hand and medial 1.5 fingers


Loss of sensation from digital (superficial) branches of ulnar n.:
Distal 1/3rd of medial palm and medial 1.5 digits.

Loss of sensation from palmar branch of ulnar n., from forearm:
Proximal 1/3rd of medial palm



Palmar view


Loss of sensation from dorsal cutaneous branch of ulnar n., from forearm:
Medial 1/3rd of dorsum of hand and medial 1.5 digits



Dorsal view


Neuropathy of ulnar n. at wrist:

- Guyon's Canal (ulnar tunnel) syndrome
- Motor loss in hand same as lesion at elbow



Neuropathy of ulnar n. at wrist:

Damage to ulnar n. at wrist leaves flexor carpi ulnaris m. and medial half of flexor digitorum profundus m. intact.
Wrist flexion NOT affected



Summary of motor damage to ulnar n. at wrist:

1. Paralysis of most intrinsic mm. of the hand
 - Exception-thenar eminence mm. and lateral two lumbrical mm.
 - Opposition of thumb intact
2. Consequences of paralysis:
 - Loss of adduction-abduction of fingers due to loss of interosseous mm.
 - Impairment of flexion of MCP joints and extension of PIP and DIP joints of digits 4-5 due to loss of interosseous mm. and medial two lumbrical mm.
3. Not affected:
 - Actions of flexor carpi ulnaris m. and all of flexor digitorum profundus m. intact due to ulnar n. innervation occurring above lesion at wrist.

Digits 2-3 not affected



Flexion of PIP and DIP joints of digits 4-5 due to loss of medial two lumbrical mm. and interosseous mm. and unopposed action of FDP and FDS


Hyperextension of MCP joints of digits 4-5 occurs due to unopposed action of extensor digitorum m. due to loss of interossei mm. and medial two lumbrical mm.

Lesion of ulnar n. at wrist:

Loss of cutaneous innervation to distal medial 1/3rd of hand and medial 1.5 fingers


Loss of sensation from digital (superficial) branches of ulnar n.:
Distal 1/3rd of medial palm and medial 1.5 digits.

NO LOSS of sensation from palmar branch of ulnar n., from forearm:
Proximal 1/3rd of medial palm



Palmar view

NO LOSS of sensation from dorsal cutaneous branch of ulnar n., from forearm:
Medial 1/3rd of dorsum of hand and medial 1.5 digits



Dorsal view

Summary of different claw effects from ulnar n. lesions at different levels

1. Klumpke's Palsy (damage to inferior trunk of brachial plexus)


- All four MCP joints extended due to loss of all intrinsic muscles of the hand from C8-T1 fibers from median and ulnar nn. and unopposed action of extensor digitorum m.
- PIP joints of digits 2-3 flexed due to unopposed action of flexor digitorum superficialis, DIP joint 2-3 flexion depends on functionality of lateral half of flexor digitorum profundus and severity of injury.
- PIP of digits 4-5 flexed because of unopposed action of FDS, DIP not flexed because no action from medial side of FDP.
- Result: total claw hand, digits 2-3 possibly clawed worse.

2. High lesion of ulnar n. (i.e. at elbow region, broken humerus or cubital tunnel syndrome)

- MCP of digits 4-5 extended because of loss of interosseous mm. and medial two lumbrical mm.
- MCP of digits 2-3 not extended because lateral two lumbrical mm. are still intact.
- PIP and DIP of digits 2-3 not flexed because lateral lumbrical mm. are still intact.
- PIP and DIP of digits 4-5 same as above. PIP flexed, DIP not flexed.
- Result: "ulnar paradox", higher lesion (=more serious) than wrist lesion, but claw effect is lessened due to lack of action from lateral side of FDP. Claw effect will worsen as nerve heals and action of FDP is restored.

3. Low lesion of ulnar n. (i.e. at wrist, Guyon's canal (ulnar tunnel) entrapment)

- Same as above, except both PIP and DIP of digits 4-5 are flexed because of unopposed action of FDP.
- Result: Ulnar claw





Brachial Plexus: Lesions and Neuropathies

Principles and Practices of Osteopathic Medicine I

2020

Matthew Mihlbachler, Ph.D.

Associate Professor, Anatomy

mmihlbac@nyit.edu

Part 5: example cases, questions and summary

A 21-year-old male varsity collegiate wrestler reported left elbow pain and loss of strength and motor control in his left upper limb, most notably in his wrist and hand. He also reports tingling sensations on digit 5 and the medial half of digit 4.

He did not recall experiencing a specific injury that caused the problem. However, he complains that the symptoms worsen after wrestling practice. Point tenderness was found along the medial midshaft of the humerus, the medial epicondyle, and the olecranon process.

Q1.

Which injury is most likely?

- a. cubital tunnel syndrome
- b. Erb-Duchenne palsy
- c. radial neuropathy within the axilla
- d. carpal tunnel syndrome
- e. axillary nerve lesion

Q2.

Which motor function do you expect the patient to mostly likely have difficulty with?

- a. extension of digits 1 and 2
- b. extension of the wrist
- c. abduction of the arm
- d. pronation
- e. adduction of the fingers

A 21-year-old male varsity collegiate wrestler reported left elbow pain and loss of strength and motor control in his left upper limb, most notably in his wrist and hand. He also reports tingling sensations on digit 5 and the medial half of digit 4.

He did not recall experiencing a specific injury that caused the problem. However, he complains that the symptoms worsen after wrestling practice. Point tenderness was found along the medial midshaft of the humerus, the medial epicondyle, and the olecranon process.

Q1.

Which injury is most likely?

- a. *cubital tunnel syndrome
- b. Erb-Duchenne palsy
- c. radial neuropathy within the axilla
- d. carpal tunnel syndrome
- e. axillary nerve lesion in the quadrangular space

Q2.

Which motor function do you expect the patient to have difficulty with?

- a. extension of digits 1 and 2
- b. extension of the wrist
- c. abduction of the arm
- d. pronation
- e. *adduction of the fingers

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1319819/pdf/jathtrain00014-0057.pdf>

Cubital Tunnel Syndrome in a Collegiate Wrestler: A Case Report

[Scott L. Bruce](#), MS, ATC

Scott L. Bruce is Head Athletic Trainer at Slippery Rock University, Slippery Rock, PA.

[Noah Wasielewski](#), ATC

Noah Wasielewski is Graduate Assistant Athletic Trainer at Auburn University, Auburn, AL.

[Richard L. Hawke](#), ATC

Richard L. Hawke is Graduate Assistant Athletic Trainer at University of Virginia, Charlottesville, VA.




Fig 1. The athlete's arm before surgery. (A) ulnar nerve; (B) ulnar collateral ligament; (C) flexor carpi ulnaris muscle; (D) flexor carpi ulnaris tendon.




Fig 2. The athlete's arm after surgery. (A) transposed ulnar nerve; (B) ulnar collateral ligament; (C) flexor carpi ulnaris muscle; (D) flexor carpi ulnaris tendon.

A 25 day-old neonate is presented to the ED. The parents are concerned about the lack of movement in the right upper limb.

A few days after birth, the patient underwent a right postero-lateral thoracotomy to repair an esophageal birth defect. At presentation, the neonate's upper right limb maintains an adducted and pronated posture. Shoulder abduction, supination, and flexion of the elbow were not observed; however, the patient is capable of palmar grasp and distal finger movements.



Q1

A lesion of which nerve best explains this presentation?

- a. inferior trunk of brachial plexus
- b. median nerve
- c. long thoracic nerve
- d. superior trunk of brachial plexus
- e. ulnar nerve

Q2

Based on the neonate's presentation, where might you expect to find a sensory deficiency?

- a. medial side of forearm and hand
- b. lateral side of shoulder
- c. lateral side of arm, forearm and hand
- d. 5th digit and medial half of 4th digit
- e. axilla

A 25 day-old neonate is presented to the ED. The parents are concerned about the lack of movement in the right upper limb.

A few days after birth, the patient underwent a right postero-lateral thoracotomy to repair an esophageal birth defect. At presentation, the neonate's upper right limb maintains an adducted and pronated posture. Shoulder abduction, supination, and flexion of the elbow were not observed; however, the patient is capable of palmar grasp and distal finger movements.



Q1

A lesion of which nerve best explains this presentation?

- a. inferior trunk of brachial plexus
- b. median nerve
- c. long thoracic nerve
- d. *superior trunk of brachial plexus
- e. ulnar nerve

Q2

Based on the neonate's presentation, where might you expect to find a sensory deficiency?

- a. medial side of forearm and hand
- b. lateral side of shoulder
- c. *lateral side of arm, forearm and hand
- d. 5th digit and medial half of 4th digit
- e. axilla

An Unusual Case of Erb's Palsy in a Neonate: A Case Report

Journal of Child Neurology
25(10) 1261-1264
© The Author(s) 2010
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/0883073809356809
<http://jcn.sagepub.com>


Deepak Louis, MD,¹ Venkateshan Sundaram, DM,¹ and
Ram Samujh, MCh, FRCS²

Abstract


Erb's palsy occurs in neonates following traumatic delivery, where excessive traction on the neck stretches these nerve roots. Nonobstetric causes of Erb's palsy are rare in neonates. The authors report the presentation of a female neonate with Erb's palsy following a postero-lateral thoracotomy. The infant underwent surgery on day 3 of life for esophageal atresia and presented with right upper limb weakness on day 21 of life. She demonstrated features of Erb's palsy with normal higher mental functions. An electromyography and nerve conduction study confirmed Erb's palsy. The surgical procedure in the index case did not involve the brachial plexus. However, her right upper limb was positioned hyperabducted during the intraoperative period, which possibly had led to the palsy. The key message of this report is that prolonged stretching of the brachial plexus roots during surgery of the neck and thorax can be an important nonobstetric cause of Erb's palsy in neonates

Keywords

Erb's palsy, thoracotomy, hyperabduction, brachial plexus

Received October 13, 2009. Accepted for publication November 9, 2009.

<https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.1008.4727&rep=rep1&type=pdf>



Left deltoid muscle atrophy


A 43 year-old man presents with conspicuous atrophy of his left deltoid muscle. MRI reveals atrophy of both the deltoid and teres minor. No other notable muscular atrophy was found. He says his deformity is a result of shoulder injury sustained in a car accident 25 years ago. Although he reports some upper limb weakness, he has learned to compensate for his injury and is able to function without any major disabilities.

Q1. Which nerve was injured in the car accident?

- a. long thoracic
- b. superior trunk
- c. median
- d. axillary
- e. suprascapular

Q2. What muscle enables the patient to abduct his left arm?

- a. triceps
- b. supraspinatus
- c. subscapularis
- d. coracobrachialis
- e. rhomboid major



Left deltoid muscle atrophy

A 43 year-old man presents with conspicuous muscular atrophy on his left shoulder. MRI reveals atrophy of both the deltoid and teres minor. No other notable muscular atrophy was discovered. He has diminished cutaneous sensation on the lateral side of his left shoulder. He says his deformity is a result of shoulder injury sustained in a car accident 25 years ago. Although he reports some upper limb weakness, he has learned to compensate for his injury and feels that he is able to function without any major disability.

Q1. Which nerve was injured in the car accident?

- a. long thoracic
- b. superior trunk
- c. median
- d. *axillary
- e. suprascapular

Q2. What muscle enables the patient to abduct his left arm?

- a. triceps
- b. *supraspinatus
- c. subscapularis
- d. coracobrachialis
- e. rhomboid major

Outcomes Following Closed Axillary Nerve Injury: A Case Report and Review of the Literature

CPT Joseph W. Galvin, MC USA; LTC Josef K. Eichinger, MC USA

ABSTRACT We report a case of a 43-year-old male who sustained an axillary nerve injury secondary to a glenohumeral joint dislocation at a young age, and who has served over 20 years in the military with near normal shoulder function. In addition, we review the literature for the natural history of axillary nerve injury. A 43-year-old male sustained a left anterior glenohumeral dislocation in a motor vehicle accident as an 18-year-old. Following prompt manual reduction and subsequent physical therapy, the patient developed a permanent axillary nerve palsy. Despite the development of complete atrophy of his deltoid musculature and persistent sensory loss in the axillary nerve distribution, he experienced restoration of function with minimal to no deficit. Ultimately, he enlisted in the military 4 years after the injury and has served 22 years, which includes combat deployments with normal shoulder function and absence of pain. Axillary nerve injury is a relatively common injury after anterior glenohumeral joint dislocation. There is little known about the long-term outcome of patient's with permanent axillary nerve injury. This case suggests that it is possible for a young athletic individual to function at a high level of activity after permanent loss of axillary nerve function.




FIGURE 2. (A–C) Pictures demonstrating full active forward elevation, external rotation, and internal rotation. (D and E) Pictures demonstrating performance of a push-up without difficulty.

<https://pubmed.ncbi.nlm.nih.gov/26926757/>

Nerve Injuries of the Upper Limb

- Injury to spinal nerves: dermatomes and myotomes
- Injury to proximal brachial plexus
 - Superior trunk
 - Inferior trunk
- Injury to the long thoracic nerve (nerve to serratus anterior)
- Injury to the axillary nerve
- Injury to the radial nerve
 - Different effects depending on location of injury
- Injury to median nerve
 - Different effects depending on location of injury
- Injury to ulnar nerve
 - Different effects depending on location of injury

Lecture Feedback Form:

<https://comresearchdata.nyit.edu/redcap/surveys/?s=HRCY448FWYXREL4R>