

Session Objectives


1. Contrast the major divisions of the hair complex and its relation to neighboring tissue.
2. Explain the development and life cycle of the hair complex.
3. Contrast the major divisions of the nail complex and their relation to neighboring tissue.
4. Explain the development of nails.
5. Describe the regional distribution of sweat and sebaceous glands, contrast their functions, and relate these functions to the cellular composition.
6. Locate and describe the variety of sensory receptors throughout the skin.



Appendages of the skin

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



▲ Schematic of skin and its appendages that shows the epidermis, dermis, and subcutaneous tissue.

Source: “Integument” in Netter’s Essential Histology (2013)

Epidermis is derived from ectoderm during embryonic development. Cells of ectoderm (forming epidermis) grow downward into the dermis and give rise to:

- Hair follicles
- Fingernails
- Toenails
- Sweat glands
- Sebaceous glands
- Mammary glands



Hence, “epidermal appendages”.

Planting the seeds (hair)

1. Develop in 3rd month of fetal life.
2. Epidermis sends down-growth into dermis.
3. Down-growth becomes hair follicle. Invading epidermis forms a cap over the connective tissue papilla.
4. As development proceeds a mature adult follicle is formed and can continuously grow via keratinization.


*similar developmental program as epidemis



Source: WebMD

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine




Three major segments of the hair follicle

Infundibulum - from surface opening of follicle to opening of its sebaceous gland

Isthmus - from infundibulum to level of insertion of the arrector pili muscle.


Inferior segment - growing follicle portion; at base expands to bulb invaginated by vascularized loose connective tissue called dermal papilla.




Follicular development

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




- Follicular bulge houses epidermal stem cells which give rise to sebaceous gland and the hair follicle.
- Deepest downgrowth becomes germinal matrix of hair, includes keratinocytes and melanocytes
- Gives rise to the hair shaft and internal root sheath



Follicular development


NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



- Keratinization occurs after cells leave germinal matrix in region known as keratogenous zone
- When hair emerges from follicle it is entirely keratinized as hard keratin (analogous to entering stratum corneum), does not desquamate.
- Internal root sheath is soft keratin, is broken down/desquamates at opening of sebaceous gland


Multiple layers of the hair shaft



1. Medulla – central, soft keratin, large vacuolated cells; only present in thick/terminal hair
2. Cortex – peripheral to medulla, cells differentiate into keratin-filled cells; contains pigment, 80% mass
3. Cuticle of hair shaft – outermost layer of hair composed of very thin scale-like cells



Multiple layers of the internal root sheath



Need not remember these divisions


4. IRS cuticle: squamous cells

5. Huxley's layer: flattened cells


6. Henley's layer: cuboidal cells in direct contact with outermost part of hair follicle, the external root sheath

The internal root sheath is soft keratin and is broken down at infundibulum level where sebaceous secretions enter follicle


The external root sheath is epidermis and its basement membrane is called the glassy membrane



M - medulla
Cx - cortex
Cu - cuticle
E - epidermis



H - hair
IRS - internal root sheath
ERS - external root sheath
GM - glassy membrane




HP - hair papilla
GC - germinal cells
Me - melanocytes
GM - connective tissues



Vellus versus terminal hair

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Vellus - thin fiber, follicle housed in dermis, distributed over the body except portion of external genitals, palms, and soles

Terminal - thick fiber, follicle housed in hypodermis, distributed to head, eyebrows, end of eyelids, genitals, axilla and other regions impacted by androgens during puberty

Arrector Pili (muscles for hair)


Bundle of smooth muscle fibers that attaches to the connective tissue sheath of a hair follicle

When muscle contracts, pulls hair follicle outwards and makes hair stand up; hairs exit at an angle.

Produces “goose bumps”/dimpling of skin

Also squeezes on sebaceous glands causes the secretions to be emptied

Activated by autonomic nervous system.




Source: “Integument” in Netter’s Essential Histology (2013)

Planting the seeds (Nails)


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Hair



Nail




3rd month of embryonic life, epidermis of fingers and toes “invade” underlying dermis along a curved line to form the nail

Proximal part of nail, the nail root, is buried in fold of epidermis covering cells of the germinative zone.

Cells become fully keratinized.

Source: “Integumentary System”
in Ross and Pawlina
Histology, 7th Ed. (2015)




Matrix of Nails


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Hair



Nail




Matrix of nails has stem cells, melanocytes, Merkel's and Langerhans' cells

The stem cells divide, migrate toward the root and differentiate and produce keratin of the nail which is hard keratin, does not desquamate – consists of densely packed keratin filaments embedded in a matrix of amorphous keratin with a high sulfur content, which makes nails hard.

Crescent-shaped white area near nail root is the lunula: color is from thick, opaque layer of partially keratinized matrix cells

For the nail,
but not of
the nail



NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine

Eponychium (cuticle) – edge of skin fold covering nail root, also hard keratin, does not desquamate

Hyponychium – epidermis underlying distal tip of nail where stratum corneum of distal free edge contacts the nail; thickened epidermal layer which secures the free edge of the nail plate at fingertip

Source: "Skin and its appendages"
in Gray's Anatomy (2010)

Planting the seeds (sebaceous glands)






Cells of external root sheath in ~ upper 1/3 of follicle grow out into adjacent dermis and differentiate into sebaceous glands

Sebaceous cells develop a product that kills the cell.

See a theme?



Eccrine gland



ANK INSTITUTE
IOLOGY

osteopathic

Simple tubular gland with a secretory portion and excretory duct. Secretory part housed in deep dermis or subcutaneous tissue. Excretory duct goes directly to skin surface.


Reabsorption of some Na and water in duct results in release of hypotonic sweat low in protein with different amounts of NaCl, urea, uric acid, and ammonia



Major role in temperature regulation by evaporation of water from sweat on body surface

Source: "Integumentary System" in Ross and Pawlina Histology, 7th Ed. (2015)

Apocrine glands



Develop from epidermal downgrowths for hair follicles;
connection to follicle remains
Coiled with secretory portion deep in dermis or hypodermis

Lumens of secretory portion are much wider than those of
eccrine glands

Apocrine glands store secretory products in their lumens,
eccrine glands do not.

Also, reabsorption does not take place in duct, unlike eccrine
glands

Source: "Integumentary System" in Ross and Pawlina Histology, 7th Ed. (2015)

Cells of the apocrine glands

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Originally thought to be apocrine secretion (membrane budding)
– actually merocrine.

Only one secretory cell type – eosinophilic with bleblike apical protrusions

Myoepithelial cells are also present in secretory portion (same as eccrine)

Produce secretion with protein, carbohydrate, ammonia, lipid, and some organic compounds that may give milky color
Secretion develops odor via bacterial action on skin,

Become functional at puberty – linked to pheromones?


Source: “Integumentary System” in Ross and Pawlina Histology, 7th Ed. (2015)



Free nerve endings

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Present in both throughout epidermis and dermis

Lack a connective tissue or Schwann cell investment, i.e. “free”

Heat, cold, itch, and pain

Surround most hair follicles
mechanoreceptors (e.g. “whiskers”)



Summary

- Epidermal appendages include hair, nails, sebaceous glands, and sweat glands.
- Develop from epidermal downgrowth and are housed deep in skin.
- Hair, nails, and sebaceous glands express similar developmental sequence as epidermis.
- Eccrine and apocrine sweat glands differ in many ways.
- The skin has an extensive nerve supply.

