

Learning Objectives:

1. Basic Newborn Visit Interviewing/History Taking
2. Newborn Growth
3. Newborn Development
4. Vaccine Schedule
5. How to examine a newborn
6. Physical exam pathologies
7. Children – toddlers to 5 years of age history questions
8. Early AAP Screenings
9. Tips and Tricks to examine young children
10. Resources

At delivery -

A P G A R score (at 1 min and 5 mins of life)

At initial newborn visit -

Always ask birth history:

Gestational age

Birth weight if they have it

Complications during delivery?




Did baby need NICU care/oxygen/phototherapy?

Mode of delivery? And why?

GBS status of mother or if mom had any diseases

Passed hearing, CCHD, got hep B, erythromycin, vit K? Blood type /coombs status

Target Oxygen Saturation Table	
2 minutes	65%-70%
3 minutes	70%-75%
4 minutes	75%-80%
5 minutes	80%-85%
10 minutes	85%-95%

The Apgar Score				
Indicator	0 Points	1 Points	2 Points	Total Points
A Appearance (skin color)	Blue (cyanotic) Pale (white) all over 	Pink body blue extremities (Acrocyanosis) 	Pink body 	—
P Pulse	Absent No pulse 	< 100 bpm 	> 100 bpm 	—
G Grimace Reflex Irritability	Floppy No response 	Minimal Response when Stimulated 	Prompt Response Sneezing, Crying or Pulling Away when stimulated 	—
A Activity (Muscle tone)	No Movement Flaccid 	Flexed arms and legs 	Active motion 	—
R Respiration	Absent No breathing 	Weak or Slow Irregular 	Vigorous Strong Cry 	—
Severely depressed Moderately depressed Excellent condition				0 - 3 4 - 6 7 - 10

Approach to the normal newborn

Weight: should be gaining 20-30g/day - should double weight by 4 months. Triple by 1 year
Normal FT Babies have 14 days to get back to their birth weight, want NO MORE THAN 10% weight loss

Asymmetric vs symmetric SGA, LGA

HC: Birth 35cm: growth fastest during newborn period - 2cm/month for the 3 months, then 1 cm/month for next 3 months - slows down to 0.5cm/month for 6 months. 1 year - fontanelles fuse


During this time: also important to check newborn screening results - if normal - can take up to 2 weeks to come back

If abnormal: usually clinic gets faxed important and emergent - might need redraw or intervention

Ask about feeding what and how much and how often, ask about elimination and urination, always ask about Safe Sleep and how mom and dad are doing. Use Edinburgh Postnatal depression scale

Happy mommy = Happy baby

SIDS Rate and Back Sleeping (1988 – 2006)



SIDS Rate Source: CDC, National Center for Health Statistics,
Sleep Position Data: NICHD, National Infant Sleep Position Study.



Approach to the normal newborn

Contraindications to Breast Feeding

Baby reasons: galactosemia, maple syrup urine disease, PKU

Parent reasons: HIV ½ (depending on treatment and viral load)

Active TB

HSV breast lesions

Ebola

Certain medications

Radioactive compounds

Breast feeding/formula exclusively until 4 months

4-6 months can start some purees/solids (one food at a time, early peanut exposure)

By age 1 can start cow's milk (max 24oz/day)

Stool colors you don't want: RED/WHITE/BLACK
all others are okay

Urine: urate crystals common

TABLE 2

Resources for Prescribing Medications During Lactation

For prescribers

LactMed

Free, peer reviewed, authoritative, and regularly updated e-Lactancia

Free, available in English and Spanish, peer reviewed, and regularly updated

Medications and Mothers' Milk

Authoritative textbook written by Thomas Hale, website and smartphone application require a subscription

For patients


Drugs.com

Free general pharmaceutical database, includes well researched lactation information






Mothertobaby.org

Free information on medications in pregnancy and lactation, available in English and Spanish

Information from references 2 and 4-7.



Growth charts



Newborn Exam

Start with Head - fontanelles, shape of head


Eyes: RR reflex, spacing of eyes, strabismus, tracking by 2-4 months

Ears: pits, tags, low set vs not

Nose: Patent nares

Mouth: outer Deformities, check suck reflex and hard palate intact

Clavicles (newborn): for deformities/step offs



Types of scalp findings from delivery in newborn

Caput succedaneum – like a CAP swelling is throughout, crosses suture lines, usually prolonged delivery – due to birth canal or vacuum assisted. Resolves 24-48 hours

Cephalohematoma – DOES not cross suture lines, often with bruising, skin discoloration, Resolves 2-4 weeks, +can cause jaundice



Subgaleal hemorrhage: dangerous, crosses sutures, Can accumulate a lot of blood causing anemia, Hypovolemia, +positive fluid wave

Caput succedaneum V/S Cephalohematoma



Dr. Rekha Thaddanee

Extracranial Hemorrhage or Edema in Newborn



Newborn Exam

Normal Red Reflex:



Abnormal Red Reflex:



Etiologies: congenital cataracts, retinoblastoma, chorioretinal coloboma, scarring from retinopathy of prematurity

TABLE 99.1

Visual System Milestones

From Edward DP, Kaufman LM. Anatomy, development, and physiology of the visual system. 2003;50:1–23.

Description	Age
Pupillary light reaction present	30 weeks' gestation
Pupillary light reaction well developed	1 month
Lid closure in response to bright light	30 weeks' gestation
Blink response to visual threat	2–5 months
Visual fixation present	Birth
Fixation well developed	2 months
Visual following well developed	3 months
Accommodation well developed	4 months
Visual evoked potential acuity at adult level	6 months
Grating acuity preferential looking at adult level	2 years
Snellen letter acuity at adult level	2 years
Color vision present	2 months
Color vision at adult level	6 months
Stereopsis developed	6 months
Stereoacuity at adult level	7 years
End of critical period for monocular visual deprivation	10 years

Newborn Exam

Neonatal conjunctivitis:

Incidence 1-21% depending on region

Gonococcal Infections: Copious purulent discharge (EARLY) can start 2-4 days of birth. Tx: IM dose of 3rd generation cephalosporin

Chlamydia Infections: Watery discharge (LATER) will start 5-14 days from birth. Tx: oral azithromycin (or erythromycin), not preventable with erythromycin ointment at birth

HSV: can cause infectious keratitis, scarring of the cornea

Swabs helpful!

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



FIG. 20.57

Ophthalmia neonatorum, a hyperacute bacterial conjunctivitis, with thick purulent discharge and red swollen lids.



Newborn Exam

Upper Extremities: tone, strength, 5 fingers on each hand, creases, grasp, moro (disappears by about 2-4 months)

Chest: deformities, heart and lung sounds

Abdomen: distension, soft, masses, HSM, hernias


GU: femoral pulses bilaterally, ambiguous genitalia, males descended testes, anus patent!

Lower Extremities: Hips - Ortolani & Bartlow, leg length, tone, strength, deformities, 5 toes, plantar reflex

Back: Spine deformities, tufts of hair, sacral dimples

Neuro: reflexes

Skin: rashes, jaundice, etc



Newborn Exam

Prominent xyphoid



Chest


Pectus excavatum



Supernumerary nipple

Pulses

120-160



Femoral pulses very important! if abnormal = coarctation of aorta


Abdomen



Check for hepatosplenomegaly and abdominal masses/hernias



Umbilical cord care




Falls off on it's own in 10-14 days – cannot submerge in water til then

Delayed separation: immune disorder

DO NOT cover

If bleeding or foul smelling discharge, alert PCP

Umbilical hernia – nothing to do until age 3-6 years, as long as it's compressible




[Enlarge](#) [Print](#)

Selected Differential Diagnosis of Neonatal Umbilical Masses

Condition	Characteristics
Umbilical pyogenic granuloma	Small, red, moist, velvety mass
Umbilical polyp	Firm, red mass; does not resolve after application of silver nitrate
Umbilical hernia	Covered with skin, usually reducible
Patent urachus	Discharge (urine) from the umbilicus
Omphalomesenteric duct remnant	Bilious or fecal discharge from umbilicus
Omphalocele	Membranous sac containing viscera herniates through the umbilicus
Gastroschisis	Loops of bowel protrude lateral to an intact umbilicus

Umbilical granuloma



Umbilical hernia

Newborn Exam

Heart:

Fast Rate

Murmurs hard to hear when subtle

Grading (1-6)

Systolic vs Diastolic

Better with positioning?

Cyanosis





Lungs:

Lung sounds


RR 30-60 normal in newborns

Signs of respiratory distress


Accessory muscle use



Hip Exam: checking for Hip Dysplasia



- Risk Factors: Female, Breech, Pos Family history
- **Barlow sign "Back"**
- Maneuver attempts to subluxate or dislocate unstable hip from acetabulum with gentle hip **adduction**
- **Ortolani sign "Out"**
- Maneuver attempts to reduce a subluxated or dislocated femoral head back into acetabulum with gentle hip **abduction**
- Asymmetry
- Asymmetrical gluteal or thigh skinfolds
- Specificity is low; finding is present in up to 25% of unaffected infants
- Younger than 4 months – b/I Hip US
- After 4 months – do bilateral hip xray



Positive Galeazzi Sign ↑



GU exam

Females:

Labia – make sure not fused

Clitoromegaly

Some white vaginal discharge + blood - normal – mom's hormones

Males:

Testicular exam – both descended ** **Cryptorchidism**

Urethra placement – hypospadias vs epispadias

Scrotal sac – make sure not bifid

Hydroceles give you positive transillumination test

In BOTH: Anus patent

If Ambiguous Genitalia in a Newborn : have to rule out classic **Congenital Adrenal Hyperplasia (CAH)

When to suspect

- Micropenis: Stretched penile length < 2.5cm in a term newborn
- Asymmetry of labioscrotal folds
- A penis with B/L nonpalpable testis
- U/L cryptorchidism with hypospadias
- Perineoscrotal or penoscrotal hypospadias with or without microphallus even if testes are descended.
- Apparently female external genitalia with clitoromegaly or inguinal hernia
- Overtly abnormal genitals like cloacal exstrophy
- Discordance of external genitalia with prenatal karyotype.




Imperforated Anus

If a baby is born without an anus (the opening at the end of the digestive tract), then this birth defect is known as an imperforated anus.

Baby is not able to pass stools from the body.


It is an **acute emergency**.

Surgery is required to repair imperforated anus.
Initial surgery to reroute the intestines (colostomy)
Second surgery for colostomy closure.



Skin:

1. Infantile Eczema
2. Hemangioma
3. Cradle Cap (seborrheic dermatitis)
4. Nevus Simplex
5. Erythema toxicum neonatorum (ETN) (Etox)
6. Pustular melanosis
7. Congenital Dermal Melanocytosis
8. Milia



Children - toddlers up to 5

Ask milestone questions: Gross motor, Fine motor, Language, Social Emotional

Screenings: **M-CHAT scoring** at 18 months, 24 months
Developmental screening: 9months, 18 months, 30 months (Ages and Stages Ques) vs surveillance at each well child visit

How is school/daycare going? Changes in social situation?

Safety questioning: guns, smoking, seat belts, car seats/booster seats, helmets while riding bike, smoke detectors, carbon monoxide detectors, poison control info at hand for toddlers

Anticipatory guidance: baby proofing house, locking away meds, cabinet + oven clamps

Hearing and Vision screenings in clinic – older than 3-4 years of age

AAP recommends **Lead and Iron testing at 1 yr plus/minus 2 yr** (pre-mie/WIC 9 months), BP measurement starting at age 3

Toilet Training: usually >18 months when child is ready



Do you know the four stages of car seat safety?



Rear Facing

Rear facing provides the best protection for a young child's spine. Rear face for a minimum of 2 years, ideally until the seat is outgrown.



Forward Facing

When a child has outgrown their rear facing convertible seat and is at least 2 years old, move to a forward facing 5 point harness.



Booster

When a child has outgrown their harnessed seat, is at least 5 years old, and is mature enough to sit properly, move to a belt positioning booster.



Adult Seat Belt

When a child is at least 10 years old and can pass the 5 step test, move to the seat belt alone.