Objectives

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Be able to describe the autonomic processes of the cardiovascular system and how they affect the physiology of a patient and relate to viscerosomatic reflexes
- Be able to describe the lymphatic and musculoskeletal structures related to the cardiovascular system and how changes to these structures may affect the system.
- Be able to identify and describe the location of Chapman's points related to the cardiovascular system and how these can be used to diagnose and treat cardiovascular problems.
- The student will be able to describe the goals for treating a patient with osteopathic principles and be able to apply the 5 models of osteopathic treatment to treating patients with cardiovascular disease.

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

Clinical Relevance:

You will be able to look for somatic dysfunctions in specific areas based on the patient's cardiac disease process and understand how and why to treat the patient's autonomic nervous system, lymphatic system and musculoskeletal system .

Models of Osteopathic Manipulation

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

Biomechanical
(Postural/Musculoskeletal)

Behavioral
(Psychobehavioral)



Neurological

Metabolic-
Energy
(Bioenergy)


Respiratory/Circulatory

Neurological Model of Osteopathic Manipulation

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

Biomechanical
(Postural/Musculoskeletal)

Behavioral
(Psychobeavioral)



Neurological

Metabolic-
Energy
(Bioenergy)

Respiratory/Circulatory




- Increased sympathetic tone → predisposition for ectopic beats and ventricular fibrillation.
- Increased sympathetic tone to the coronary vessels has been linked to coronary vasospasm.

- Prolonged increased sympathetic tone to the heart decreases the heart's ability to form collateral circulation.

- An imbalance between the right and left stellate ganglionic sympathetic impulses to the heart has been postulated as being the mechanism for causing some ventricular arrhythmias in some long Q-T syndromes.

- Vagus exits the skull through the jugular foramen
- Basi-occiput & petrous portion of the temporal bone are borders of the jugular formamen
- Treatment of the O-A is an often-cited method for optimizing the Vagal Supply to the Viscera



3d4 Medical 2020

Respiratory/Circulatory Model

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

Biomechanical
(Postural/Musculoskeletal)

Behavioral
(Psychbehavioral)



Neurological

Metabolic-
Energy
(Bioenergy)

Respiratory/Circulatory (includes lymphatics!)

Cardiovascular Lymphatics (cont')

- The thoracic duct and large lymphatic vessels are under sympathetic control
  - Therefore: increased sympathetics can → **increase pumping of lymph. This effect is theorized to <u>be more to modify immune response rather than increase flow through vessels.</u>**

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Study: “Active Drainage of Cardiac Lymph in Relation to Reduction in Size of Myocardial Infarction: - An Experimental Study”
  - Akira T, Uehara K, Fukuda S, Takenaka K, and Koga M
  - Angiology 1990 Dec; 41(12): 1029-36

(CONTINUED…)

Thoughts:


- Treating the body to maintain optimal lymphatic flow in a patient at risk for a myocardial infarction (MI) may help decrease the size of an MI if one does occur (research needed)
- Or, treating the lymphatics post-infarct may help decrease the area ultimately affected by the infarct. (research needed)

Biomechanical Model of Osteopathic Manipulation

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

Biomechanical (Postural/Musculoskeletal)

Behavioral (Psychobehavioral)



Neurological

Metabolic-Energy (Bioenergy)

Respiratory/Circulatory

Musculoskeletal findings for Cardiac disease

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Viscerosomatic reflexes at T1-5 on left (heart)
- Acute viscerosomatic reflexes
  - Warm, moist, boggy, edematous, + skin drag test, + erthyema test
- Chronic viscerosomatic reflexes
  - firm, dry sponginess, joint is “really stuck” , rubbery end-point
- Chapman’s reflexes at the left second intercostal space (between ribs 2 & 3) near the sternum and posteriorly between the transverse processes (TPs) T2 and T3


Somatic Findings in Hypertension

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Adrenal Function Chapman's Point
  - Anterior: 1 inch lateral & 2 inches superior to umbilicus
  - Posterior: Intertransverse space between T11 & T12
- Kidney Chapman's points
  - Posterior: Intertransverse space between T12 & L1
  - Anterior: 1 inch superior and 1 inch lateral to the umbilicus.
- Viscerosomatic reflexes from the kidneys at T10 and T11

Remember there are fascial relations to the heart

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine




- The fibrous pericardium becomes the mediastinal fascia, and is continuous into the neck, after attaching firmly to the sternum, manubrium, and clavicles.
- In the neck it is called the pretrachial or anterior cervical fascia.
- This attaches firmly to the cranial base.

Treat fascial restrictions in the neck and thorax to optimize mediastinal motion and lymphatic flow

Pectoralis Trigger Point for Arrhythmias


- Pectoralis Major Trigger point on right has been associated with supraventricular tachyarrhythmias.
- Located below the Right Rib 5 margin, half-way between the sternal margin & the nipple line




Kuchera. Osteopathic Considerations in Systemic Dysfunction. P 57

Left sided Pectoralis Major trigger point


NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine




clavicular



sternal





costal

- Trigger Points of the Left Pectoralis Major May (somewhat) Resemble “Referred” Pain Pattern of Heart
- Hallmark of a Trigger Point is that the pain travels from the site of Tenderness in a characteristic pattern
- Reproducible Muscular Pain and characteristic pattern may aid clinical Dx & Tx

Travell, Simons & Amp; Simons’ Myofascial Pain and Dysfunction: The Trigger Point Manual, 3e, 2019, Ch 42

Copyright © Wolters Kluwer

Behavioral Model

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Consider that osteopathic care (medical, manipulation, and counseling) can help
  - Decrease anxiety and stress – treating the patient allows patient to get back to work, interact with family, perform activities of daily living, etc.
  - Encourage healthy lifestyle –proper diet, exercise and sleep behaviors

Metabolic Model

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- One example of how OMT can affect the Metabolic-Energy model is that by treating the sympathetic nervous system will help to decrease peripheral vascular resistance, thereby decreasing the workload of the heart which will, in turn, make total body energy consumption less.

Treatment of Myocardial Infarction


NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Treat T2-3 on left, rib raising (reducing sympathetics.)
- Post-chest compressions- ribs/sternum-indirect
- C2 & Cranial base dysfunction (vagal…parasymp)
  - Use indirect if you can
- Paraspinal inhibition to decrease the total peripheral resistance and cardiac workload (sympathetics)
- Thoracic inlet (rib 1, clavicles, scalenes): (lymphatics, stellate ganglion)
- Hyoid fascial connection to mediastinum (lymphatics)
- Pectoral traction to increase lymphatic return
- Doming of diaphragm (better motion; lymphatics)
- CV4 fluid homeostasis, decrease stress levels (parasympathetics)

Treating the Hypertensive Patient

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- T1-T5 (sympathetics to heart)
- T10-11 (sympathetics to kidneys)
- Treat All sympathetics (rib raising, chain ganglia) to decrease total peripheral resistance
- Chapman's points for the adrenal glands and kidneys.
- OMT should be slow, gentle, augment rest and relaxation, promote autonomic balance and release fascial contractures.
- OA, OM (parasympathetics)
- CV4, Fascia- Lymphatics, body fluids
- Ribs, gentle OMT- ANS, stress reduction/relaxation
- T-L junction/chapman's – ANS, balance rennin-angiotensin



Treating Arrhythmias

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Decrease sympathetic tone
  - Cervical ganglion, rib raising T1-T5
- Modify vagal tone
  - ocular pressure, carotid massage, valsalva maneuver
  - treat occiput, OA-AA, C2
- Tx pectoral trigger point (if present)



Treating Congestive Heart Failure

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Lymphatics- lymphatic pump, effleurage
  - Help to decrease inefficiency of the dilated thoracic duct
  - Restore lymph flow to more physiologic levels
    - Open up a path for the fluid to move, don't force the fluid back and overload an already compromised heart (ie – don't do a pedal pump)
- Treat the diaphragm and thoracic cage
  - Optimal respiratory function is vital for the maintenance of treatment effects directed at lymphatic and venous return.

Summary Slide

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- In summary, when treating any cardiovascular patient osteopathically, in addition to administering all other treatments and counseling according to the standard of care, one must address the sympathetics, parasympathetics, lymphatics and any somatic components related to the other three in order to improve homeostasis and promote physical and mental healing.
  - You are expected to be able to use the principles discussed to osteopathically evaluate and treat a patient with any cardiac disease, not only the ones mentioned.