

Session objectives

- 1) Students will be able to understand the divisions of the medulla, pons, and midbrain from external views and transverse sections.
- 2) Students will be able to identify the major structures, reflecting nuclei and tracts, on the external anatomy of the brainstem, along with cranial nerves and ventricles.
- 3) Students will be able to identify the major nuclei on transverse sections of the brainstem and cerebellum.
- 4) Students will be able to describe the placement of the cranial nerves on the brainstem.

The Human Brain: An introduction to its functional anatomy by J. Nolte
Chapter 11 “Organization of the brainstem”

<http://arktos.nyit.edu/login?url=https://www-clinicalkey-com.arktos.nyit.edu/#!/content/book/3-s2.0-B9780323653985000114>

The brainstem involves many systems & functions


- Cranial nerves and cranial nerve nuclei
- Assorted major nuclei (e.g. substantia nigra)
- Reticular formation
- Ascending & descending pathways (e.g. corticospinal tract)

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



The largest divisions of the brainstem: medulla, pons, and midbrain



**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

First identity global region,
then narrow down to
specifics.

Important differences within
the rostral-caudal extent of
the major divisions


The cerebellum and vesicles
are complementary to the
brainstem.


External views of the tracts & nuclei of the brainstem

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Ask yourself:
“What external features
reflect a structure
underneath?”





Cerebral peduncle – thick columns
Anterior portion = crus cerebri which
includes descending tracts from
cerebral cortex.

Basilar pons (BP) – bulb-shaped

Medulla – spinal cord-like
Medullary pyramids (inside is the corticospinal tract)
Pyramidal decussation (d)
Olive (O) (inside is the inferior olivary nucleus)

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Ventral brainstem

Stained sections of the brainstem

nuclei: light area in myelin-stained slides

axons: dark area in myelin-stained slides

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Anterior-posterior regions

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine

Tectum is the superior and inferior colliculi



Tegmentum is the region surrounding (and largely anterior) to the 4th ventricle & cerebral aqueduct.

It includes cranial nerve nuclei, reticular formation and some assorted tracts.

Basilar region is anterior to the tegmentum and includes major descending tracts.



Medulla



rostral

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Overall tubular shape.
Opens at top to the 4th ventricle.

Would inferior olivary nucleus be dorsal or ventral, rostral or caudal?

What about gracile and cuneate nuclei?
Hint: also called dorsal column nuclei.

caudal

Midbrain

NEW YORK INSTITUTE OF TECHNOLOGY


College of Osteopathic
Medicine

rostral

Tectum is the superior and inferior colliculi

Tegmentum is the region surrounding (and largely anterior) to the cerebral aqueduct and contains cranial nerve nuclei, reticular formation and some assorted tracts


caudal



Where sections align with the external brainstem

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine





Nuclei to identify (remember their function)


Midbrain	red nucleus	motor-related (contralateral)
	superior colliculus	saccadic eye movements
	inferior colliculus	hearing
	substantia nigra	motor-related (contralateral) <i>linked to basal-ganglia</i>
	periaqueductal gray	pain processing
Pons	pontine nuclei	cortical-cerebellar relay (contralateral)
	dentate nuclei	motor coordination (ipsilateral) <i>pons-level cerebellar nucleus</i>
Medulla	gracile & cuneate nuclei	touch, proprioception (ipsilateral)
	inferior olivary nucleus	motor coordination (contralateral)
	spinal trigeminal nucleus	pain, temperature for face (ipsilateral)

*note that light areas indicate nuclei in myelin-stained slides and this will include cranial nerve nuclei

Medulla




Medulla



Pons

NEW YORK INSTITUTE OF TECHNOLOGY





College of Osteopathic
Medicine



dentate nucleus

4th ventricle

pontine nuclei




Pons



4th ventricle

pontine nuclei



NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine


Pons

cerebral
aqueduct/
4th ventricle

pontine nuclei

NEW YORK INSTITUTE
OF TECHNOLOGY





College of Osteopathic
Medicine



Name the nuclei!

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Cranial nerves


NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

Ventral View



Dorsal View



NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine

Nerves are part of the PNS.
Tracts are part of the CNS.

All cranial nerves innervate
ipsilateral tissue, whether they are
afferents, efferents, or both.

Ipsilateral. Ipsilateral. Ipsilateral.

This is not a hard rule for tracts.
They can convey ipsilateral
information, contralateral
information, or both.

CN 3 →

CN 5 →

CN 6 →

CN 7 →

CN 8 →

CN 9 →

CN 10 →

CN 12

CN 11 →

Summary

- The brainstem has distinctive external features for the medulla, pons, and midbrain.
- The tectum is in the dorsal midbrain only. Tegmentum is further ventral and at all divisions. And the basilar division is most anterior/ventral and present at all levels.
- Distinct nuclei are present at each level of the brainstem.
- There is a rational order for numbering the cranial nerves.