

Supraspinal motor systems: Cerebellum

Isaac Kurtzer, Ph.D.
Department of Biomedical Science
ikurtzer@nyit.edu

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Do.
Make.
Heal.
Innovate.
Reinvent the Future.

Session Objectives

1. Identify the lobes, nuclei, and peduncles of the cerebellum along with associated brainstem nuclei.
2. Identify the pathways linking the cerebellar cortex and nuclei to the spinal cord, brainstem, and cerebrum.
3. Describe the basic cerebellar circuit, the input, internal organization, and output.
4. Understand how the functional specialization of different cerebellar regions relate to various abilities along with the origin and expression of cerebellar disorders.

Nolte's The Human Brain: An introduction to its functional anatomy

Chapter 20 Cerebellum

<http://arktos.nyit.edu/login?url=https://www-clinicalkey-com.arktos.nyit.edu/#!/content/book/3-s2.0-B9780323653985000205>

General features of the cerebellum

NEW YORK INSTITUTE
OF TECHNOLOGY



- “Little brain” composed of cortical sheet and nuclei
80% of surface area of cerebral cortex!
80% of neurons in the brain!
- Critical for motor coordination.
- Damage results in poor coordination.
- Does not directly drive motor neurons.
- Less clear relation to cognitive & affective function.




Gordon Holmes studied wounded British soldiers (WW1); the standard helmet did not protect the back of their heads unlike German soldiers.

Holmes GM (1917) The symptoms of acute cerebellar injuries due to gunshot injuries. *Brain* 40:461–535

Impressively understudied




- about 10% of the research on cerebral cortex
- very few studies examine the variety of tasks needed to uncover cognition and emotion



Wang (2025) Ignoring the cerebellum is hindering progress in neuroscience

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




King (2019) Functional boundaries in the cerebellum revealed by multi-domain task battery

Gross cerebellar anatomy

- Lobes and regions - functional divisions of the cortical sheet
- Peduncles - fiber tracts conveying info to and from
- Internal Nuclei – the primary output nuclei of the cerebellum
- Associated external nuclei – input nuclei to the cerebellum

Lobes and regions: superior view



Lobes and regions: inferior view

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine


Flocculonodular lobe (Flocculus)



Vermis

Medial
hemisphere

Lateral
Hemisphere




Posterior Lobe



Lobes and regions: sagittal view

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine



The internal nuclei (deep cerebellar nuclei)

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Relay information from the cerebellar cortex to brainstem and cerebral cortex

The internal nuclei (deep cerebellar nuclei)


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



All output (with one exception) are axons from the deep cerebellar nuclei.


Relay information to the brainstem and cerebral cortex.



External brainstem nuclei linked to cerebellum

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine



Red nucleus (midbrain) – receives output from the contralateral cerebellum

Pontine nuclei (pons) – transmits information from ipsilateral cerebrum to contralateral cerebellum

Vestibular nucleus (medulla) – receives output from ipsilateral cerebellum

Inferior Olive (medulla) - send inputs/afferents to contralateral cerebellum


External brainstem nuclei linked to cerebellum

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Red nucleus (midbrain) –
receives output from the
contralateral cerebellum



Pontine nuclei (pons) – transmits information from
ipsilateral cerebrum to contralateral cerebellum



Vestibular nucleus (medulla) –
receives output from ipsilateral cerebellum




Inferior olive (medulla) - send inputs/afferents to
contralateral cerebellum

Cerebellar peduncles

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




(A from Nolte J: The human brain, ed 5, St. Louis, 2002, Mosby.)

Cerebellum communicates with three major portions of the nervous system


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine


Spinal Cord



Brainstem



Cerebral cortex



Remember that cerebellum controls the ipsilateral body. Hence, signals (to and from) must cross side to communicate with the cerebellum. A mix from the spinal cord and brainstem.

Afferents and efferents of the peduncles

Inferior peduncle

Afferent tracts

Cuneocerebellar(arm)
Dorsal spinocerebellar(leg)
Olivocerebellar
Rostral spinocerebellar(arm)
Reticulocerebellar
Trigeminosensoricerebellar
Vestibulocerebellar

Efferent tracts

Cerebellovestibular
Cerebelloreticular

Middle Peduncle

Afferent tracts

Corticopontocerebellar

Superior Penduncle

Efferent tracts

Cerebellarthalamic
Cerebellorubral
Interpositotectal

Afferent tracts

Ventral spinocerebellar*(leg)
Tectocerebellar

*no need to remember all tracts but recognize the breadth of connections

A reasonable simplification

Inferior peduncle

Afferent tracts

All remaining afferents

Middle Peduncle

Afferent tracts

Corticopontocerebellar

Superior Penduncle

Efferent tracts

Everything

Three functional divisions of the cerebellum

Vestibulocerebellum

Flocculonodular lobe receives input from vestibular nuclei and nerves.

Head-eye coordination and balance.



Spinocerebellum

Vermal and paravermal regions of anterior & posterior lobes receive spinal inputs.

Axial control and gait.




Cerebrocerebellum

Lateral hemispheres of anterior and posterior lobes receive cerebral cortical input via pontine nuclei.


Voluntary movements especially learned and skillful movements

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine




Functional divisions of the cerebellum



How cerebral cortex communicates with ipsilateral thalamus and contralateral cerebellum

**NEW YORK INSTITUTE
OF TECHNOLOGY**


College of Osteopathic
Medicine



How red nucleus and inferior olive communicate with cerebellum

Left

Right



How vestibular nucleus communicates with cerebellum


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine


Left

Right

midline




Basic cerebellar circuit



Deep cerebellar nuclei receives excitatory input (raw/“measured state” of the body and motor commands) and a processed inhibitory input (processed/“predicted state” of the body) and the two are compared.

Layers of cerebellar cortex

NEW YORK INSTITUTE
OF TECHNOLOGY



Molecular layer is closest to the surface - contains dendrites of Purkinje cells and axons of granule cells

Granule layer is the deepest layer and more neurons than all others in the nervous system.

Purkinje cell layer is between the two; one layer of Purkinje cell bodies.

Exclusive output from the cerebellar cortex. GABAergic/inhibitory.

Summary of internal connections

- Climbing fibers
 - olivocerebellar tract
 - excite cerebellar cortex and deep cerebellar nuclei
- Mossy fibers
 - all other tracts
 - excite cerebellar cortex and deep cerebellar nuclei
- Purkinje cells
 - output of cerebellar cortex
- Deep cerebellar nuclei
 - excited by mossy fibers and climbing fibers (“raw” measured state)
 - inhibited by Purkinje cells (highly processed “predicted state”)
 - difference of this two is the final output of cerebellum

Cerebellar disorders can result from a wide variety of causes

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

Environmental: alcohol abuse, lithium, gluten intolerance

Acute diseases: tumors, stroke (lateral medullary syndrome), trauma, & viral infections (chicken pox)

Congenital malformations: Arnold-Chiari malformations & Dandy-Walker

Progressive hereditary disorders: Friedreich's ataxia, multiple sclerosis

ATAXIA is the most common symptom - dyscoordination in timing & magnitude -> overshoots, misdirected movements, and oscillations.

Not weakness (palsy, paralysis), or spasticity!

Similar to alcohol intoxication and are scrutinized in "Standard Field Sobriety Test".

Damage to just one region is rare, e.g. each cerebellar artery supplies different functional divisions.

Damage specifically on one side would impact coordination on the same side.


Cerebellum controls the ipsilateral body.

Cerebrocerebellar Dysfunction


NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

Intention tremor



Problems with rapid alternating movements*



- Symptoms**
- Ataxia incl. misdirections and overshoots by the upper limb
 - Intention Tremor (increasing tremor as hand approaches the movement target) NOT resting tremor
 - *Dysdiadochokinesia (DDK), pronunciation → dis-die-a-dah-ko-kinesia
 - Slurred speech
 - Hypotonia (low muscle tone)


Causes – Head trauma affecting dentate nuclei or lateral hemisphere, pontine degeneration

Spinocerebellar dysfunction

Anterior lobe degeneration



Dandy-Walker



**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Symptoms – Ataxic gait (feet too far apart or too close in irregular progression, stagger, and reel in all directions as if intoxicated by alcohol)

Causes – Anterior lobe degeneration (chronic alcoholism)
– Dandy-Walker syndrome (underdeveloped vermis)
– Arnold-Chiari syndrome (displaced tonsils)




Ataxic gait

Vestibulocerebellar dysfunction


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Balance impairment



Nystagmus (oscillating eye movements)



- Symptoms**
- Wide stance, loss of balance* -> fall to affected side, *coordinated gait if the body is supported!
 - Nystagmus, inaccurate eye tracking & saccades, blurred vision
 - Vertigo and nausea

- Causes**
- Medulloblastoma, vestibular cerebellar syndrome, viral infection

Handy mnemonic! and some video examples

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

DANISH

- Dysdiadochokinesia <https://www.youtube.com/watch?v=ylcWsoBOGzk>
- Ataxia https://www.youtube.com/watch?v=vka_GwXd1os
- Nystagmus
- Intention tremor <https://www.youtube.com/watch?v=-dFMisBI1aM>
- Speech
- Hypotonia



Essential Tremor

Most common motor disorder, ~2% of population

Present during postural hold or movement

(not rest tremor nor intention tremor)

Nearly always involves the upper limb

(sometimes the vocal cords and head)

Not associated with Parkinson's dysfunction

Appears around 40-50 yro

Progressive


Strong genetic component (50% inheritance)

No definitive cause **BUT** the consensus opinion points to cerebellar dysfunction


**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

Spiral Drawing Task



Essential Tremor



Healthy Control

Ali (2022) Wearable sensors during drawing tasks to measure severity of essential tremor

<https://www.youtube.com/watch?v=U3Ktz-UbqDo>

Summary

- Cerebellum is composed of a cortical sheet and internal nuclei.
- Processes information from the spinal cord, brainstem and cerebral cortex.
- Functional divisions defined by cortical territory, output nuclei, and capabilities.
- Information processing involves mossy fibers, climbing fibers, granule cells, Purkinje cells and deep cerebellar nuclei.
- Cerebellum is essential for coordination. Dysfunction is manifest as ataxia.

Feedback is appreciated!

<https://comresearchdata.nyit.edu/redcap/surveys/?s=HRCY448FWYXREL4R>

