NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic Medicine

Arterial Blood Gas Interpretation and Management of Acute Respiratory Distress Syndrome

Michael Gindi, MD, FACEP
Associate Professor of Emergency Medicine
mgindi@nyit.edu

Do.
Make.
Heal.
Innovate.
Reinvent the Future.

Session Objectives

At the conclusion of the lecture, the student should be able to:

1. **Determine which acid-base disorder(s) is/are affecting a patient using arterial blood gas results**
2. **Identify anion gap and non-anion gap metabolic acidosis related conditions**
3. **List the clinical disorders commonly associated with Acute Respiratory Distress Syndrome (ARDS)**
4. **Develop a treatment plan for treatment of ARDS**
5. **Differentiate ARDS from other types of Respiratory Failure**

Drawing an Arterial Blood Gas (ABG)

- Palpate the radial (or alternate) artery
  - Test for collateral circulation with the Allen test
- Prepare the skin
- Insert the needle/syringe at a 40-60 degree angle into the pulsatile artery
- Allow syringe to self-fill with 1cc of arterial sourced blood. *The syringe is heparinized and vented. Don't pull up on syringe because an air bubble can't cause errors.*
- Record FiO2 (0.21 on room air)
- Place specimen on ice and send to lab ***immediately***
- Apply compression for 5 minutes

Before or after the lecture, please watch the video "Harrison's Percutaneous Arterial Blood Gas Sampling" accessible via NYIT COM's subscription to Access Medicine

NORMAL ARTERIAL BLOOD GAS VALUES

Results will be listed as:

pH*	7.35 – 7.45
$PaCO_2$	35 – 45 mmHg
PaO_2	80 – 100 mmHg
HCO_3^{-}**	22 – 26 mEq/L
O_2 % sat	96 – 99%

*Normal pH: 7.35 – 7.45. The body DOES NOT OVERCORRECT

**Also called “bicarb”

ABG's are Painful and Dangerous

- Fewer ABG's are being done than ever before*
- Modern pulse oximetry is very accurate – the main exceptions are carbon monoxide poisoning, severe anemia and dark nail polish
- Modern end-tidal CO2 measurements are very accurate
- Venous Ph is *about* .05 less than arterial
- The bicarb value reported in a serum electrolyte panel (usually drawn from a venous sample) is **measured**. The bicarb value reported in an ABG is **calculated** and thus is less reliable anyway.

*From: https://litfl.com/vbg-versus-abg/

Step-wise approach to Simple Acid-Base Disorders

1. **First: Evaluate the pH:** If the pH < 7.4 then the patient is acid**emic.** If the pH > 7.4 then alkal**emic.** (If pH is normal, then there is either no acid base disorder or a mixed acid-base disorder)

2. **Next: Check the PC02 and HCO3-**

3. **Then ask: Which of those two values could cause that Ph abnormality?** Then that is the one causing the primary process (“osis” is the Greek for suffix for “process)

4. Then the **OTHER process** is the one **COMPENSATING** for the primary process

Of course, in a real patient, there can be a mixed picture with two primary processes. However, for your exam or the Boards we would not write a question that complicated.

Distinguishing Primary (Initial) Change from Compensatory Response

PH	Acid Base Disorder	Initial Chemical Change	Compensatory Response
<7.4 -	Respiratory Acidosis	$\uparrow PCO_2$	$\uparrow HCO_3^-$
>7.4 -	Respiratory Alkalosis	$\downarrow PCO_2$	$\downarrow HCO_3^-$
<7.4 -	Metabolic Acidosis	$\downarrow HCO_3^-$	$\downarrow PCO_2$
>7.4 -	Metabolic Alkalosis	$\uparrow HCO_3^-$	$\uparrow PCO_2$


Formulas* for Determining Expected Changes and Their Chronicity. NO NEED TO MEMORIZE these but have them available.

Metabolic Acidosis	$Pa_{CO2} = (1.5 \times HCO_3^-) + 8 \pm 2$
Metabolic Alkalosis	Pa_{CO2} will ↑0.75 mmHg per mmol/L ↑in $[HCO_3^-]$
Respiratory Alkalosis	
Acute	$[HCO_3^-]$ will ↓0.2 mmol/L per mmHg ↓in Pa_{CO2}
Chronic	$[HCO_3^-]$ will ↓0.4 mmol/L per mmHg ↓in Pa_{CO2}
Respiratory Acidosis	
Acute	$[HCO_3^-]$ will ↑0.1 mmol/L per mmHg ↑in Pa_{CO2}
Chronic	$[HCO_3^-]$ will ↑0.4 mmol/L per mmHg ↑in Pa_{CO2}

The Davenport Diagram

Watch this video to learn how to use this diagram: https://www.youtube.com/watch?v=snp5oq-hLpg

Example ABG:
PH = 7.3
HCO3- = 16
PCO2 = 30



Source: Longo DL, Fauci AS, Kasper DL, Hauser SL, Jameson JL, Loscalzo J: *Harrison's Principles of Internal Medicine, 18th Edition*: www.accessmedicine.com

Copyright © The McGraw-Hill Companies, Inc. All rights reserved.

What is the anion gap (AG)?

It represents unmeasured anions in the plasma.

Normal range is 8-16
(depends on your lab)



Image from: https://elite.com.au/static-electricity-carpet-tips-pain/

How do you calculate it?

$AG = Na^+ - (Cl^- + HCO_3^-)$

What does it tell you?

Increased AG: Excess anions are being added to the plasma

High Anion Gap Metabolic Acidosis
Differential Diagnosis

- M Methanol
- U Uremia
- D Diabetic/Starvation/Alcoholic Ketoacidosis
- P Paraldehyde
- I Iron, Isoniazid, Inborn Errors
- L Lactic Acidosis (THE most common)
- E Ethylene Glycol
- S Salicylates

Normal or Non-Anion Gap Metabolic Acidosis

- GI Bicarb loss
  - Diarrhea
  - Drugs(Laxatives)
- Renal cause
  - RTA type 1 and 2
  - Hyperkalemia
- Drugs
  - Diuretics, ACE-I, ARB
  - NSAIDS
  - Cyclosporine
  - Trimethroprim
  - Pentamidine

Management: Metabolic Acidosis

First and foremost treat the underlying cause

Although it intuitively make sense to administer bicarb, **it is almost never indicated** solely to improve PH.

*It is only indicated in specific scenarios: e.g hyperkalemia, cyclic antidepressant overdose, other Na+ channel blockades, acute kidney injury, alkalinization of urine for certain kidney stones and toxicities, certain non-anion gap acidoses.

*A Review of Bicarbonate Use in Common Clinical Scenarios
Gabriel Wardi, MD,MPH*,† gwardi@ucsd.edu · Sarah Holgren, MD‡ · Arnav Gupta, MD* Alex Pearce, MD† · Atul Malhotra, MD† · Christopher Tainter, MD‡ JEM, Volume 65, Issue 2 E71-E80August 2023

BICARB IS

- **NOT** for DKA – it actually worsens outcome
- **NOT** for rhabdomyolysis – its only benefit is a DOO not POO. It decreases urinary casts that can obstruct the renal tubules, BUT ***there is no difference in acute kidney injury rate or mortality.***
- **NOT** because you are unhappy with the Ph

ETHYLENE GLYCOL

- Source: Antifreeze
- Cause of Toxicity:

Alcohol Dehydrogenase

↓

Glycoaldehyde

↓

Glycolic, Oxalic, Glyoxylic Acids

↓

Metabolic acidosis

METHANOL

- Source: solvents, paint removers, moonshine (100,000 cases of blindness during Prohibition)*
- Cause of Toxicity:

Alcohol Dehydrogenase

↓

Formaldehyde

↓

Formate and lactate

↓

Metabolic acidosis

*From: https://medium.com/creative-science/methanol-in-the-moonshine-b7e38d695717

Blind Blues Musicians

- Why were so many blues musicians blind?*
- Blind Willie Johnson, Blind Willie McTell, Blind Lemon Jefferson, Blind Joe Fuller, Blind Joe Reynolds etc. . .
- Multifactorial: poverty, poor healthcare access, environmental toxins, malnutrition
- **Improper distillation of ethanol (Moonshine) leading to retinopathy**
- I'd sing the blues too

*https://allthatsinteresting.com/blind-blues-musicians

ETHYLENE GLYCOL OR METHANOL TREATMENT:

- Fomepizole (Antizol™) – inhibits alcohol dehydrogenase so the liver has a chance to metabolize the toxins
- Thiamine and pyrodoxine replacement
  - Act as cofactors in ethylene glycol metabolism
- Saline diuresis
- Ethanol IV (or PO) to compete with enzymatic metabolism
- Dialysis



Image from:https://spiritedgifts.com/absolut-vodka

Salicylate Overdose – one example of a mixed picture



Initially salicylates stimulate the respiratory center

↓

Respiratory alkalosis initially

then

Uncoupling of oxidative phosphorylation

↓

Disrupts glucose and fatty acid metabolism

↓

Metabolic acidosis

SALICYLATES TREATMENT:

- Consider gastric lavage if massive ingestion that happened less than 30 minutes ago
- Activated charcoal
- Intravenous $NaHCO_3^-$ to alkalinize the urine – enhances salicylate excretion
- Dialysis
- Consider acetazolamide if respiratory alkalosis predominates (MOA on next slide)

Metabolic Alkalosis Causes

- Antacids
- Baking soda
- Diuretics
- Certain laxatives
- Steroids

Treatment: underlying cause*

*Some cases may require acetazolamide, which causes a metabolic acidosis. It works in acute mountain sickness by triggering a compensatory respiratory alkalosis by increasing minute ventilation which, in turn, increases oxygenation

From: https://my.clevelandclinic.org/health/diseases/21529-metabolic-alkalosis

Respiratory acidosis treatment is aimed at improving ventilation

- Limit sedative and CNS depressants use
- Treat COPD exacerbations carefully because they, unlike healthy individuals, rely on a hypoxic drive, because they are chronically hypercapnic
- Poor ventilation may require noninvasive positive pressure ventilation (CPAP or BIPAP) or intubation with mechanical ventilation

End Organ Effects of Respiratory Alkalosis

- Early: lightheadedness, confusion, syncope, tachycardia, anxiety, dyspnea, **binding of Ca_2^+ to albumin** → *circumoral or acral paresthesias, seizures, tetany*
  - *Breathing into a paper bag does may not raise CO2 and may cause hypoxia.* So you better be right that it's only a panic attack.*
- Late: Apathy, confusion, neuromuscular instability, cardiac arrhythmias (often due to hypokalemia)

*HYPOXIC HAZARDS OF TRADITIONAL PAPER BAG REBREATHING IN HYPERVENTILATING PATIENTS
Callaham, M., Ann Emerg Med 18(6):622, June 1989

Management: Respiratory Alkalosis

- Treat underlying cause (Are you seeing a pattern here?) – most often is by calm reassurance
- Don’t infuse acid
- Hypocapnia may be iatrogenic, so check your ventilatory settings

Acid-Base Case Study 1

A 60-year-old patient with COPD on multiple inhalers, hypertension, GERD, peptic ulcer disease, and CKD presents with altered mental status from home. He is on omeprazole for GERD, but his son notes he often takes over-the-counter antacids as well.

Source: Case Studies: Clinical Scenario 23, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 1

A 60-year-old patient with COPD, GERD, PUD, and CKD presents with altered mental status from home. He is supposed to take omeprazole for GERD, but his son notes he often takes over-the-counter antacids instead.

- COPD → Respiratory Acidosis (chronic)
- CKD → Metabolic Acidosis
- OTC antacids (if TUMS…) → Metabolic alkalosis (milk-alkali syndrome)

This case would be present with a challenging ABG

Source: Case Studies: Clinical Scenario 23, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 2

A 44-year-old homeless man suffering from schizophrenia and alcoholism presents to the emergency department with altered mental status and renal failure. He is tachypneic and is intubated shortly after arrival due to an inability to protect his airway. There is no additional history available.

Source: Case Studies: Clinical Scenario 1, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 2

A 44-year-old homeless person with schizophrenia and alcoholism presents to the emergency department with altered mental status and renal failure. He is intubated shortly after arrival due to an inability to protect his airway. There is no additional history available. He is tachypneic with a respiratory rate in the 30s prior to intubation.

- Ethanol ingestion/Methanol toxicity → Metabolic acidosis, anion gap
- Renal Failure → Metabolic Acidosis, non-anion gap
- Respiratory failure, acute→ Respiratory acidosis (acute)

This case would be present with another very challenging ABG but you could guess that ALL of these process would cause acidosis resulting in acidemia.

Source: Case Studies: Clinical Scenario 1, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 3

A 24-year-old woman with chronic migraines presents to the ED with altered mental status and what her husband describes as "her breathing looking uncomfortable." She has been having more frequent headaches in the last few weeks. She takes no prescribed medications.

Source: Case Studies: Clinical Scenario 3, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 3

A 24-year-old woman with chronic migraines presents to the emergency department with altered mental status and what her husband describes as "her breathing looks uncomfortable." She has been having more issues with headaches in the last few weeks, but otherwise the husband notes no other problems. She takes no prescribed medications.

HINT: think of potential ways she might have tried to treat her headache

- **Many analgesics contain aspirin** – mixed disorder of metabolic acidosis and respiratory alkalosis as noted in reading and lecture
- Some analgesics contain opioids – respiratory acidosis from respiratory suppression or opioid-induced pulmonary edema

Source: Case Studies: Clinical Scenario 3, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 4

A 77-year-old male with COPD presents to the ED in respiratory distress.

BP 80/40 mm Hg, HR120 bpm, RR 30, Temp 102.4°F, WBC is 22.3k with 20% bands. He has a large infiltrate in the right lower lobe on chest x-ray.

A month prior, during an outpatient pulmonary appointment, the patient had an ABG: pH 7.35, $PaCO_2$ 65 mm Hg, HCO_3 33 mEq/L.

Source: Case Studies: Clinical Scenario 37, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 4

A 77-year-old male with chronic hypoxic respiratory failure presents to the emergency department with respiratory distress.

His blood pressure is 80/40 mm Hg, and his heart rate is 120 bpm. Temperature is 102.4°F, and his WBC is 22.3k with 20% bands. He has a large infiltrate in the right lower lobe on chest x-ray.

A month prior, during an outpatient pulmonary appointment, the patient had an ABG: pH 7.35, $PaCO_2$ 65 mm Hg, HCO_3 33 mEq/L.

- Pneumonia → Respiratory Acidosis, not ventilating well acutely
- COPD → Respiratory Acidosis, not ventilating well chronically
- Sepsis → Metabolic Acidosis

This case is very challenging (above and beyond what would be expected)

NOTE: You will not need to be able to do this on an exam or boards, but perhaps on wards next year…

Source: Case Studies: Clinical Scenario 37, Author: Zachary Healy from AccessMedicine

Respiratory Failure: Case 1

A 73 y F arrives to the ED complaining of dry cough and breathlessness for 2 weeks. She has a RR of 28, HR of 98, BP of 100/60, SpO2 of 88%, and temperature of 40 C. Her WBC is 15k and CXR shows infiltrates in all lung fields. She decompensates and dies soon after. At autopsy, both lungs were voluminous and microscopic examination revealed acute-stage, severe diffuse alveolar damage with remarkable *mononuclear cell infiltration and hyaline membrane* formation in the lungs. **Which of the following is most likely cause of death?**

A. Pulmonary embolism
B. Bilateral interstitial pneumonia
C. COPD exacerbation
D. Acute respiratory distress syndrome (ARDS)
E. Hypersensitivity pneumonitis

ARDS: Epidemiologic Features

10-86/100,000 with highest rates reports in the USA and Australia.

ARDS likely to be underreported in low-income countries

Under recognized in high-income countries:

- 459 ICUs in 50 countries revealed recognition rates of
  - 51% for mild ARDS
  - 79% for severe ARDS

Decreased recognition may be due to confusion surrounding radiographic findings

NEJM 2017; 277:562-572

Incidence of ARDS in the ICU

(the fellowship is called “Pulmonary and Critical Care”)

- **10.4% of all ICU patients**
- **67.2% of patients with acute hypoxemic respiratory failure**
- **ARDS was frequently undiagnosed (only 60.2% recognized by clinician) or late diagnosed (34.0% recognized at the time of criteria fulfillment)**
- **Mortality rate ~ 38%**


The Berlin Definition/Criteria for ARDS

ARDS in noncardiogenic pulmonary edema due to permeability of pulmonary capillaries leading to increased barrier to O2 exchange and hypoxemia

Criteria	Definition
Onset	Within 7 days of known clinical insult or worsening respiratory symptoms
Chest X-ray	Bilateral opacities consistent with pulmonary edema and not fully explained by effusions, atelectasis, or nodules
Edema	Respiratory failure with pulmonary edema due to non-cardiogenic causes (PCWP ≤ 18 mmHg)
Oxygenation	PaO2/FiO2 < 300 mmHg w/ minimum PEEP or CPAP > 5 cm H2O

NEJM 2017; 277:562-572 https://www-uptodate-com.arktos.nyit.edu/contents/noncardiogenic-pulmonary-edema?search=ARDS%20and%20PCWP&source=search_result&selectedTitle=4~150&usage_type=default&display_rank=4


Chest X ray



Chest radiograph showing opacifications in both lung fields, compatible with acute respiratory distress syndrome (white out)

Case courtesy of Dr. Craig Hacking, Radiopaedia.org, rID: 53758

Pulmonary Capillary Wedge Pressure (PCWP)



Source: Maxine A. Papadakis, Stephen J. McPhee, Michael W. Rabow:
Current Medical Diagnosis and Treatment 2021
Copyright © McGraw Hill. All rights reserved.

PCWP measurement using a Swan-Ganz catheter:

Normal PCWP = 4-12 mmHg

PCWP > 18 mmHg is consistent with heart failure

PCWP ≤ 18 mmHg is consistent with ARDS

Current Medical Diagnosis and Treatment 2021
https://www-uptodate-com.arktos.nyit.edu/contents/noncardiogenic-pulmonary-edema?search=ARDS%20and%20PCWP&source=search_result&selectedTitle=4~150&usage_type=default&display_rank=4

Staging and Prognosis

Stage	PaO2:FiO2	Mortality
Normal	> 300 mmHg	N/A
Mild	201 – 300 mmHg	27%
Moderate	101 – 200 mmHg	32%
Severe	<100 mmHg	45%

Example calculation of a NORMAL patient:

- **PaO2 (partial pressure of O2) on ABG= 90mmHg**
- **FiO2 = (fraction of inspired O2) on room air = 0.21**
  PaO2:FiO2 = 90/0.21 = 428 mmHg

NEJM 2017; 277:562-572

Risk Factors

Direct Lung injury	Indirect Lung injury
Pneumonia (bacterial, viral, fungal, opportunistic)*	**Sepsis (non-pulmonary source)***
Aspiration of gastric contents*	Disseminated intravascular coagulation (DIC)
Pulmonary contusion	Shock
Inhalation injury	Acute pancreatitis
Near drowning	Major burn injury
Diffuse lung disease	Drug overdose
Air embolism	Transfusion of blood products

* accounts for over 85% of ARDS cases

NEJM 2017; 277:562-572

ARDS vs. CHF

	ARDS	CHF
Exacerbation	Sepsis, pneumonia, aspiration, near drowning, transfusion	HTN, MI, fluid overload
Chest X-ray	Severe bilateral infiltrates, “white out”	Bilateral pleural effusions
PCWP	Normal (4-12 mmHg)	Elevated (> 18 mmHg)
LV Function	Normal	

NEJM 2017; 277:562-572
https://www-uptodate-com.arktos.nyit.edu/contents/noncardiogenic-pulmonary-edema?search=ARDS%20and%20PCWP&source=search_result&selectedTitle=4~150&usage_type=default&display_rank=4

Treatment – Starts with Prevention

Good ICU practices:

- **Lower tidal volumes** for all mechanically ventilated patients - avoids barotrauma
- **Early volume resuscitation and antibiotics** for sepsis
- **Restrictive use of blood** products - reduces risk of transfusion-associated lung injury & volume overload
- **Intensivist involvement** to help prevent development of nosocomial ARDS

NEJM 2017; 277:562-572

Treatment

- **Treat the underlying disease**
- **Supportive therapy: focused on limiting further injury**
  - Lung-protective ventilation to prevent ventilator-associated lung injury
  - Use tidal volumes of only 4-8 ml/kg
  - Judicious fluid therapy to prevent lung edema

Treatment (continued)

- **No pharmacologic therapy for ARDS has been shown to reduce short or long-term mortality**
- **Use PEEP** (Positive End Expiratory Pressure) in your ventilator settings of **at least 5 cm H2O**
  - PEEP < 5 resulted increased mortality vs. PEEP > 16 adults with moderate to severe ARDS*

NEJM 2017; 277:562-572
https://www.ficm.ac.uk/sites/default/files/prone_position_in_adult_critical_care_2019.pdf
https://www.uptodate.com/contents/prone-ventilation-for-adult-patients-with-acute-respiratory-distress-syndrome

Role of PEEP in Recruiting Alveoli



Normal

ARDS with exudate (protein)

ARDS with PEEP

Treatment (continued)

Ventilating in prone position keeps heart and mediastinum from pressing against lungs



COVID-19 and ARDS: What did we learn?

- **Pathology of severely ill COVID-19 patients is consistent with that of early-stage ARDS during autopsy***
  - Hyaline membrane formation and micro vessel thrombosis
  - Anticoagulation helps in severe cases**
- **Management of severe COVID -19 is similar to ARDS**
  - Mechanical ventilation with low tidal volume and high peep
  - But add dexamethasone
  - ?Inhaled nitric oxide: vasodilation, antiviral, anti-inflammatory. Better O2 saturation but same mortality- another DOO not POO

*https://www-uptodate-com.arktos.nyit.edu/contents/covid-19-epidemiology-clinical-features-and-prognosis-of-the-critically-ill-adult?search=COVID%20pneumonia%20and%20ARDS&source=search_result&selectedTitle=3~150&usage_type=default&display_rank=3

Ann Transl Med. 2018 Jan;6(2):36. doi: 10.21037/atm.2018.01.08 **Anticoagulant therapy in acute respiratory distress syndrome
Marta Camprubí-Rimblas [1,2], Neus Tantinyà [3], Josep Bringué [1,2,3], Raquel Guillamat-Prats [1,3], Antonio Artigas[1,2,3,4,✉]

Summary

- Approach ABG interpretation in a step-wise fashion, but try to predict what the disorder SHOULD be based on the clinical picture
- If the ABG does not show a simple process, then use your formulas or the Davenport Diagram
- ARDS is an extremely common, heterogeneous and dangerous syndrome
- Focus on treating and/or avoiding the insults which lead to ARDS
- Good ICU practices coupled with early treatment can prevent ARDS
- Early lung-protective ventilation and sensible fluid management remain the mainstay of ARDS treatment

Lecture Feedback Form:

https://comresearchdata.nyit.edu/redcap/surveys/?s=HRCY448FWYXREL4R