Session Objectives

At the conclusion of the lecture, the student should be able to:

1. **Determine which acid-base disorder(s) is/are affecting a patient using arterial blood gas results**
2. **Identify anion gap and non-anion gap metabolic acidosis related conditions**
3. **List the clinical disorders commonly associated with Acute Respiratory Distress Syndrome (ARDS)**
4. **Develop a treatment plan for treatment of ARDS**
5. **Differentiate ARDS from other types of Respiratory Failure**

Step-wise approach to Simple Acid-Base Disorders

1. **First: Evaluate the pH:** If the pH < 7.4 then the patient is acid**emic.** If the pH > 7.4 then alkal**emic.** (If pH is normal, then there is either no acid base disorder or a mixed acid-base disorder)

2. **Next: Check the PC02 and HCO3-**

4. Then the **OTHER process** is the one **COMPENSATING** for the primary process

Of course, in a real patient, there can be a mixed picture with two primary processes. However, for your exam or the Boards we would not write a question that complicated.

Distinguishing Primary (Initial) Change from Compensatory Response

PH	Acid Base Disorder	Initial Chemical Change	Compensatory Response
<7.4 -	Respiratory Acidosis	$\uparrow PCO_2$	$\uparrow HCO_3^-$
>7.4 -	Respiratory Alkalosis	$\downarrow PCO_2$	$\downarrow HCO_3^-$
<7.4 -	Metabolic Acidosis	$\downarrow HCO_3^-$	$\downarrow PCO_2$
>7.4 -	Metabolic Alkalosis	$\uparrow HCO_3^-$	$\uparrow PCO_2$

High Anion Gap Metabolic Acidosis
Differential Diagnosis

- M Methanol
- U Uremia
- D Diabetic/Starvation/Alcoholic Ketoacidosis
- P Paraldehyde
- I Iron, Isoniazid, Inborn Errors
- L Lactic Acidosis (THE most common)
- E Ethylene Glycol
- S Salicylates

Normal or Non-Anion Gap Metabolic Acidosis

- GI Bicarb loss
  - Diarrhea
  - Drugs(Laxatives)
- Renal cause
  - RTA type 1 and 2
  - Hyperkalemia
- Drugs
  - Diuretics, ACE-I, ARB
  - NSAIDS
  - Cyclosporine
  - Trimethroprim
  - Pentamidine

Salicylate Overdose – one example of a mixed picture



Initially salicylates stimulate the respiratory center

↓

Respiratory alkalosis initially

then

Uncoupling of oxidative phosphorylation

↓

Disrupts glucose and fatty acid metabolism

↓

Metabolic acidosis

Respiratory acidosis treatment is aimed at improving ventilation

- Limit sedative and CNS depressants use
- Treat COPD exacerbations carefully because they, unlike healthy individuals, rely on a hypoxic drive, because they are chronically hypercapnic
- Poor ventilation may require noninvasive positive pressure ventilation (CPAP or BIPAP) or intubation with mechanical ventilation

End Organ Effects of Respiratory Alkalosis

- Early: lightheadedness, confusion, syncope, tachycardia, anxiety, dyspnea, **binding of Ca_2^+ to albumin** → *circumoral or acral paresthesias, seizures, tetany*
  - *Breathing into a paper bag does may not raise CO2 and may cause hypoxia.* So you better be right that it's only a panic attack.*
- Late: Apathy, confusion, neuromuscular instability, cardiac arrhythmias (often due to hypokalemia)

*HYPOXIC HAZARDS OF TRADITIONAL PAPER BAG REBREATHING IN HYPERVENTILATING PATIENTS
Callaham, M., Ann Emerg Med 18(6):622, June 1989

Acid-Base Case Study 1

A 60-year-old patient with COPD on multiple inhalers, hypertension, GERD, peptic ulcer disease, and CKD presents with altered mental status from home. He is on omeprazole for GERD, but his son notes he often takes over-the-counter antacids as well.

Source: Case Studies: Clinical Scenario 23, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 1

A 60-year-old patient with COPD, GERD, PUD, and CKD presents with altered mental status from home. He is supposed to take omeprazole for GERD, but his son notes he often takes over-the-counter antacids instead.

- COPD → Respiratory Acidosis (chronic)
- CKD → Metabolic Acidosis
- OTC antacids (if TUMS…) → Metabolic alkalosis (milk-alkali syndrome)

This case would be present with a challenging ABG

Source: Case Studies: Clinical Scenario 23, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 2

A 44-year-old homeless man suffering from schizophrenia and alcoholism presents to the emergency department with altered mental status and renal failure. He is tachypneic and is intubated shortly after arrival due to an inability to protect his airway. There is no additional history available.

Source: Case Studies: Clinical Scenario 1, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 2

A 44-year-old homeless person with schizophrenia and alcoholism presents to the emergency department with altered mental status and renal failure. He is intubated shortly after arrival due to an inability to protect his airway. There is no additional history available. He is tachypneic with a respiratory rate in the 30s prior to intubation.

- Ethanol ingestion/Methanol toxicity → Metabolic acidosis, anion gap
- Renal Failure → Metabolic Acidosis, non-anion gap
- Respiratory failure, acute→ Respiratory acidosis (acute)

This case would be present with another very challenging ABG but you could guess that ALL of these process would cause acidosis resulting in acidemia.

Source: Case Studies: Clinical Scenario 1, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 3

A 24-year-old woman with chronic migraines presents to the ED with altered mental status and what her husband describes as "her breathing looking uncomfortable." She has been having more frequent headaches in the last few weeks. She takes no prescribed medications.

Source: Case Studies: Clinical Scenario 3, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 3

A 24-year-old woman with chronic migraines presents to the emergency department with altered mental status and what her husband describes as "her breathing looks uncomfortable." She has been having more issues with headaches in the last few weeks, but otherwise the husband notes no other problems. She takes no prescribed medications.

HINT: think of potential ways she might have tried to treat her headache

- **Many analgesics contain aspirin** – mixed disorder of metabolic acidosis and respiratory alkalosis as noted in reading and lecture
- Some analgesics contain opioids – respiratory acidosis from respiratory suppression or opioid-induced pulmonary edema

Source: Case Studies: Clinical Scenario 3, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 4

A 77-year-old male with COPD presents to the ED in respiratory distress.

BP 80/40 mm Hg, HR120 bpm, RR 30, Temp 102.4°F, WBC is 22.3k with 20% bands. He has a large infiltrate in the right lower lobe on chest x-ray.

A month prior, during an outpatient pulmonary appointment, the patient had an ABG: pH 7.35, $PaCO_2$ 65 mm Hg, HCO_3 33 mEq/L.

Source: Case Studies: Clinical Scenario 37, Author: Zachary Healy from AccessMedicine

Acid-Base Case Study 4

A 77-year-old male with chronic hypoxic respiratory failure presents to the emergency department with respiratory distress.

His blood pressure is 80/40 mm Hg, and his heart rate is 120 bpm. Temperature is 102.4°F, and his WBC is 22.3k with 20% bands. He has a large infiltrate in the right lower lobe on chest x-ray.

A month prior, during an outpatient pulmonary appointment, the patient had an ABG: pH 7.35, $PaCO_2$ 65 mm Hg, HCO_3 33 mEq/L.

- Pneumonia → Respiratory Acidosis, not ventilating well acutely
- COPD → Respiratory Acidosis, not ventilating well chronically
- Sepsis → Metabolic Acidosis

This case is very challenging (above and beyond what would be expected)

NOTE: You will not need to be able to do this on an exam or boards, but perhaps on wards next year…

Source: Case Studies: Clinical Scenario 37, Author: Zachary Healy from AccessMedicine

Respiratory Failure: Case 1

A 73 y F arrives to the ED complaining of dry cough and breathlessness for 2 weeks. She has a RR of 28, HR of 98, BP of 100/60, SpO2 of 88%, and temperature of 40 C. Her WBC is 15k and CXR shows infiltrates in all lung fields. She decompensates and dies soon after. At autopsy, both lungs were voluminous and microscopic examination revealed acute-stage, severe diffuse alveolar damage with remarkable *mononuclear cell infiltration and hyaline membrane* formation in the lungs. **Which of the following is most likely cause of death?**


A. Pulmonary embolism
B. Bilateral interstitial pneumonia
C. COPD exacerbation
D. Acute respiratory distress syndrome (ARDS)
E. Hypersensitivity pneumonitis

Incidence of ARDS in the ICU

(the fellowship is called “Pulmonary and Critical Care”)

- **10.4% of all ICU patients**
- **67.2% of patients with acute hypoxemic respiratory failure**
- **ARDS was frequently undiagnosed (only 60.2% recognized by clinician) or late diagnosed (34.0% recognized at the time of criteria fulfillment)**
- **Mortality rate ~ 38%**

Chest X ray



Chest radiograph showing opacifications in both lung fields, compatible with acute respiratory distress syndrome (white out)

Case courtesy of Dr. Craig Hacking, Radiopaedia.org, rID: 53758

Staging and Prognosis

Stage	PaO2:FiO2	Mortality
Normal	> 300 mmHg	N/A
Mild	201 – 300 mmHg	27%
Moderate	101 – 200 mmHg	32%
Severe	<100 mmHg	45%

Example calculation of a NORMAL patient:

- **PaO2 (partial pressure of O2) on ABG= 90mmHg**
- **FiO2 = (fraction of inspired O2) on room air = 0.21**
  PaO2:FiO2 = 90/0.21 = 428 mmHg

NEJM 2017; 277:562-572

Risk Factors

Direct Lung injury	Indirect Lung injury
Pneumonia (bacterial, viral, fungal, opportunistic)*	**Sepsis (non-pulmonary source)***
Aspiration of gastric contents*	Disseminated intravascular coagulation (DIC)
Pulmonary contusion	Shock
Inhalation injury	Acute pancreatitis
Near drowning	Major burn injury
Diffuse lung disease	Drug overdose
Air embolism	Transfusion of blood products

* accounts for over 85% of ARDS cases

NEJM 2017; 277:562-572

Treatment – Starts with Prevention

Good ICU practices:

- **Lower tidal volumes** for all mechanically ventilated patients - avoids barotrauma
- **Early volume resuscitation and antibiotics** for sepsis
- **Restrictive use of blood** products - reduces risk of transfusion-associated lung injury & volume overload
- **Intensivist involvement** to help prevent development of nosocomial ARDS

NEJM 2017; 277:562-572

Summary

- Approach ABG interpretation in a step-wise fashion, but try to predict what the disorder SHOULD be based on the clinical picture
- If the ABG does not show a simple process, then use your formulas or the Davenport Diagram
- ARDS is an extremely common, heterogeneous and dangerous syndrome
- Focus on treating and/or avoiding the insults which lead to ARDS
- Good ICU practices coupled with early treatment can prevent ARDS
- Early lung-protective ventilation and sensible fluid management remain the mainstay of ARDS treatment