Session Objectives

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

1. Identify the properties of the most common primary fungal pathogens of the respiratory tract.
2. Identify the etiology, clinical presentation, diagnosis, and treatment for each of the fungal respiratory pathogens.
3. Describe the epidemiology involved in the spread and acquisition of respiratory fungal diseases.
4. Define the pathogenesis of the fungal agents.
5. Outline the laboratory techniques used for the identification and diagnosis of these fungal pathogens.

Systemic Fungal Infections

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- Uncommon
  - Most superficial
- Natural immunity is high
  - Innate and cell mediated
- Cases have increased over the past few decades
- High mortality rate, treatment is challenging
- Most are asymptomatic or self-limiting
- Most acquired through respiratory track

Coccidioidomycosis

NEW YORK INSTITUTE OF TECHNOLOGY

- **Risks Factors**
  - Ethnicity (African Americans, Hispanic & Filipinos)
  - Age: Extremes more susceptible
  - Sex: Males more susceptible
  - Pregnancy: 3rd trimester
  - Immunosuppression (HIV, transplants, etc.)
- **Diagnosis**
  - Histopathology: spherules or spores seen in sputum, exudates or tissue is sufficient
  - Culture: dangerous! (direct microscopic observation preferred)
  - Serology
- **Treatment**
  - Fluconazole or itraconazole for mild
  - Amphotericin B for disseminated
  - Meningeal coccidioidomycosis is managed with fluconazole







Histoplasmosis

NEW YORK INSTITUTE


- *H. capsulatum* (US); var: *duboisii* (Africa)
- Endemic areas:
  Central & Midwestern states (most cases in Ohio Valley & Mississippi Valley)
- Dimorphic/soil saprophyte
  - Hyphae contain macro & microconidia
- Birds/bats droppings (guano)
- Cave disease




Yeast @ 37°C




Conidia @ 25 °C





Histoplasmosis

- 90% of cases are asymptomatic
  - Rare cases show flu-like symptoms
- **Immune Response**
  - Phagocytic activity of macrophage is considered an important component of resistance
    - Survival in macrophages—elevates pH of phagosomes
    - Macrophages spread yeast
  - Cell-mediated response help clear infection
    - Activated macrophage can kill yeast cells
    - Granuloma formation (potential for reactivation)
- Disseminated histoplasmosis can involve many systems
  - Diagnosed frequently in patients with AIDS living in the central U.S.





Histoplasmosis

NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic Medicine

- **Diagnosis**
  - Direct histology & culture of blood or bone marrow
    - Visualization of yeast cells inside macrophages
  - Serological testing for antibody or Ag in blood & urine
  - PCR

- **Treatment**
  - Itraconazole for mild or moderate disease
  - Amphotericin B for disseminated and severe pulmonary histoplasmosis



Blastomycosis

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine


- **Pathogenesis and Immune response:**
  - Alveolar macrophage provide a first line of defense
    - Conidia are more sensitive to killing because yeast are too big
  - Escapes phagocytosis
  - T-cells stimulate neutrophils to kill *Blastomyces* cells
  - Pyogranulomas form
    - Granulomatous infection predominantly involves lungs
- Dissemination
  - Commonly skin, bones, GU tract



Paracoccidiodomycosis

NEW YORK INSTITUTE OF TECHNOLOGY

- *Paracoccidioides brasiliensis*
- Distributed in Central and South America
- Dimorphic
    - Yeast have multiple buds (captain's wheel)
- Lung infection, common in older men (farmers, rural laborers)
- Dissemination uncommon (oral mucosa most common site)
- Treatment:
    - Itraconazole
    - Sulfonamides
    - Amphotericin B



Source: P. Chin-Hong, E. A. Joyce, M. Karandikar, M. Matloubian, L. A. Rubio, B. S. Schwartz, W. Levinson: Levinson's Review of Medical Microbiology & Immunology: A Guide to Clinical Infectious Diseases, 19th Edition: Copyright © McGraw Hill. All rights reserved.





Talaromycosis

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- *Talaromyces marneffei*
    - Former genus: *Penicillium*
- Endemic in Southeast Asia
- Disease develops in immunocompromised, especially those with HIV
- Dimorphic/soil; Opportunistic
- Yeast is elongate with central septum
- Dissemination possible
- Treatment: Amphotericin B and/or itraconazole



Summary

NEW YORK INSTITUTE OF TECHNOLOGY
College of Osteopathic Medicine

- The endemic mycoses are common infections that exist within restricted geographical areas
- Primary fungal pathogens that cause systemic disease are typically thermally dimorphic
- Clinical manifestations are unique to each endemic fungus, but the lung is prominently involved
- Cell-mediated immunity is essential for successful resolution of these infections
- Impaired immunity can facilitate systemic fungal disease