

Session Objectives

- 1 Students will know the risk and protective factors for suicide.
- 2 Students will be able to apply the characteristics of the suicidal patient to clinical scenarios.
- 3 Students will learn the components of a suicide and dangerous behavior assessment.
- 4 Students will understand the privilege of confidentiality and its exceptions.
- 5 Students will learn the best practices to safely assess the violent, aggressive patient

Suicide Data: United States

Suicide is a public health problem and leading cause of death in the United States. Suicide can also be prevented – more investment in suicide prevention, education, and research will prevent the untimely deaths of thousands of Americans each year. Unless otherwise noted, this fact sheet reports 2022 data from the CDC, the most current verified data available at time of publication (May 2024).

49,476 Americans died by suicide making it the **11th leading cause of death**.



- **3rd** leading cause of death for ages 15-24
- **2nd** leading cause of death for ages 25-34
- **4th** leading cause of death for ages 35-44
- **Over one third** of people who died by suicide were 55 or older



4.8% of adult Americans have thought about suicide.

1.6 million Americans attempted suicide.

54% of Americans have been affected by suicide in some way.

Men died by suicide **3.85 times** more often than females.

Women were **1.3 times** more likely to attempt suicide.

56% of firearm deaths were suicides.

55% of all suicides were by firearms.

In 2021, the suicide rate was **2 times higher for Veterans** than for non-Veteran adults over the age of 18.



90% of those who died by suicide had a diagnosable mental health condition at the time of their death.

46% of Americans ages 18+ living with a mental health condition received treatment in the past year.

73% of communities in the United States did not have enough mental health providers to serve residents in 2022, according to federal guidelines.




See full list of citations at afsp.org/statistics.



Suicide Data: United States

- States in the top third of suicide rates in the U.S.
- States in the middle third of suicide rates in the U.S.
- States in the lower third of suicide rates in the U.S.



See full list of citations at afsp.org/statistics.

10 Leading Causes of Death, United States

2022, All Deaths with drilldown to ICD codes, Both Sexes, All Races, All Ethnicities



NEW YORK INSTITUTE OF TECHNOLOGY

College of Osteopathic
Medicine

Ten Leading Causes of Death By Age Group, United States, 2022. CENTERS FOR DISEASE CONTROL AND PREVENTION, NATIONAL CENTER FOR INJURY PREVENTION AND CONTROL. WEB-BASED INJURY STATISTICS QUERY AND REPORTING SYSTEM (WISQARS) [ONLINE] AVAILABLE FROM URL: WWW.WISQARS.CDC.GOV

Suicide Data: Arkansas



Suicide is a public health problem and leading cause of death in the United States. Suicide can also be prevented – more investment in suicide prevention, education, and research will prevent the untimely deaths of thousands of Americans each year. Unless otherwise noted, this fact sheet reports 2021 data from the CDC, the most current verified data available at time of publication (January 2024).

11th leading cause of death in Arkansas

3rd leading

cause of death for ages 10-24

2nd leading

cause of death for ages 25-34

5th leading

cause of death for ages 35-44

8th leading

cause of death for ages 45-54

11th leading

cause of death for ages 55-64

18th leading

cause of death for ages 65+

Suicide Death Rates

	Number of Deaths by Suicide	Rate per 100,000 Population	State Rank
Arkansas	618	20.61	11
Nationally	48,183	14.04	

See full list of citations at afsp.org/statistics.

66% of communities did not have enough mental health providers to serve residents in 2023, according to federal guidelines.

Over **three times** as many people died by suicide than in alcohol related motor vehicle accidents.

The total deaths to suicide reflected a total of **12,812 years** of potential life lost (YPLL) before age 65.

56% of firearm deaths were suicides.

63% of all suicides were by firearms.

Suicide Data: New York



Suicide is a public health problem and leading cause of death in the United States. Suicide can also be prevented – more investment in suicide prevention, education, and research will prevent the untimely deaths of thousands of Americans each year. Unless otherwise noted, this fact sheet reports 2021 data from the CDC, the most current verified data available at time of publication (January 2024).

15th leading cause of death in New York

3rd leading

cause of death for ages 10-24

2nd leading

cause of death for ages 25-34

5th leading

cause of death for ages 35-44

6th leading

cause of death for ages 45-54

11th leading

cause of death for ages 55-64

19th leading

cause of death for ages 65+

Suicide Death Rates

	Number of Deaths by Suicide	Rate per 100,000 Population	State Rank
New York	1,660	7.92	49
Nationally	48,183	14.04	

See full list of citations at afsp.org/statistics.

49% of communities did not have enough mental health providers to serve residents in 2023, according to federal guidelines.

Over **four times** as many people died by suicide than in alcohol related motor vehicle accidents.

The total deaths to suicide reflected a total of **32,406 years** of potential life lost (YPLL) before age 65.

41% of firearm deaths were suicides.

26% of all suicides were by firearms.

THE SUICIDAL PATIENT

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

Demographic characteristics

- White
- Male
- Older age (> 55 years) middle age
- Divorced, never married, or widowed
- Unemployed
- Age 15-34 years (young)

THE SUICIDAL PATIENT

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

Patient history characteristics

- Previous suicide attempts, especially with serious intent, lethal means, or disappointment about survival
- Family history of suicide
- Victim of physical or sexual abuse

THE SUICIDAL PATIENT

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

Psychiatric characteristics

- Diagnosis: affective disorder, alcoholism, panic disorder, psychotic disorders, conduct disorder, severe personality disorder (especially antisocial and borderline).
- Symptoms: suicidal or homicidal ideation; depression, especially with hopelessness, helplessness, anhedonia, delusions, agitation; mixed mania and depression; psychotic symptoms, including command hallucinations and persecutory delusions.
- Current use of alcohol or illicit drugs.
- Recent psychiatric hospitalization.

THE SUICIDAL PATIENT

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

Environmental characteristics

- Recent loss (e.g., of a spouse or job)
- Social isolation
- Access to guns or other lethal weapons
- Social acceptance of suicide

Behavioral characteristics

- Antisocial acts
- Poor impulse control, risk taking, and aggressiveness
- Preparing for death (e.g., making a will, giving away possessions, stockpiling lethal medication)
- Well-developed, detailed suicide plan
- Statements of intent to inflict harm on self or others

THE SUICIDAL PATIENT

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- All potentially fatal threats, gestures, and attempts should be taken seriously.
- Be aware of the possibility. If you don't consider the diagnosis, you won't make the diagnosis.
- Be empathic, try to establish a rapport before honing in on the issue of suicide.
- Perform a mental status examination.
- Ask about suicidal thoughts (e.g., ideas, wishes, motives).
- Ask about suicidal intent

THE SUICIDAL PATIENT

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

- Ask about suicide plans. Is there a detailed plan? Are the means available? Are the means lethal? Is there a chance for rescue?
- Are there any realistic plans for the future?
- Why now? Is there a precipitant? Anger? Loss? Helplessness? A social network?
- Information may be difficult to obtain if privacy not assured. However, information from friends or relatives may be essential.
- Review for the presence of risk factors (especially depression, alcoholism, psychosis, medical illness, and a history of prior attempts.)

Columbia-Suicide Severity Rating Scale

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

COLUMBIA-SUICIDE SEVERITY RATING SCALE Screen Version

SUICIDE IDEATION DEFINITIONS AND PROMPTS	Past month	
	YES	NO
Ask questions that are bolded and <u>underlined</u>.		
Ask Questions 1 and 2		
1) Wish to be Dead: Person endorses thoughts about a wish to be dead or not alive anymore, or wish to fall asleep and not wake up. <u>Have you wished you were dead or wished you could go to sleep and not wake up?</u>		
2) Suicidal Thoughts: General non-specific thoughts of wanting to end one's life/commit suicide, "I've thought about killing myself" without general thoughts of ways to kill oneself/associated methods, intent, or plan. <u>Have you actually had any thoughts of killing yourself?</u>		
If YES to 2, ask questions 3, 4, 5, and 6. If NO to 2, go directly to question 6.		
3) Suicidal Thoughts with Method (without Specific Plan or Intent to Act): Person endorses thoughts of suicide and has thought of a least one method during the assessment period. This is different than a specific plan with time, place or method details worked out. "I thought about taking an overdose but I never made a specific plan as to when where or how I would actually do it....and I would never go through with it." <u>Have you been thinking about how you might kill yourself?</u>		
4) Suicidal Intent (without Specific Plan): Active suicidal thoughts of killing oneself and patient reports having <u>some intent to act on such thoughts</u> , as opposed to "I have the thoughts but I definitely will not do anything about them." <u>Have you had these thoughts and had some intention of acting on them?</u>		
5) Suicide Intent with Specific Plan: Thoughts of killing oneself with details of plan fully or partially worked out and person has some intent to carry it out. <u>Have you started to work out or worked out the details of how to kill yourself? Do you intend to carry out this plan?</u>		
6) Suicide Behavior Question: <u>Have you ever done anything, started to do anything, or prepared to do anything to end your life?</u> Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc. If YES, ask: <u>How long ago did you do any of these?</u> • Over a year ago? • Between three months and a year ago? • Within the last three months?		

For inquiries and training information contact: Kelly Posner, Ph.D.
 New York State Psychiatric Institute, 1051 Riverside Drive, New York, New York, 10632; posnerk@nyspi.columbia.edu
 © 2008 The Research Foundation for Mental Hygiene, Inc.

SAFETY PLAN

NEW YORK INSTITUTE OF TECHNOLOGY


College of Osteopathic
Medicine

Patient Safety Plan Template	
Step 1: Warning signs (thoughts, images, mood, situation, behavior) that a crisis may be developing:	
1.	_____
2.	_____
3.	_____
Step 2: Internal coping strategies – Things I can do to take my mind off my problems without contacting another person (relaxation technique, physical activity):	
1.	_____
2.	_____
3.	_____
Step 3: People and social settings that provide distraction:	
1. Name _____	Phone _____
2. Name _____	Phone _____
3. Place _____	4. Place _____
Step 4: People whom I can ask for help:	
1. Name _____	Phone _____
2. Name _____	Phone _____
3. Name _____	Phone _____
Step 5: Professionals or agencies I can contact during a crisis:	
1. Clinician Name _____	Phone _____
Clinician Pager or Emergency Contact # _____	
2. Clinician Name _____	Phone _____
Clinician Pager or Emergency Contact # _____	
3. Local Urgent Care Services _____	
Urgent Care Services Address _____	
Urgent Care Services Phone _____	
4. Suicide Prevention Lifeline Phone: 1-800-273-TALK (8255)	
Step 6: Making the environment safe:	
1.	_____
2.	_____
Safety Plan Template 5/2008 Barbara Stanley and Gregory K. Brown, is reprinted with the express permission of the authors. No portion of this Safety Plan Template may be reproduced without their express, written permission. You can contact the authors at bbs52@cornell.edu or gkbrown@gmail.com at nyinstitute.edu.	
The one thing that is most important to me and worth living for is:	

The Safety Plan Application

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine



Source: Stanley-Brown Safety Plan on the App



Confidentiality

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

1. DEFINITION

In the American Psychiatric Association's Principles of Medical Ethics with Annotations Especially Applicable to Psychiatry, Annotation 1 reads, *"Confidentiality is essential to psychiatric treatment. This is based in part on the special nature of psychiatric therapy as well as on the traditional ethical relationship between physician and patient."*

Confidentially is defined mnemonically as the clinician's obligation to keep information learned in that relationship unavailable to third parties.



Confidentiality (*cont'd*)

NEW YORK INSTITUTE
OF TECHNOLOGY

College of Osteopathic
Medicine

4. Principles to follow when breaching confidentiality is necessary

- Notify patient whenever possible
- Utilize therapeutic alliance
- Encourage patient to inform (e.g., duty to protect)
- Document rationale for action

Physically Approaching the Patient (contd.)

- Be careful about direct eye contact
- Keep your facial expressions neutral, accepting, and attentive.
- Keep your hands to the midline of your body where they will be more ready to ward off a blow, => DO NOT fold them akimbo or at your waist <= Confrontational
- Avoid easy striking range. At least 1 ½ arms length away so patient will have to move toward you to strike.
- =>Never turn your back and stay near the door or an escape route.
- Attitude toward punishment or other negative consequences
- Degree to which violence is normal in current social circle
- Suicidal thoughts or plans

Safe and Supportive Stance

1. Safest shoulder side
2. Distancing
3. Front foot at 12:00 position; back foot at 10:00 or 2:00
4. Blading
5. Knees slightly bent
6. Hands above belt line

12:00



2:00



Verbally Approaching the Patient

**NEW YORK INSTITUTE
OF TECHNOLOGY**

College of Osteopathic
Medicine

- Interview in a calm, direct reassuring manner
- Refer to patients by surname with title
- Explain your own behavior
- Keep questions, requests and comments very simple, brief, and repeat information to patient several times.
- Constantly clarify that talking and feeling are acceptable that acting violent is not.
- Make it clear patient will be held responsible for his or her actions.
- Acknowledge patient's ability to harm others, then establish your position that this would not be acceptable and that the staff will act to prevent harm to the patient or to others.
- Do not let this sound like a threat but state it as a matter of fact and balance it with a statement that you will do what you can to help.