

Approximating the Score Function using Optimal Transport

Curves and Surfaces 2026

Quentin Giton

Laboratoire de Mathématiques d'Orsay

June 11th

(Badly and Slowly) Approximating the Score Function using Optimal Transport


Curves and Surfaces 2026

Quentin Giton

Laboratoire de Mathématiques d'Orsay

June 11th


Quick reminders on Optimal Transport 1/3



- ▶ Given ground cost $c(x, y)$
- ▶ Among all maps T satisfying $\forall B, \nu(B) = \mu(T^{-1}(B))$
- ▶ Find the one minimising the average transport cost

$$\int_{\mathbb{R}^d} c(x, T(x)) \mu(dx)$$


Quick reminders on Optimal Transport 1/3



- ▶ Given ground cost $c(x, y)$
- ▶ Among all maps T satisfying $\forall B, \nu(B) = \mu(T^{-1}(B))$
Among all **transport plans** π with marginals μ and ν
- ▶ Find the one minimising the average transport cost

$$\int_{\mathbb{R}^d} c(x, T(x)) \mu(dx) \longrightarrow \int_{\mathbb{R}^d \times \mathbb{R}^d} c(x, y) \pi(dx, dy)$$


Quick reminders on Optimal Transport 2/3



► p -Wasserstein distance:

$$W_p(\mu, \nu)^p = \inf_{\pi \in \Pi(\mu, \nu)} \int d(x, y)^p \gamma(dx, dy)$$

Quick reminders on Optimal Transport 2/3



- p -Wasserstein distance:


$$W_p(\mu, \nu)^p = \inf_{\pi \in \Pi(\mu, \nu)} \int d(x, y)^p \gamma(dx, dy)$$

- Kantorovich duality

$$\inf_{\pi \in \Pi(\mu, \nu)} \int c(x, y) \gamma(dx, dy) = \sup_{\phi} \int \phi(x) \mu(dx) + \int \phi^c(y) \nu(dy),$$

where $\phi^c(y) := \inf_x c(x, y) - \phi(x)$.


Quick reminders on Optimal Transport 3/3



► Kantorovich duality

$$T : \inf_{T_{\#}\mu=\nu} \int_{\mathbb{R}^d} c(x, T(x)) \mu(dx) = \sup_{\phi} \int \phi(x) \mu(dx) + \int \phi^c(y) \nu(dy)$$

Quick reminders on Optimal Transport 3/3



► Kantorovich duality

$$T : \inf_{\# \mu = \nu} \int_{\mathbb{R}^d} c(x, T(x)) \mu(dx) = \sup_{\phi} \int \phi(x) \mu(dx) + \int \phi^c(y) \nu(dy)$$

► If an optimal transport map T^* exists, there also exists an optimal Kantorovich potential ϕ^* and

$$y = T^*(x) = \nabla \left(\frac{\|\cdot\|^2}{2} - \phi^* \right) (x) = x - \nabla \phi^*(x),$$

hence $-\nabla \phi^*(x) = T^*(x) - x = y - x$ is the **displacement** of the particle x .

What is the score function?

- **Forward process:** Ornstein–Uhlenbeck (OU) process:

$$d\vec{X}_t = -\vec{X}_t dt + \sqrt{2}d\omega_t, \quad \vec{X}_0 \sim \mu, \quad t \in [0, T].$$

Let $\vec{\rho}_t := \text{Law}(\vec{X}_t)$ and $\gamma \propto e^{-\|\cdot\|^2/2}$. Then $\vec{\rho}_t$ solves the Fokker–Planck equation

$$\partial_t \vec{\rho}_t(x) = \Delta \vec{\rho}_t(x) + \text{div}(\vec{\rho}_t(x)x).$$

whose stationary solution is γ . The process $(\vec{X}_t)_t$ is a **diffusive** process.

What is the score function?

- **Forward process:** Ornstein–Uhlenbeck (OU) process:

$$d\vec{X}_t = -\vec{X}_t dt + \sqrt{2}d\omega_t, \quad \vec{X}_0 \sim \mu, \quad t \in [0, T].$$

Let $\vec{\rho}_t := \text{Law}(\vec{X}_t)$ and $\gamma \propto e^{-\|\cdot\|^2/2}$. Then $\vec{\rho}_t$ solves the Fokker–Planck equation

$$\partial_t \vec{\rho}_t(x) = \Delta \vec{\rho}_t(x) + \text{div}(\vec{\rho}_t(x)x).$$

whose stationary solution is γ . The process $(\vec{X}_t)_t$ is a **diffusive** process.

- **Backward process:** $\tilde{X}_s := \vec{X}_{T-s}$ for $s \in [0, T]$. If $(\vec{X}_t)_{t \in [0, T]}$ is an OU process, then its time reversal $(\tilde{X}_s)_{s \in [0, T]}$ solves

$$d\tilde{X}_s = \left[\tilde{X}_s + 2 \underbrace{\nabla \ln(\vec{\rho}_{T-s}(\tilde{X}_s))}_{\text{The score function}} \right] ds + \sqrt{2}d\omega_s, \quad \tilde{X}_0 \sim \vec{\rho}_T.$$

Using OT to learn the score 1/2

- The Fokker–Planck equation is the 2-Wasserstein Gradient Flow (WGF) of the relative entropy $\mathcal{H}(\rho|\gamma) := \int V\rho + \int \rho \ln(\rho)$. That is:

$$\partial_t \rho_t(x) = \Delta \rho_t(x) + \operatorname{div}(\rho_t(x)x) = -\nabla \cdot (\rho_t(x)v_t(x)),$$

where $v_t(x) = -x - \nabla \ln(\rho_t(x))$.

Using OT to learn the score 1/2

- ▶ The Fokker–Planck equation is the 2-Wasserstein Gradient Flow (WGF) of the relative entropy $\mathcal{H}(\rho|\gamma) := \int V\rho + \int \rho \ln(\rho)$. That is:

$$\partial_t \rho_t(x) = \Delta \rho_t(x) + \operatorname{div}(\rho_t(x)x) = -\nabla \cdot (\rho_t(x)v_t(x)),$$

where $v_t(x) = -x - \nabla \ln(\rho_t(x))$.

- ▶ Given a time-step $\tau > 0$, such a gradient flow can be discretized in time using the **JKO scheme**

$$\begin{cases} \rho_0 = \mu \\ \rho_{n+1}^\tau := \arg \min \mathcal{H}(\rho|\gamma) + \frac{1}{2\tau} W_2(\rho_n^\tau, \rho)^2 \end{cases}$$

Using OT to learn the score 2/2

► Solving

$$\inf_{\rho} \mathcal{H}(\rho|\gamma) + \frac{1}{2\tau} W_2(\rho_n^\tau, \rho)^2$$

amounts to finding the optimal transport map between

$$\rho_n^\tau \quad \text{and} \quad \rho_{n+1}^\tau \propto \exp\left(-\phi^{c/2\tau}\right) d\gamma,$$

where ϕ solves

$$\inf_{\phi} \int e^{-\phi^{c/2\tau}} d\gamma - \int \phi d\rho_n^\tau.$$

Using OT to learn the score 2/2

► Solving

$$\inf_{\rho} \mathcal{H}(\rho|\gamma) + \frac{1}{2\tau} W_2(\rho_n^\tau, \rho)^2$$

amounts to finding the optimal transport map between

$$\rho_n^\tau \quad \text{and} \quad \rho_{n+1}^\tau \propto \exp\left(-\phi^{c/2\tau}\right) d\gamma,$$

where ϕ solves

$$\inf_{\phi} \int e^{-\phi^{c/2\tau}} d\gamma - \int \phi d\rho_n^\tau.$$

► The optimal potential $\phi^\star$ satisfies

$$-\nabla \phi^\star(x) = \nabla(\phi^\star)^{c/2\tau}(y) = y - x.$$

Using OT to learn the score 2/2

- Solving

$$\inf_{\rho} \mathcal{H}(\rho|\gamma) + \frac{1}{2\tau} W_2(\rho_n^\tau, \rho)^2$$

amounts to finding the optimal transport map between

$$\rho_n^\tau \quad \text{and} \quad \rho_{n+1}^\tau \propto \exp\left(-\phi^{c/2\tau}\right) d\gamma,$$

where ϕ solves

$$\inf_{\phi} \int e^{-\phi^{c/2\tau}} d\gamma - \int \phi d\rho_n^\tau.$$

- The optimal potential ϕ^* satisfies

$$-\nabla\phi^*(x) = \nabla(\phi^*)^{c/2\tau}(y) = y - x.$$

- The displacement of the particle x being $v_t(x)$, we obtain

$$-\nabla\phi^*(x) = v_t(x) = -x - \nabla\ln(\rho_{n+1}^\tau(x)) \iff \boxed{\nabla\ln(\rho_{n+1}^\tau(x)) = x + \nabla\phi^*(x)}.$$