

Approximating the Score Function using Optimal Transport

Curves and Surfaces 2026

Quentin Giton

Laboratoire de Mathématiques d'Orsay

June 11th


 (Badly and Slowly) Approximating the Score Function
using Optimal Transport
Curves and Surfaces 2026

Quentin Giton

Laboratoire de Mathématiques d'Orsay

June 11th


Quick reminders on Optimal Transport 1/3



- ▶ Given ground cost $c(x, y)$
- ▶ Among all maps T satisfying $\forall B, \nu(B) = \mu(T^{-1}(B))$
- ▶ Find the one minimising the average transport cost

$$\int_{\mathbb{R}^d} c(x, T(x)) \mu(dx)$$


Quick reminders on Optimal Transport 1/3



- ▶ Given ground cost $c(x, y)$
- ▶ Among all maps T satisfying $\forall B, \nu(B) = \mu(T^{-1}(B))$
Among all **transport plans** π with marginals μ and ν
- ▶ Find the one minimising the average transport cost

$$\int_{\mathbb{R}^d} c(x, T(x)) \mu(dx) \longrightarrow \int_{\mathbb{R}^d \times \mathbb{R}^d} c(x, y) \pi(dx, dy)$$


Quick reminders on Optimal Transport 2/3



► p -Wasserstein distance:

$$W_p(\mu, \nu)^p = \inf_{\pi \in \Pi(\mu, \nu)} \int d(x, y)^p \gamma(dx, dy)$$

Quick reminders on Optimal Transport 2/3



- p -Wasserstein distance:


$$W_p(\mu, \nu)^p = \inf_{\pi \in \Pi(\mu, \nu)} \int d(x, y)^p \gamma(dx, dy)$$

- Kantorovich duality

$$\inf_{\pi \in \Pi(\mu, \nu)} \int c(x, y) \gamma(dx, dy) = \sup_{\phi} \int \phi(x) \mu(dx) + \int \phi^c(y) \nu(dy),$$

where $\phi^c(y) := \inf_x c(x, y) - \phi(x)$.


Quick reminders on Optimal Transport 3/3



► Kantorovich duality

$$\inf_{T : T_{\#}\mu = \nu} \int_{\mathbb{R}^d} c(x, T(x)) \mu(dx) = \sup_{\phi} \int \phi(x) \mu(dx) + \int \phi^c(y) \nu(dy)$$

Quick reminders on Optimal Transport 3/3



- Kantorovich duality

$$\inf_{T: T_{\#}\mu=\nu} \int_{\mathbb{R}^d} c(x, T(x)) \mu(dx) = \sup_{\phi} \int \phi(x) \mu(dx) + \int \phi^c(y) \nu(dy)$$

- If an optimal transport map T^* exists, there also exists an optimal Kantorovich potential ϕ^* and

$$y = T^*(x) = \nabla \left(\frac{\|\cdot\|^2}{2} - \phi^* \right) (x) = x - \nabla \phi^*(x),$$

hence $-\nabla \phi^*(x) = T^*(x) - x = y - x$ is the **displacement** of the particle x .

What is the score function?

- **Forward process:** Ornstein–Uhlenbeck (OU) process:

$$d\vec{X}_t = -\vec{X}_t dt + \sqrt{2}d\omega_t, \quad \vec{X}_0 \sim \mu, \quad t \in [0, T].$$

Let $\vec{\rho}_t := \text{Law}(\vec{X}_t)$ and $\gamma \propto e^{-\|\cdot\|^2/2}$. Then $\vec{\rho}_t$ solves the Fokker–Planck equation

$$\partial_t \vec{\rho}_t(x) = \Delta \vec{\rho}_t(x) + \text{div}(\vec{\rho}_t(x)x).$$

whose stationary solution is γ . The process $(\vec{X}_t)_t$ is a **diffusive** process.

What is the score function?

- **Forward process:** Ornstein–Uhlenbeck (OU) process:

$$d\vec{X}_t = -\vec{X}_t dt + \sqrt{2}d\omega_t, \quad \vec{X}_0 \sim \mu, \quad t \in [0, T].$$

Let $\vec{\rho}_t := \text{Law}(\vec{X}_t)$ and $\gamma \propto e^{-\|\cdot\|^2/2}$. Then $\vec{\rho}_t$ solves the Fokker–Planck equation

$$\partial_t \vec{\rho}_t(x) = \Delta \vec{\rho}_t(x) + \text{div}(\vec{\rho}_t(x)x).$$

whose stationary solution is γ . The process $(\vec{X}_t)_t$ is a **diffusive** process.

- **Backward process:** $\tilde{X}_s := \vec{X}_{T-s}$ for $s \in [0, T]$. If $(\vec{X}_t)_{t \in [0, T]}$ is an OU process, then its time reversal $(\tilde{X}_s)_{s \in [0, T]}$ solves

$$d\tilde{X}_s = \left[\tilde{X}_s + 2 \underbrace{\nabla \ln \left(\vec{\rho}_{T-s}(\tilde{X}_s) \right)}_{\text{The score function}} \right] ds + \sqrt{2}d\omega_s, \quad \tilde{X}_0 \sim \vec{\rho}_T.$$

Using OT to learn the score 1/2

- The Fokker–Planck equation is the 2-Wasserstein Gradient Flow (WGF) of the relative entropy $\mathcal{H}(\rho|\gamma) := \int V\rho + \int \rho \ln(\rho)$. That is:

$$\partial_t \rho_t(x) = \Delta \rho_t(x) + \operatorname{div}(\rho_t(x)x) = -\nabla \cdot (\rho_t(x)v_t(x)),$$

where $v_t(x) = -x - \nabla \ln(\rho_t(x))$.

Using OT to learn the score 1/2

- ▶ The Fokker–Planck equation is the 2-Wasserstein Gradient Flow (WGF) of the relative entropy $\mathcal{H}(\rho|\gamma) := \int V\rho + \int \rho \ln(\rho)$. That is:

$$\partial_t \rho_t(x) = \Delta \rho_t(x) + \operatorname{div}(\rho_t(x)v_t(x)) = -\nabla \cdot (\rho_t(x)v_t(x)),$$

where $v_t(x) = -x - \nabla \ln(\rho_t(x))$.

- ▶ Given a time-step $\tau > 0$, such a gradient flow can be discretised in time using the **JKO scheme**

$$\begin{cases} \rho_0 = \mu \\ \rho_{n+1}^\tau := \arg \min \mathcal{H}(\rho|\gamma) + \frac{1}{2\tau} W_2(\rho_n^\tau, \rho)^2 \end{cases}$$

Using OT to learn the score 2/2

► Solving

$$\inf_{\rho} \mathcal{H}(\rho|\gamma) + \frac{1}{2\tau} W_2(\rho_n^\tau, \rho)^2$$

amounts to finding the optimal transport map between

$$\rho_n^\tau \quad \text{and} \quad \rho_{n+1}^\tau \propto \exp\left(-\phi^{c/2\tau}\right) d\gamma,$$

where ϕ solves

$$\inf_{\phi} \tilde{J}(\phi) := \ln\left(\int e^{-\phi^{c/2\tau}} d\gamma\right) - \int \phi d\rho_n^\tau.$$

Using OT to learn the score 2/2

► Solving

$$\inf_{\rho} \mathcal{H}(\rho|\gamma) + \frac{1}{2\tau} W_2(\rho_n^{\tau}, \rho)^2$$

amounts to finding the optimal transport map between

$$\rho_n^{\tau} \quad \text{and} \quad \rho_{n+1}^{\tau} \propto \exp\left(-\phi^{c/2\tau}\right) d\gamma,$$

where ϕ solves

$$\inf_{\phi} \tilde{J}(\phi) := \ln\left(\int e^{-\phi^{c/2\tau}} d\gamma\right) - \int \phi d\rho_n^{\tau}.$$

► The optimal potential $\phi^{\star}$ satisfies

$$-\nabla\phi^{\star}(x) = \nabla(\phi^{\star})^{c/2\tau}(y) = \frac{y - x}{\tau}.$$

Using OT to learn the score 2/2

- Solving

$$\inf_{\rho} \mathcal{H}(\rho|\gamma) + \frac{1}{2\tau} W_2(\rho_n^\tau, \rho)^2$$

amounts to finding the optimal transport map between

$$\rho_n^\tau \quad \text{and} \quad \rho_{n+1}^\tau \propto \exp\left(-\phi^{c/2\tau}\right) d\gamma,$$

where ϕ solves

$$\inf_{\phi} \tilde{J}(\phi) := \ln\left(\int e^{-\phi^{c/2\tau}} d\gamma\right) - \int \phi d\rho_n^\tau.$$

- The optimal potential $\phi^\star$ satisfies

$$-\nabla\phi^\star(x) = \nabla(\phi^\star)^{c/2\tau}(y) = \frac{y - x}{\tau}.$$

- The displacement of the particle x being $v_t(x)$, we obtain

$$-\nabla\phi^\star(x) = v_t(x) = -x - \nabla\ln(\rho_{n+1}^\tau(x)) \iff \boxed{\nabla\ln(\rho_{n+1}^\tau(x)) = \nabla\phi^\star(x) - x}.$$

General contributions

Proposition (G.)

The functional $\tilde{J}(\phi) := \ln \left(\int e^{-\phi^{c/2\tau}} d\gamma \right) - \int \phi d\rho_n^\tau$ is convex.

General contributions

Proposition (G.)

The functional $\tilde{J}(\phi) := \ln \left(\int e^{-\phi^{c/2\tau}} d\gamma \right) - \int \phi d\rho_n^\tau$ is convex.

Theorem (G.)

Let

$$J(\phi) := \int e^{-\phi^{c/2\tau}} d\gamma - \int \phi d\rho_n^\tau.$$

Then,

$$\arg \min_{\phi} J(\phi) \subseteq \arg \min_{\phi} \tilde{J}(\phi) \quad \text{and} \quad \arg \min_{\int e^{-\phi^{c/2\tau}} d\gamma = 1} \tilde{J}(\phi) = \arg \min_{\phi} J(\psi).$$

Stochastic gradient descent 1/2

$$\blacktriangleright J(\phi) = \mathbb{E}_{\mathbf{y} \sim \gamma} \left[\exp \left(-\phi^{c/2\tau}(\mathbf{y}) \right) - \int \phi(\mathbf{x}) \rho_n^\tau(\mathrm{d}\mathbf{x}) \right]$$

Stochastic gradient descent 1/2

- ▶ $J(\phi) = \mathbb{E}_{\mathbf{y} \sim \gamma} \left[\exp \left(-\phi^{c/2\tau}(\mathbf{y}) \right) - \int \phi(\mathbf{x}) \rho_n^\tau(\mathrm{d}\mathbf{x}) \right]$
- ▶ Choice of parametrisation: semi-discrete optimal transport using

$$\hat{\rho}_n^\tau = \frac{1}{N} \sum_{i=1}^N \delta_{\hat{\mathbf{x}}_i^{(n)}}$$

Stochastic gradient descent 1/2

- ▶ $J(\phi) = \mathbb{E}_{\mathbf{y} \sim \gamma} \left[\exp \left(-\phi^{c/2\tau}(\mathbf{y}) \right) - \int \phi(\mathbf{x}) \rho_n^\tau(\mathrm{d}\mathbf{x}) \right]$
- ▶ Choice of parametrisation: semi-discrete optimal transport using

$$\hat{\rho}_n^\tau = \frac{1}{N} \sum_{i=1}^N \delta_{\hat{\mathbf{x}}_i^{(n)}}$$

- ▶ $G(\theta) = \mathbb{E}_{\mathbf{y} \sim \gamma} \left[\max_{1 \leq i \leq N} \left\{ \exp \left(\theta_i - \frac{c(\hat{\mathbf{x}}_i^{(n)}, \mathbf{y})}{2\tau} \right) \right\} - \frac{1}{N} \sum_{i=1}^N \theta_i \right]$

Stochastic gradient descent 1/2

- ▶ $J(\phi) = \mathbb{E}_{\mathbf{y} \sim \gamma} \left[\exp \left(-\phi^{c/2\tau}(\mathbf{y}) \right) - \int \phi(\mathbf{x}) \rho_n^\tau(\mathrm{d}\mathbf{x}) \right]$
- ▶ Choice of parametrisation: semi-discrete optimal transport using

$$\hat{\rho}_n^\tau = \frac{1}{N} \sum_{i=1}^N \delta_{\hat{\mathbf{x}}_i^{(n)}}$$

- ▶ $G(\theta) = \mathbb{E}_{\mathbf{y} \sim \gamma} \left[\max_{1 \leq i \leq N} \left\{ \exp \left(\theta_i - \frac{c(\hat{\mathbf{x}}_i^{(n)}, \mathbf{y})}{2\tau} \right) \right\} - \frac{1}{N} \sum_{i=1}^N \theta_i \right]$
- ▶ Projected SGD algorithm ($c_\tau(\mathbf{x}, \mathbf{y}) := \frac{c(\mathbf{x}, \mathbf{y})}{2\tau}$)

$$\left\{ \begin{array}{l} \text{pick } \mathbf{y} \sim \gamma \\ \text{let } i := \arg \min_j c_\tau(\hat{\mathbf{x}}_j^{(n)}, \mathbf{y}) - \theta_j \\ \text{define } \mathbf{e}_i \text{ as the } i\text{-th canonical basis vector of } \mathbb{R}^N \\ \theta^{(k+1)} := \text{proj}_{B(0, R_{\text{proj}})} \left(\theta^{(k)} - \alpha_k e^{\theta_i^{(k)} - c_\tau(\hat{\mathbf{x}}_j^{(n)}, \mathbf{y})} \mathbf{e}_i + \frac{\alpha_k}{N} \mathbf{1} \right) \end{array} \right.$$

Stochastic gradient descent 2/2

Theorem (G .)

Let $(\theta^{(k)})_{k=0}^K$ be the iterates obtained with step-sizes $\alpha_k = \frac{R_{\text{proj}}}{B\sqrt{k}}$ starting from some initial $\theta^{(0)} \in B(0, R_{\text{proj}})$. Define $\bar{\theta}_K$ to be the arithmetic mean of all iterates up to time K . Then, it holds that

$$0 \leq \mathbb{E}[G(\bar{\theta}_K)] - G(\theta^*) \leq BR_{\text{proj}} \left(\frac{2}{K} + \frac{1}{\sqrt{K}} \right) = \mathcal{O}(K^{-\frac{1}{2}}),$$

where θ^* denotes the unique minimiser of G .

Approximating the next JKO iterate

- Recall that the next measure is

$$\rho_{n+1}^\tau(dy) \propto \max_i e^{\theta_i - c_\tau(\hat{x}_i^{(n)}, y)} \gamma(dy)$$

Approximating the next JKO iterate

- Recall that the next measure is

$$\rho_{n+1}^\tau(dy) \propto \max_i e^{\theta_i - c_\tau(\hat{x}_i^{(n)}, y)} \gamma(dy)$$

- It is approximated by

$$\rho_{n+1}^\tau(dy) \propto \max_i e^{[\bar{\theta}_K]_i - c_\tau(\hat{x}_i^{(n)}, y)} \gamma(dy)$$

Approximating the next JKO iterate

- Recall that the next measure is

$$\rho_{n+1}^\tau(dy) \propto \max_i e^{\theta_i - c_\tau(\hat{x}_i^{(n)}, y)} \gamma(dy)$$

- It is approximated by

$$\rho_{n+1}^\tau(dy) \propto \max_i e^{[\bar{\theta}_K]_i - c_\tau(\hat{x}_i^{(n)}, y)} \gamma(dy)$$

- We discretise it by

$$\hat{x}_i^{(n+1)} :=$$