


ORNL DAAC
DISTRIBUTED ACTIVE ARCHIVE CENTER
FOR BIOGEOCHEMICAL DYNAMICS





About Us

Get Data

Submit Data

Tools

Resources

Help

Sign in

[DAAC Home](#) >
[Get Data](#) >
[NASA Projects](#) >
[North American Carbon Program \(NACP\)](#) >
User guide

Methane Emissions from Dairy Sources (Vista-CA), State of California, USA, 2019

Get Data

Documentation Revision Date: 2020-09-23

Dataset Version: 1

Summary

This dataset provides estimates of methane (CH₄) emissions from dairies in California at a resolution of 0.1 degrees (~ 10 km x 10 km) for the year 2019. The mapped sources of dairy CH₄ emissions are enteric fermentation and manure management reported in gigagrams per year (Gg/yr). The sum of the two sources is also provided. These data are in the succession of Vista California (Vista-CA) spatial datasets that have identified and classified potential methane source emitters in California and were created utilizing an assortment of publicly available data sources from local, state, and federal agencies. This dataset can serve as a planning tool for mitigation, a prior for atmospheric observation-based emissions estimates, attribution of emissions to a specific facility, and to validate CH₄ emissions reductions from management changes.

The locations of 1,727 dairy farms in California were determined by inspection of Google Earth satellite imagery. Enteric emissions at each dairy were derived from the number of cattle, dry matter intake, neutral detergent fiber in the diet, and milkfat at the dairy. Estimates of herd numbers and demographic categories came from three sources, the Regional Water Quality Control Board, San Joaquin Valley Air Pollution Control Board, and the US Department of Agriculture National Agricultural Statistics Survey. Manure management emissions were estimated by applying equations of the California Air Resources Board to the facility level with data from the San Joaquin Valley Air Pollution Control District permits, local animal management data, and regional differences in manure management.

There are 3 files in GeoTIFF (*.tif) format.




Figure 1. The sum of manure management and enteric fermentation methane emissions (in gigagrams per year) from dairy farms in California. Source: total-dairy_ch4_CA_2019.tif

Citation

Marklein, A.R., D. Meyer, M.L. Fischer, S. Jeong, T. Rafiq, M. Carr, and F.M. Hopkins. 2020. Methane Emissions from Dairy Sources (Vista-CA), State of California, USA, 2019. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAAC/1814>

Table of Contents

1. Dataset Overview
2. Data Characteristics
3. Application and Derivation
4. Quality Assessment
5. Data Acquisition, Materials, and Methods
6. Data Access
7. References

1. Dataset Overview

This dataset provides estimates of methane (CH_4) emissions from dairies in California at a resolution of 0.1 degrees ($\sim 10 \text{ km} \times 10 \text{ km}$) for the year 2019. The mapped sources of dairy CH_4 emissions are enteric fermentation and manure management reported in gigagrams per year (Gg/yr). The sum of the two sources is also provided. These data are in the succession of Vista California (Vista-CA) spatial datasets that have identified and classified potential methane source emitters in California and were created utilizing an assortment of publicly available data sources from local, state, and federal agencies. This dataset can serve as a planning tool for mitigation, a prior for atmospheric observation-based emissions estimates, attribution of emissions to a specific facility, and to validate CH_4 emissions reductions from management changes.

The locations of 1,727 dairy farms in California were determined by inspection of Google Earth satellite imagery. Enteric emissions at each dairy were derived from the number of cattle, dry matter intake, neutral detergent fiber in the diet, and milk fat at the dairy. Estimates of herd numbers and demographic categories came from three sources, the Regional Water Quality Control Board, San Joaquin Valley Air Pollution Control Board, and the US Department of Agriculture National Agricultural Statistics Survey. Manure

management emissions were estimated by applying equations of the California Air Resources Board to the facility level with data from the San Joaquin Valley Air Pollution Control District permits, local animal management data, and regional differences in manure management.

Project: [North American Carbon Program](#)

The North American Carbon Program (NACP) is a multidisciplinary research program designed to improve understanding of North America's carbon sources, sinks, and stocks. The central objective is to measure and understand the sources and sinks of Carbon Dioxide (CO₂), Methane (CH₄), and Carbon Monoxide (CO) in North America and adjacent oceans. The NACP is supported by a number of different federal agencies.

Related Publication:

Marklein, A. R., Meyer, D., Fischer, M. L., Jeong, S., Rafiq, T., Carr, M., and Hopkins, F. M. 2020. Facility scale inventory of dairy methane emissions in California: Implications for mitigation, Earth Syst. Sci. Data Discuss., <https://doi.org/10.5194/essd-2020-133>, in review.

Related Datasets:

Carranza, V., T. Rafiq, I. Frausto-Vicencio, F. Hopkins, K.R. Verhulst, P. Rao, R.M. Duren, and C.E. Miller. 2018. Sources of Methane Emissions (Vista-LA), South Coast Air Basin, California, USA. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1525>

Hopkins, F.M., T. Rafiq, and R.M. Duren. 2019. Sources of Methane Emissions (Vista-CA), State of California, USA. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1726>

Thorpe, A.K., B.D. Bue, D.R. Thompson, and R.M. Duren. 2019. Methane Plumes Derived from AVIRIS-NG over Point Sources across California, 2016-2017. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1727>

Acknowledgments:

This work was supported by the University of California Office of the President (grant LFR-18-548581) and NASA's Advancing Collaborative Connections for Earth System Science (ACCESS) Methane Source Finder (grant 15-ACCESS15-0034). The authors acknowledge the dairy farmers who provided information to the permits and reports, as well as the San Joaquin Valley and Santa Ana Air Quality Control Boards, California Integrated Water Quality System, and Regional Water Quality Control Boards.

2. Data Characteristics

Spatial Coverage: California, USA

Spatial Resolution: 0.1 degree (10 km)

Temporal Coverage: 2019-01-01 to 2019-12-31

Temporal Resolution: One-time annual estimate

Study Area: Latitude and longitude are given in decimal degrees.

Site	Northernmost Latitude	Southernmost Latitude	Easternmost Longitude	Westernmost Longitude
California	42.05	32.55	-114.15	-124.45

Data File Information

There are 3 files in GeoTIFF (*.tiff) format (Table 1).

Table 1. File names and descriptions.

File Names	Units	Description
enteric-dairy_ch4_CA_2019.tif	Gg/yr	enteric fermentation methane (CH ₄) emissions (in gigagrams per year) from dairy farms in California
manure-dairy_ch4_CA_2019.tif	Gg/yr	manure management methane (CH ₄) emissions (in gigagrams per year) from dairy farms in California
total-dairy_ch4_CA_2019.tif	Gg/yr	the sum of manure management and enteric fermentation methane (CH ₄) emissions (in gigagrams per year) from dairy farms in California

Data File Details

For all files:

- The data type is Float64.
- Missing values are represented by -9999.
- Each file contains 95 rows and 103 columns.
- The projection used is "WGS 84" (EPSG:4326).

3. Application and Derivation

Dairies emit roughly half of total CH₄ emissions in California, generating CH₄ from enteric fermentation by ruminant gut microbes and anaerobic decomposition of manure. Representation of these emission processes is essential for the management and mitigation of CH₄ emissions and is typically done using standardized emission factors applied at large spatial scales (e.g., state-level). However, CH₄-emitting activities and management decisions vary across facilities, and current inventories do not have a sufficiently high spatial resolution to capture changes at this scale. In addition to serving as a planning tool for mitigation, this dataset is useful as a prior for atmospheric observation-based emissions estimates, attribution of emissions to a specific facility, and to validate CH₄ emissions reductions from management changes.

4. Quality Assessment

To estimate facility-scale uncertainty for enteric fermentation emissions, the standard error in dry matter intake, neutral detergent fiber, milkfat was calculated. The facility-scale uncertainty in manure management emissions was estimated by propagating uncertainty in terms of the number of cows, the fraction of time on concrete, volatile solids production, the methane conversion factor (MCF), and the fraction of material used as bedding. The standard error of each variable was determined through the emissions calculations equations, assuming the errors were uncorrelated.

5. Data Acquisition, Materials, and Methods

Vista-California

Vista-CA spatial datasets identify and classify potential methane source emitters in California. Vista-CA spatial datasets were created utilizing an assortment of publicly available data sources ranging from local, state, and federal agencies. See Marklein et al. (2020) for more details.

Methane Emissions from Dairies

This data set provides spatial data products with enteric fermentation and manure methane emissions from dairies in California. Dairies emit roughly half of total methane (CH₄) emissions in California, generating CH₄ from both enteric fermentation by ruminant gut microbes and anaerobic decomposition of manure. Representation of these emission processes is essential for management and mitigation of CH₄ emissions and is typically done using standardized emission factors applied at large spatial scales (e.g., state-level). However, CH₄-emitting activities and management decisions vary across facilities, and current inventories do not have sufficiently high spatial resolution to capture changes at this scale. To create this data set, we disaggregate the California Air Resources Board (CARB) inventory to the facility level by 1) developing a spatially-explicit map of dairy locations, 2) applying facility-level information from regulatory permit data and county-level animal inventories to estimate herd sizes; and 3) estimating enteric and manure CH₄ emissions from dairy facilities based on manure management from permit data and regional norms.

Data Sources

Google Earth satellite imagery was used to determine the locations of 1,727 dairy farms in California, by identifying metal-topped shelters alongside manure lagoons and corrals (Duren et al., 2019).

Data from three sources were used to estimate herd numbers and demographic categories at each dairy. First, the Regional Water Quality Control Board reports provided the number of milk cows, dry cows, heifers, and calves for dairies in the Central Valley and Southern California for the year 2005 (RWQCB, 2013). Second, San Joaquin Valley Air Pollution Control Board permits include the maximum number of cattle in each class at a given facility, based on facility housing in 2011, rather than the number of animals (SJVAPCD, 2004). Third, the 2017 US Department of Agriculture (USDA) National Agricultural Statistics Survey (NASS) provided the number of farms and the number of cows in different dairy size classes in each county (USDA NASS, 2017).

Estimates of enteric fermentation CH₄ emissions were based on calculations by Appuhamy et al. (2019). Enteric emissions included the number of cattle, dry matter intake, neutral detergent fiber in the diet, and milk fat. Manure management emissions were estimated using equations by the California Air Resources Board to the facility level (CARB, 2014), data from the San Joaquin Valley Air Pollution Control District permits (SJVAPCD, 2004), animal management data (Meyer, 2019), and regional differences in manure management.

Processing

Results were aggregated to a resolution of 0.1 degrees to protect the identity and characteristics of dairy farms. Dairy locations are available in Carranza et al. (2018) (for SCAB) and Hopkins et al. (2019) (state-wide). For more information, refer to Marklein et al. (2020).

Uncertainty Analysis:

Total uncertainty in CH₄ emissions at the facility scale is 14.4%; 84.1% of the uncertainty is due to uncertainty in manure emissions, while 15.9% of the uncertainty is due to enteric emissions. We report the statewide uncertainty in enteric emissions to be 20%. The facility-level standard errors for enteric fermentation we calculated are 35.6%, and is most sensitive to dry matter intake, followed by the number of animals. We report the statewide uncertainty in manure emissions to be 30%. The facility-level standard errors for manure emissions we calculated is 55.4%, and is most sensitive to the lagoon methane conversion factor, followed by the number of animals, and the fraction of manure allocated to bedding.

6. Data Access

These data are available through the Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC).

[Methane Emissions from Dairy Sources \(Vista-CA\), State of California, USA, 2019](#)

Contact for Data Center Access Information:

- E-mail: uso@daac.ornl.gov
- Telephone: +1 (865) 241-3952

7. References

Appuhamy, Ranga, and E. Kebreab. 2018. Characterizing California-specific Cattle Feed Rations and Improve Modeling of Enteric Fermentation for California's Greenhouse Gas Inventory. California Environmental Protection Agency, Air Resources Board, Research Division. <https://ww2.arb.ca.gov/sites/default/files/classic/research/apr/past/16rd001.pdf>

CARB. 2014. California's 2000-2012 Greenhouse Gas Emissions Inventory Technical Support Document, 2014 Edition. California Air Resources Board, Air Quality Planning and Science Division. https://ww3.arb.ca.gov/cc/inventory/pubs/reports/2000_2012/ghg_inventory_00-12_technical_support_document.pdf

RWQCB. 2013. Reissued Waste Discharge Requirements General Order for Existing Milk Cow Dairies (Reissued Dairy General Order). Confined Animal Facilities - Dairy Program Regulations and Requirements, Order R5-2013-0122. Regional Water Quality Control Board, Central Valley. https://www.waterboards.ca.gov/centralvalley/board_decisions/adopted_orders/general_orders/r5-2013-0122.pdf

Carranza, V., T. Rafiq, I. Frausto-Vicencio, F. Hopkins, K.R. Verhulst, P. Rao, R.M. Duren, and C.E. Miller. 2018. Sources of Methane Emissions (Vista-LA), South Coast Air Basin, California, USA. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1525>

Carranza, V., T. Rafiq, I. Frausto-Vicencio, F. Hopkins, K.R. Verhulst, P. Rao, R.M. Duren, and C.E. Miller. 2018. Vista-LA: Mapping methane-emitting infrastructure in the Los Angeles megacity. Earth System Science Data, 10(1):653-676, <https://doi.org/10.5194/essd-10-653-2018>

10-653-2018.

Duren, R.M., A.K. Thorpe, K.T. Foster, T. Rafiq, F.M. Hopkins, V. Yadav, B.D. Bue, D.R. Thompson, S. Conley, N.K. Colombi, C. Frankenberg, I.B. McCubbin, M.L. Eastwood, M. Falk, J.D. Herner, B.E. Croes, R.O. Green, and C.E. Miller. 2019. California's methane super-emitters. *Nature* 575:180-184. <https://doi.org/10.1038/s41586-019-1720-3>

Hopkins, F.M., T. Rafiq, and R.M. Duren. 2019. Sources of Methane Emissions (Vista-CA), State of California, USA. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1726>

Marklein, A.R., D. Meyer, M.L. Fischer, S. Jeong, T. Rafiq, M. Carr, and F.M. Hopkins. 2020 Facility scale inventory of dairy methane emissions in California: Implications for mitigation. Submitted to *Earth System Science Data*.

Meyer, D, J. Heguy, B. Karle, and P. Robinson. 2019. Characterize Physical and Chemical Properties of Manure in California Dairy Systems to Improve Greenhouse Gas Emission Estimates. California Environmental Protection Agency, Air Resources Board. <https://ww2.arb.ca.gov/sites/default/files/classic/research/apr/past/16rd002.pdf>

SJVAPCD. 2004. Best Available Control Technology (BACT) Dairy Operations. San Joaquin Valley Air Pollution Control Board. https://www.valleyair.org/farmpermits/updates/draft_dairy_bact.pdf

USDA NASS. 2017. Census of Agriculture. United States Department of Agriculture National Agricultural Statistics Service. <https://www.nass.usda.gov/Publications/AgCensus/2017/index.php>



Privacy Policy | Feedback | Help

Home

About Us

Mission
Data Use and Citation Policy
User Working Group
Partners

Get Data

Science Themes
NASA Projects
All Datasets

Submit Data

Submit Data Form
Data Scope and Acceptance
Data Authorship Policy
Data Publication Timeline
Detailed Submission
Guidelines

Tools

MODIS
THREDDS
SDAT
Daymet
CARVE Data Viewer
Soil Moisture Visualizer
Land - Water Checker

Resources

Learning
Data Management
News

Contact Us