

# Lab 1 Activity

PSYC 6802

Lab activities are usually much shorter than this. You are not expected to turn in this lab activity, but I do encourage you to attempt solving the questions below to better familiarize yourself with R.

## Basic Functions and Calculations

1. Create an object (1D numeric vector) that contains all the *even numbers* from 1 to 10 (10 included). Name the object **even\_10**.
2. Create an object (1D numeric vector) that contains all the *odd numbers* from 1 to 10. Name the object **odd\_10**.
3. Create an object (1D numeric vector) that contains the multiplication between the respective elements of **even\_10** and **odd\_10**. Name the object **mult\_10**. Additionally, calculate the sum of the elements of the **mult\_10** object.

**HINT:** You can apply mathematical operations to vectors of the same length. Mathematical operations will be performed between the respective elements of each vector.

4. Calculate the *mean* of all of the numbers contained in the **even\_10**, **odd\_10**, and **mult\_10** (so only 1 mean, not 3). Use the **mean()** function for this.

**HINT:** the **mean()** function only takes in one object at a time, maybe you can get creative with the **c()** function?

- 4.1. Calculate the same mean, but do so without using the **mean()** function! The mathematical formula for the mean is  $Mean = \frac{\sum x_i}{n}$ , where the numerator is the sum of all of your values, and the denominator is how many values you have.

**HINT:** there is a function that you can use to count how many elements there are in an object.

**5.** Calculate the *standard deviation* of all of the numbers contained in the **even\_10**, **odd\_10**, and **mult\_10** (so only 1 standard deviation, not 3). You will also need to find the function that calculates the standard deviation.

**5.1.** Calculate the same standard deviation without using the standard deviation function! The mathematical formula for the mean is  $SD = \sqrt{\frac{\sum (x_i - \bar{x})^2}{n-1}}$ , where  $x_i$  represents every single values,  $\bar{x}$  represent the mean. You will also need to find the function that calculates the square root.

**HINT:** here you need to use “()” to tell R the correct order of operations and functions.

**More Practice questions on the next page**

## Importing Data and Subsetting

6. Click [here](https://www.openintro.org/data/index.php?data=mammals) to download the **Mammal\_Sleep.csv** file, import it into R, and name it **dat**. You can find the description of the variables in the data set here (<https://www.openintro.org/data/index.php?data=mammals>). Additionally, there is an extra variable, *primate*, that specifies whether the mammal is a primate or not. Explore the data either visually or with the `str()` function to get a better sense of what you are looking at!

7. The `summary()` function has MANY uses in R (the output is different depending on what object you use as input). When applied to a `data.frame` object, `summary()` calculates some descriptive statistics for numeric variables. Run the following code:

```
sum_tab <- summary(dat)
```

Now, extract *only* the means of the **BrainWt** and **TotalSleep** variables from the **sum\_tab** object.

**HINT:** You can investigate what and how information is stored in the **sum\_tab** object by just running `sum_tab`, which will print all of the stored information. Additionally, note that this is a subsetting problem, so try to identify what the dimensions of the **sum\_tab** object are (looking at the environment may help!).

8. How many animals in the data are primates?

**HINT:** In the lecture we looked at a function for counting elements.

8.1. can you find a way to output *only* the number for primates?

**HINT:** This question has to do with dimensions and subsetting.

9. Can you create a new variable in the **dat** data set that is the proportion of body weight that brain weight takes up? That is, if body weight is 2 and brain weight is 0.2, then brain weight takes up .1 ( $0.2/2 = .1$ , or 10%) of the total body weight. Name the new variable **br\_to\_bd\_weight**.

**HINT 1:** you should be able to calculate the proportion in a really short line of code, a hint for one of the previous questions may help you out!

**HINT 2:** you can create a new variable in a `data.frame` as follows:

```
# this is not runnable code, just a conceptual example
```

```
name_of_data$new_variable <- the variable that you want to add to the data
```