Discovering epistasis between germline mutator alleles in mice

This manuscript (<u>permalink</u>) was automatically generated from <u>quinlan-lab/mutator-epistasis-manuscript@077e8f2</u> on February 22, 2023.

Authors

- Thomas A. Sasani
 - © 0000-0003-2317-1374 · ♥ tomsasani · ♥ tomsasani

 Department of Human Genetics, University of Utah · Funded by Grant XXXXXXXX
- Aaron R. Quinlan
 - © 0000-0003-1756-0859 · **У** aaronquinlan

Department of Human Genetics, University of Utah; Department of Biomedical Informatics, University of Utah

- Kelley Harris [™]
 - © 0000-0003-0302-2523 ⋅ **У** Kelley Harris

Department of Genome Sciences, University of Washington

Abstract

Maintaining genome integrity in the mammalian germline is essential and enormously complex. Hundreds of proteins comprise pathways involved in DNA replication, and hundreds more are mobilized to repair DNA damage [1]. While loss-of-function mutations in any of the genes encoding these proteins might lead to elevated mutation rates, *mutator alleles* have largely eluded detection in mammals.

DNA replication and repair proteins often recognize particular sequence motifs or excise lesions at specific nucleotides. Thus, we might expect that the spectrum of de novo mutations — i.e, the frequency of each individual mutation type (C>T, A>G, etc.) — will differ between genomes that harbor either a mutator or wild-type allele at a given locus. Previously, we used quantitative trait locus (QTL) mapping to discover a mutator allele near the DNA repair gene *Mutyh* that increases the rate of *de novo* C>A germline mutation in a collection of recombinant inbred lines (RILs) known as the BXDs [2,3].

In this study, we developed a new method to detect alleles that affect the mutation spectrum in biparental RILs. By applying this method to mutation data from the BXDs, we confirmed the activity of the germline mutator locus near *Mutyh*, and discovered an additional C>A germline mutator locus on chromosome 6 that overlaps *Ogg1*, a key partner of *Mutyh* in base-excision repair of oxidative DNA damage [4]. Strikingly, BXDs with the mutator allele near *Ogg1* do not exhibit elevated rates of C>A germline mutation unless they also possess the mutator allele near *Mutyh*, but BXDs with both alleles exhibit even higher C>A mutation rates than those with either one alone.

To our knowledge, these new methods for analyzing mutation spectra reveal the first evidence of epistasis between mammalian germline mutator alleles, and may be applicable to mutation data from humans and other model organisms.

Introduction

The germline mutation rate is a fundamental parameter in population genetics, and reflects the complex interplay between DNA replication and repair pathways, exogenous sources of DNA damage, and life-history traits. *Mutator alleles* may explain some of the within- and between-species variation in germline mutation rates [5], but have proven challenging to identify in mammalian genomes.

Germline mutator alleles are difficult to detect for a number of reasons, including the fidelity of germline genome replication and the effects of selection on mutators. On average, humans are born with 30 to 50 single-nucleotide de novo germline mutations per haploid genome [6,7]; in mice, that number is closer to 10 or 15 [8]. Due to the low baseline germline mutation rate in many mammals, it can be challenging to ascertain sequencing data from enough haplotypes to reliably detect those with significantly elevated de novo mutation counts. Moreover, in a population of sufficiently large N_e (effective population size), large-effect mutator alleles will likely be efficiently purged by negative selection. The estimated selection coefficient on a mutator allele is approximately $2s\Delta U$ [9], where s is the mean selective coefficient on a new deleterious mutation and ΔU is the excess number of new deleterious mutations caused by the mutator allele; the product of s and s is multiplied by 2 to account for the average number of generations for which mutator is linked to the excess mutations it causes.

Compared to haplotypes that harbor wild-type alleles at a particular locus, those harboring mutator alleles will likely carry an excess of total germline mutations. Indeed, candidate germline mutator loci have been discovered in human genomes by identifying haplotypes with significantly more derived

alleles than the population mean [10]. However, protein-coding genes involved in DNA replication and repair often recognize particular sequence motifs or excise lesions at specific nucleotides [5]. Thus, we might also expect the spectrum of de novo mutations — that is, the frequency of each individual mutation type (C>T, A>G, etc.) — to differ between genomes that carry either a mutator or wild-type allele at a given locus.

In 2022, we discovered a germline mutator allele in mice by analyzing whole-genome sequencing data from 152 recombinant inbred lines (RILs). Commonly known as the BXDs [3], these RILs were derived from either F2 or advanced intercrosses of C57BL/6J and DBA/2J, two laboratory strains that exhibit significant differences in their germline mutation spectra [11]. Since the BXD RILs were maintained via brother-sister mating for up to 180 generations and housed in a controlled laboratory environment, they were an ideal population for mutator allele discovery. Namely, each line accumulated hundreds or thousands of germline mutations on a nearly-homozygous linear mosaic of parental B and D haplotypes, while the effects of negative selection on new and standing variation were attenuated by strict inbreeding [12]. We used quantitative trait locus (QTL) mapping to identify a locus on chromosome 4 that was strongly associated with the C>A germline mutation rate in the BXDs [2]. The QTL overlapped *Mutyh*, which encodes a protein that normally prevents C>A mutations by repairing oxidative DNA damage [4], and we hypothesized that missense mutations in *Mutyh* were responsible for a 50% increase in the C>A mutation rate between BXDs with either parental haplotype at the QTL [2].

In this study, we developed a new method to detect alleles that affect the mutation spectrum in biparental RILs, and applied it to *de novo* germline mutation data from the BXDs. We assessed its power to detect candidate mutator alleles, re-identified the mutator near *Mutyh*, and discovered compelling evidence of epistasis between two germline mutator alleles that augment the C>A germline mutation rate.

Materials and Methods

Identifying de novo germline mutations in the BXD RILs

The BXD resource currently comprises a total of 152 recombinant inbred lines (RILs). RILs were derived from either F2 or advanced intercrosses, and subsequently inbred by brother-sister mating for up to 180 generations [3]. Previously, we analyzed whole-genome sequencing data from the BXDs and identified candidate *de novo* germline mutations in each line [2]. A detailed description of the methods used for DNA extraction, sequencing, alignment, and variant processing, as well as the characteristics of the *de novo* mutations, are available in a previous manuscript [2].

Briefly, we identified private single-nucleotide mutations in each BXD that were absent from all other RILs, as well as from the C57BL/6J and DBA/2J parents. We required each private variant to be meet the following criteria:

- genotyped as either homozygous or heterozygous for the alternate allele, with at least 90% of sequencing reads supporting the alternate allele
- supported by at least 10 sequencing reads
- Phred-scaled genotype quality of at least 20
- must not overlap regions of the genome annotated as segmental duplications or simple repeats in GRCm38/mm10

must occur on a parental haplotype that was inherited by at least one other BXD at the same locus;
 these other BXDs must be homozygous for the reference allele at the variant site

A new approach to discover germline mutator alleles

Using the existing catalog of *de novo* germline mutations in the BXDs, we developed a new approach to discover loci that affect the germline *de novo* mutation spectrum in biparental RILs (Figure 1).




Figure 1: Overview of inter-haplotype distance method for discovering mutator alleles. a) A population of four haplotypes has been genotyped at three informative markers; each haplotype also harbors private de novo germline mutations. At each informative marker, we compute an aggregate de novo germline mutation spectrum in the haplotypes that carry either parental allele, and calculate the cosine distance between the two aggregate spectra. b) We repeat the process outlined in a) for every informative marker along the genome. c) To assess the significance of any cosine distance peaks in b), we perform a permutation test by shuffling the labels associated with each haplotype's mutation data and running a genome-wide distance scan. In each of N permutations, we record the maximum distance encountered at any locus in the distance scan. Finally, we calculate the 1-p percentile of that maximum distance distribution to obtain a genome-wide cosine distance threshold at the specified value of p.

We assume that a collection of haplotypes has been genotyped at informative markers, and that *de novo* germline mutations have been identified on each haplotype.

At each informative marker, we divide haplotypes into two groups based on the parental allele that they inherited. We then compute a k-mer mutation spectrum using the aggregate mutation counts in each haplotype group. The k-mer mutation spectrum contains the frequency of every possible k-mer mutation type in a collection of mutations, and can be represented as a vector of size $6\times 4^{k-1}$ after collapsing by strand complement. For example, the 1-mer mutation spectrum is 6-element vector that contains the frequencies of C>T, C>G, C>A, A>G, A>T, and A>C mutations.

At each marker, we then calculate the cosine distance between the aggregate mutation spectra of haplotypes with either parental allele. The cosine distance between two vectors ${\bf A}$ and ${\bf B}$ is defined

$$D_C = 1 - rac{\mathbf{A} \cdot \mathbf{B}}{||\mathbf{A}|| ||\mathbf{B}||}$$

where $||\mathbf{A}||$ and $||\mathbf{B}||$ are the L^2 (or Euclidean) norms of \mathbf{A} and \mathbf{B} , respectively. The cosine distance metric has a number of favorable properties for comparing mutation spectra. Since cosine distance does not take the magnitude of vectors into account, it can be used to compare two spectra with unequal total mutation counts (even if those total counts are relatively small). Additionally, by calculating the cosine distance between mutation spectra, we avoid the need to perform separate comparisons of mutation counts at each individual k-mer mutation type.

Similar to existing methods for quantitative trait locus mapping [13], we use permutation tests to establish genome-wide cosine distance thresholds. In each of N permutation trials, we randomly shuffle the per-haplotype mutation data such that haplotype labels no longer correspond to the correct mutation counts. Using the shuffled mutation data, we perform a genome-wide distance scan as described above, and record the maximum distance observed at any locus. After N permutations (usually 10,000), we compute the 1-p percentile of the maximum distance distribution, and use that percentile value as a genome-wide significance threshold (for example, at p=0.05).

The inter-haplotype distance method was implemented in Python, and relies heavily on the following Python libraries: numpy, pandas, matplotlib, scikit-learn, pandera, seaborn, and numba [14,15,16,17,18,19,20].

Additional documentation is available on GitHub [21], along with a reproducible Snakemake [22] workflow for running the method from start to finish using the BXDs (including downloading the mutation data, downloading genotypes, and running a genome-wide distance scan).

Simulations to assess the power of the inter-haplotype distance approach

We performed a series of simple simulations to estimate our power to detect alleles that affect the germline mutation spectrum in biparental RILs using the inter-haplotype distance method.

First, we simulate the k-mer mutation spectrum in a population of h haplotypes. We assume that exactly $\frac{h}{2}$ of the haplotypes are under the effects of a mutator allele that increases the mutation rate of a particular mutation type(s) by an effect size e. We simulate m mutations on each haplotype as follows:

We first define a vector of 1-mer mutation probabilities:

$$P = (0.4, 0.1, 0.075, 0.075, 0.075, 0.275)$$

These probabilities sum to 1 and correspond to the expected frequencies of C>T, C>A, C>G, A>T, A>C, and A>G *de novo* germline mutations in mice, respectively [8].

If we are simulating the 3-mer mutation spectrum, we modify the vector of mutation probabilities P to be length 96, and assign every 3-mer mutation type a value of $\frac{P_c}{16}$, where P_c is the probability of the "central" mutation type associated with the 3-mer mutation type. In other words, each of the 16

possible NCN>NTN 3-mer mutation types would be assigned a mutation probability of $\frac{P_c}{16}=\frac{0.4}{16}=0.025$.

To simulate the mutation spectrum on the *wild-type* haplotypes, we define a matrix C of size $(\frac{h}{2},n)$, where $n=6\times 4^{k-1}$ (i.e., the number of k-mer mutation types being simulated). First, we generate a vector of lambda values by scaling the mutation probabilities by the number of mutations we wish to simulate:

$$\lambda = Pm$$

Then, we populate the matrix C by taking a single Poisson draw from the vector of λ values for each mutation type on each haplotype. Thus, for every row i in the matrix (i.e., for every haplotype), we perform the following for mutation type j:

$$C_{i,j} = \operatorname{Pois}(\lambda_j)$$

To simulate the mutation spectrum on the $\frac{h}{2}$ mutator haplotypes, we define a new matrix C' of size $(\frac{h}{2},n)$ as defined above. We then multiply the lambda value of a particular mutation type (or multiple mutation types) by the mutator effect size e to obtain λ' . Then, for every row i in the matrix:

$$C'_{i,j} = \operatorname{Pois}(\lambda'_i)$$

When k=1, we only augment the effect size of one mutation type at a time, but when k=3, we augment a fraction (25%, 50%, or 100%) of the 3-mer mutation types associated with a single "base" mutation type.

After generating mutator and wild-type haplotypes, we compute the aggregate mutation spectrum in either group by summing the columns of C and C'. We then calculate the cosine distance between the two aggregate spectra, which we call the "focal" distance D_f . To determine whether D_f is greater than what we'd expect by chance, we perform a permutation test.

First, we concatenate the matrices of wild-type and mutator haplotype spectra:

$$A = \left[egin{array}{c} C \ C' \end{array}
ight]$$

Then, in each of N=1,000 trials, we randomly permute the rows of A. In every permutation, we consider the row indices from $\left[0,\frac{h}{2}\right)$ to correspond to the wild-type haplotypes, and the row indices from $\left[\frac{h}{2},h\right)$ to correspond to the mutator haplotypes. We then compute the cosine distance between the aggregate spectra of the wild-type and mutator haplotypes. If fewer than 5% of the N permutations produces a cosine distance greater than or equal to D_f , we say that the approach successfully identified the mutator allele. For every combination of simulation parameters (h,m,e,a) and so on) we perform 100 trials and record the number of trials in which we successfully identify the mutator allele.

Applying the inter-haplotype distance method to the BXDs

We downloaded previously-generated BXD *de novo* germline mutation data from the GitHub repository associated with our previous manuscript, which was also archived at Zenodo [2,23,24], and downloaded a CSV file of BXD genotypes at 7,320 informative markers from GeneNetwork [25,26]. We also downloaded relevant metadata about each BXD RIL from the manuscript describing the updated BXD resource [3].

As in our previous manuscript [2], we included mutation data from a subset of the 152 BXDs in our inter-haplotype distance scans. We removed any BXDs that had been inbred for fewer than 20 generations, as it takes approximately 20 generations of strict brother-sister mating for an RIL genome to become ~99.8% homozygous [27]. As a result, any potential mutator allele would almost certainly be either fixed or lost after 20 generations. If fixed, the allele would remain linked to any excess mutations it causes for the duration of subsequent inbreeding, and its effects would be detectable using our methods. We also removed the BXD68 RIL from our genome-wide scans, since we previously discovered a hyper-mutator phenotype in that strain; the C>A germline mutation rate in BXD68 is over 5 times the population mean, likely due to a private deleterious nonsynonymous mutation in *Mutyh* [2]. In total, we included 94 BXD RILs in our genome-wide scans.

We used Snakemake [28] to write a reproducible workflow for running the inter-haplotype distance method on the BXD dataset, which has been deposited in the GitHub repository associated with this manuscript [21].

Identifying candidate mutator alleles overlapping the chromosome 6 peak

We investigated the region implicated by our inter-haplotype distance approach on chromosome 6 by subsetting the joint-genotyped BXD VCF file (European Nucleotide Archive accession **PRJEB45429** [29]) using bcftools [30]. To predict the functional impacts of both single-nucleotide variants and indels on splicing, protein structure, etc., we annotated variants in the BXD VCF using the following snpEff [31] command:

```
java -Xmx16g -jar /path/to/snpeff/jarfile GRCm38.75 /path/to/bxd/vcf >
/path/to/uncompressed/output/vcf
```

and used cyvcf2 [32] to iterate over the annotated VCF file in order to identify nonsynonymous fixed differences between the parental C57BL/6J and DBA/2J strains.

Comparing mutation spectra between Mouse Genomes Project strains

We downloaded mutation data from a previously published analysis [11] (Supplementary File 1, Excel Table S3) that identified strain-private mutations in 29 strains that were originally whole-genome sequenced as part of the Sanger Mouse Genomes (MGP) project [33]. When comparing counts of each mutation type between MGP strains that harbored either *D* or *B* alleles at the chromosome 4 or chromosome 6 mutator loci, we adjusted mutation counts by the number of callable A, T, C, or G nucleotides in each strain as described previously [2].

Querying GeneNetwork for evidence of eQTLs at the mutator locus

We used the online GeneNetwork resource [26], which contains array- and RNA-seq-derived expression measurements in a wide variety of tissues from numerous datasets, to find *cis*-eQTLs for the DNA repair genes we implicated under the cosine distance peak on chromosome 6. On the

GeneNetwork homepage (genenetwork.org), we selected the "BXD Family" **Group** and used the **Type** dropdown menu to select each of the specific expression datasets described in Table 2. In the **Get Any** text box, we then entered the specified gene name and clicked **Search**. After selecting the appropriate data record on the next page, we used the **Mapping Tools** dropdown to run Haley-Knott regression [34] with the following parameters: WGS-based marker genotypes, 1,000 permutations for LOD threshold calculations, and controlling for BXD genotypes at the rsm10000007390 marker.

The exact names of the expression datasets we used for each tissue are shown in Table 1 below:

Table 1: Names of gene expression datasets used for each tissue type on GeneNetwork

Tissue name	Complete name of GeneNetwork expression data	GeneNetwork trait ID	
Kidney	Mouse kidney M430v2 Sex Balanced (Aug06) RMA	1448815_at	
Gastrointestinal	UTHSC Mouse BXD Gastrointestinal Affy MoGene 1.0 ST Gene Level (Apr14) RMA	10540639	
Hematopoetic stem cells	UMCG Stem Cells ILM6v1.1 (Apr09) transformed	ILM1940279	
Hematopoetic progenitor cells	UMCG Progenitor Cells ILM6v1.1 (Apr09) transformed	ILM1940279	
Spleen	UTHSC Affy MoGene 1.0 ST Spleen (Dec10) RMA	10540639	
Liver	UTHSC BXD Liver RNA-Seq Avg (Oct19) TPM Log2	ENSMUST00000032406	
Heart	NHLBI BXD All Ages Heart RNA- Seq (Nov20) TMP Log2 **	ENSMUSG00000030271	
Eye	UTHSC BXD All Ages Eye RNA- Seq (Nov20) TPM Log2 **	ENSMUSG00000030271	

Calculating the frequencies of candidate mutator alleles in wild mice

For each of the three candidate nonsynonymous mutations we identified under the cosine distance peak on chromosome 6 (Table ??), we queried a VCF file containing genome-wide variation in 67 wild-derived mice from four species of *Mus* [35]. We calculated the allele frequency of each nonsynonymous mutation in each of the four species or subspecies (*Mus musculus domesticus*, *Mus musculus musculus, Mus musculus castaneus*, and *Mus spretus*), including genotypes that met the following criteria:

- supported by at least 10 sequencing reads
- Phred-scaled genotype quality of at least 20

Results

Benchmarking the inter-haplotype distance method using simulations

We first tested the inter-haplotype cosine distance approach using simulated data (Materials and Methods). We find that the method's power is mostly limited by the initial mutation rate of the k-mer

mutation type affected by the mutator allele and the total number of $de\ novo$ germline mutations in the dataset (that is, the product of the number of haplotypes and the mean number of mutations per haplotype) (Figure 2). For example, given 50 haplotypes with an average of 500 $de\ novo$ germline mutations each, our method has nearly 90% power detect a mutator allele that increases the C>T $de\ novo$ mutation rate by 10%. However, the method only has about 10% power to detect a mutator of identical effect size that affects the C>G mutation rate, since C>G mutations are expected to make up a much smaller fraction of all $de\ novo$ germline mutations to begin with. These simulations also demonstrate that our method is well-powered to detect large-effect mutator alleles (e.g., those that increase the mutation rate of a specific k-mer by 50%), even with a relatively small number of mutations per haplotype.




Figure 2: Simulations to assess the power of the inter-haplotype distance method. We simulated de novo germline mutations on the specified number of haplotypes, such that 50% of haplotypes were affected by a mutator allele that increased the mutation rate of the specified k-mer by the specified effect size (an effect size of 1.5 indicates a 50% increase in the mutation rate). The colors of the lines indicate the number of simulated mutations on each haplotype (before augmenting the mutation rate with a mutator allele). Given a specific combination of parameters, the y-axis denotes the fraction of 100 simulations in which the simulated mutator allele could be detected at a p-value of 0.05. Shaded areas indicate the standard deviation of that fraction.

Re-identifying the mutator allele on chromosome 4 in the BXDs

We applied our inter-haplotype distance method to 93 BXD RILs (Materials and Methods) with a total of 62,993 *de novo* germline mutations [2]. Reassuringly, we observed a large peak in cosine distance

at a locus on chromosome 4 (Figure <u>3</u>A; maximum distance of 9.20e-3 at marker ID rs52263933; position 116.75 Mbp in GRCm38/mm10 coordinates).




Figure 3: Results of inter-haplotype distance scans in the BXD RILs. a) Cosine distances between aggregate de novo mutation spectra on BXD haplotypes (n = 93 haplotypes; 62,993 total mutations) with either D or B alleles at 7,320 informative markers. Distance threshold at p=0.05 was calculated by performing 10,000 permutations of the BXD haplotype mutation data, and is shown as a dotted grey line. One outlier RIL with an extremely high C>A mutation rate (BXD68) was removed prior to running the distance scan. **b)** Cosine distances between aggregate de novo mutation spectra on BXD haplotypes with D alleles at rs52263933 (n = 55 haplotypes; 40,913 total mutations), the marker with the highest cosine distance in a) and either D or B alleles at 7,320 informative markers. Distance threshold at p=0.05 was calculated by performing 10,000 permutations of the BXD haplotype mutation data, and is shown as a dotted grey line. **c)** Fractions of de novo germline mutations in BXDs with either D or B haplotypes at markers rs52263933 and rs31001331. Total counts of each mutation type were aggregated in groups of BXDs with either of the four possible

haplotype combinations, and fractions of each mutation type were calculated in each group separately. Chi-square tests of independence were used to compare counts of individual mutation types between the D-D and D-B groups of BXDs, as well as between the B-D and B-B groups. **d)** Fractions of *de novo* germline mutations in Sanger Mouse Genome Project (MGP) strains with either *D* or *B* haplotypes at markers rs52263933 and rs31001331. Total counts of each mutation type were aggregated in groups of MGP strains with either of the four possible haplotype combinations, and fractions of each mutation type were calculated in each group separately. Chi-square tests of independence were used to compare counts of individual mutation types between the D-D and D-B groups of BXDs, as well as between the B-D and B-B groups.

In a previous analysis, we used quantitative trait locus (QTL) mapping to identify a nearly identical locus on chromosome 4 that was significantly associated with the C>A germline mutation rate in the BXDs [2]. This locus overlaps 21 protein-coding genes that are annotated by the Gene Ontology as being involved in "DNA repair," but only one of these genes contains non-synonymous differences between the two parental strains: *Mutyh. Mutyh* encodes a protein involved in the base-excision repair of 8-oxoguanine (8-oxoG), a DNA lesion caused by oxidative damage, and prevents the accumulation of C>A mutations [4,36,37]. C>A germline mutation rates are nearly 50% higher in BXDs that inherited *D* haplotypes at marker ID rs52263933 than in those that inherited *B* haplotypes [2].

An additional germline mutator allele on chromosome 6

After confirming that the inter-haplotype distance method could recover the mutator locus overlapping Mutyh, we asked if our approach could identify additional mutator loci in the BXD. To account for the effects of the large-effect C>A germline mutator locus near Mutyh, we divided the BXD RILs into those with either D (n = 55) or B (n = 38) genotypes at rs52263933 (the marker at which we observed the highest inter-haplotype cosine distance on chromosome 4), and ran a genome-wide distance scan using each group separately (Figure 3B.

Using only the BXDs with *B* genotypes at the *Mutyh* mutator locus, we did not observe any genome-wide significant peaks. But using the BXDs with *D* genotypes at the same locus, we identified a cosine distance peak on chromosome 6 (Figure 3B; maximum distance of 3.03e-3 at marker rs31001331; position 114.05 Mbp in mm10 coordinates). We queried the region underneath this peak (+/- 5 Mbp) and discovered 87 protein-coding genes. Remarkably, only one was annotated with the Gene Ontology term "DNA repair" and contained nonsynonymous differences between C57BL/6J and DBA/2J: *Ogg1*. *Ogg1* encodes a key member of the base-excision repair response to oxidative DNA damage, a pathway that also includes *Mutyh* and a related gene, *Mth1*. *Ogg1* harbors a single fixed nonsynonymous differences between the C57BL/6J and DBA/2J parental strains: p.Thr95Ala, at position 113,328,510 on chromosome 6 in GRCm38/mm10 coordinates.

We also considered the possibility that expression quantitative trait loci (eQTLs), rather than nonsynonymous mutations, could contribute to the C>A mutator phenotype linked to the locus on chromosome 6. Using GeneNetwork [26], we mapped cis-eQTLs for *Ogg1* in a number of tissues, including hematopoetic stem cells, kidney, and spleen. BXD genotypes near the cosine distance peak on chromosome 6 were significantly associated with *Ogg1* expression in many tissues, and *D* genotypes were nearly always associated with decreased gene expression (Table 2). We also queried a previously published collection of eQTLs derived from Diversity Outbred (DO) mouse embryonic stem cell (mESCs) expression data [38], but did not find any significant eQTLs for *Ogg1*.

 Table 2: Presence or absence of cis-eQTLs for Ogg1 in various tissues identified using GeneNetwork.

Tissue name	# BXDs with expression data	Top significant marker	LRS at top significant marker	Significant LRS threshold	Additive effect of D allele on expression
Kidney	53	rsm100000041 88	56.41	17.80	-0.18

Tissue name	# BXDs with expression data	Top significant marker	LRS at top significant marker	Significant LRS threshold	Additive effect of D allele on expression
Gastrointestinal	46	rsm100000034 41	30.10	16.21	-0.081
Hematopoetic stem cells	22	-	-	16.43	-
Hematopoetic progenitor cells	23	-	-	18.27	-
Spleen	79	rsm100000034	17.72	17.49	-0.056
Liver	50	rsm100000041	52.27	18.81	-0.155
Heart	73	-	-	16.12	-
Eye	87	rsm100000041 94	22.66	17.20	0.087

Evidence of epistasis between germline mutator alleles

We next compared the mutation spectra of BXDs with either B or D genotypes at the mutator loci on chromosomes 4 and 6. We observed that C>A germline mutation fractions in BXDs with D alleles at both mutator loci were significantly higher than C>A fractions in BXDs with D alleles at either locus alone (Figure 3C). However, compared to BXDs with B alleles at the chromosome 6 mutator locus, those with D alleles did not exhibit significantly higher C>A mutation fractions, indicating that the effects of the chromosome 6 mutator locus depend on the presence of a D allele at the chromosome 4 locus (Figure 3C).

To explore the effects of the two mutator loci in other inbred laboratory mice, we also compared the germline mutation spectra of Sanger Mouse Genomes Project (MGP) strains. Dumont [11] previously identified private germline mutations in 29 inbred laboratory strains; these private variants likely represent recent de novo germline mutations. Only two of the MGP strains possess D genotypes at both the chromosome 4 and chromosome 6 mutator loci: DBA/1J and DBA/2J. Compared to strains with B alleles at both mutator loci, those with D alleles at both mutator loci exhibit significantly higher C>A germline mutation fractions (p=3.5e-8, Figure 3D). MGP strains with D alleles at both mutator loci appear to have higher C>A mutation fractions than those with D alleles at either locus alone (Figure 3D), but this difference is not significant (p=0.16). Therefore, given the smaller number of MGP strains with D D0 D1. Therefore is not significant of epistasis observed in the BXDs.

The candidate Ogg1 mutator allele is present in wild mice

To determine whether the candidate mutator allele on chromosome 6 was segregating in natural populations of mice, we queried previously published sequencing data generated from 67 wild-derived mice [35]. These data include three subspecies of *Mus musculus*, as well as the outgroup *Mus spretus*. We found that the *D* allele in *Ogg1* was segregating at approximately 25% frequency in *Mus musculus domesticus*, the species from which C57BL/6J and DBA/2J derive the majority of their genomes [39], and was fixed in *Mus musculus musculus*, *Mus musculus castaneus*, and the outgroup *Mus spretus*.

Discussion

Using germline mutation spectra to identify mutator alleles

Germline mutation spectra are a rich source of information about the demographic history of populations, as well as the activity of both exogenous and endogenous sources of mutation throughout time. For example, by analyzing the 3-mer mutation spectrum in a collection of human genomes, Harris and Pritchard [40] discovered a "pulse" of TCC>TTC mutation activity in European populations that likely occurred between 15,000 and 2,000 years ago, and perhaps began even earlier [41]).

Within somatic tissues, mutation spectra can also be used to uncover the mutational processes active in particular populations of cells [42]. New computational methods have been developed to extract "mutational signatures" from large databases of somatic mutations in cancer [43]. These signatures, which describe the relative frequency of each 3-mer mutation type, can often be precisely attributed to chemotherapeutic agents, exposures to environmental mutagens, or loss-of-function mutations in genes encoding DNA repair or replication proteins [42,44,45].

Although a germline mutator allele should increase the absolute count of mutations on a linked haplotype, our results demonstrate that its effects can be more easily detectable by examining mutation *spectra* instead. For example, *D* alleles at the mutator locus on chromosome 6 augment the C>A mutation rate by a factor of approximately 1.2 (Figure 3). Since C>A mutations comprise approximately 10% of all germline mutations to begin with, *D* alleles only increase the overall germline mutation rate by about 2%. Given the depth of information that can be encoded in the mutation spectrum, we expect that mutation spectra can be further exploited to discover genetic modifiers of the mutation rate in other study systems, as well.

Epistasis between germline mutator alleles

Our results also reveal evidence of epistasis between mammalian germline mutator alleles for the first time. BXDs with *D* alleles at the mutator locus on chromosome 6 only exhibit elevated C>A mutation rates if they also carry *D* alleles at the previously-identified [2] mutator locus on chromosome 4. And BXDs with *D* alleles at both loci have significantly higher C>A germline mutation rates than lines with *D* alleles at only one mutator locus alone (Figure 3C). This raises the exciting possibility that epistasis between mutator alleles has contributed to the evolution of germline mutation rates and spectra in mammalian genomes.

Importantly, we note that we observed epistasis between germline mutator alleles in an unnatural population; the BXDs were inbred by brother-sister mating in a highly controlled laboratory environment that attenuated the effects of natural selection on all but the most deleterious alleles [12]. However, we found the *D* allele in *Ogg1* to be at nearly 25% frequency in *Mus musculus domesticus*, the strain from which C57BL/6J and DBA/2J derive most of their genomes [39]. Since the *D* mutator haplotype on chromosome 6 does not appear to increase the C>A germline mutation rate on its own (even in a homozygous state), we hypothesize that similar alleles may be at intermediate or high frequency in other natural populations.

Causal variants underlying the mutator allele

Only one DNA repair gene overlapping the C>A mutator locus on chromosome 6 also contained nonsynonymous fixed differences between the C57BL/6J and DBA/2J founder strains: *Ogg1*, a proteincoding gene that participates in base-excision repair of the oxidative DNA lesion 8-oxoguanine (8-oxoG) [4]. Both missense mutations and loss-of-heterozygosity in *Ogg1* have been associated with initiation and progression of various types of human cancer [46,47,48]. Unrepaired 8-oxoG lesions can

also lead to C>A mutations, and copy-number losses of either *Ogg1* or *Mutyh* are linked to elevated rates of spontaneous C>A mutation in human neuroblastoma [49]. Given these various lines of evidence, we believe that *Ogg1* is the most likely candidate gene to explain the additional C>A mutator phenotype in the BXDs, but it remains unclear whether the p.Thr95Ala missense mutation is the causal allele. We hypothesized that *Mutyh* missense mutations on *D* haplotypes were responsible for the large-effect C>A mutator phenotype we previously observed in the BXDs [2]. However, using high-quality long-read assemblies of inbred laboratory strains, another group recently identified a ~5 kbp mobile element insertion (MEI) within the first intron of *Mutyh* [50] that is present on *D* haplotypes and absent from *B* haplotypes. The MEI is associated with significantly reduced expression of *Mutyh* in embryonic stem cells from laboratory strains, and may therefore underlie the previous C>A germline mutator phenotype in the BXDs. In light of this new evidence, we cannot discount the possibility that eQTLs associated with decreased expression of *Ogg1* (Table 2) are responsible for the C>A mutator phenotype we observed in this study.

Discovering mutator alleles in other systems

Numerous lines of evidence suggest that mutator alleles contribute to variation in mutation rates and spectra across the tree of life. In two natural isolates of *Saccharomyces cerevisiae*, nonsynonymous variation in *OGG1* causes a substantial increase in the C>A *de novo* mutation rate [51]. A recent analysis suggested that mutator alleles and/or environmental mutagens have shaped mutation rate evolution in human genomes [52], and more generally, the mutation spectrum has evolved rapidly during great ape evolution, potentially due to the effects of *trans*-acting mutation rate modifiers [53]. The heritability of paternal *de novo* mutation counts in the human germline has also been estimated to be between 10 and 20%, demonstrating a contribution of genetic factors to germline mutation rates [54]). However, mutator discovery remains challenging in mammalian genomes.

What conditions must be met in order to detect a germline mutator allele? Presumably, one must have access to many haplotypes, each with a reasonably large number of de novo germline mutations that remain linked to the mutator allele(s) that caused them. Recently, thousands of human pedigrees have been sequenced in an effort to precisely estimate the rate of human de novo germline mutation [6,7]. Selection on germline mutator alleles will likely prevent large-effect mutators from reaching high allele frequencies; however, if multiple mutators are active in a particular population, it becomes much more likely that a subset will be detectable by sequencing human trios [55]. Current estimates of power to detect germline mutators in human pedigrees generally assume that mutators affect all mutation types equally, and that methods for mutator discovery will rely on identifying haplotypes with excess total mutation counts [55]. However, our results in the BXD suggest that germline mutators often exert their effects on a small number of k-mer mutation types, and may be far more amenable to detection by analyzing mutation spectra instead.

References

1. Mechanisms of DNA damage, repair, and mutagenesis.

Nimrat Chatterjee, Graham C Walker

Environmental and molecular mutagenesis (2017-05-09)

https://www.ncbi.nlm.nih.gov/pubmed/28485537

DOI: 10.1002/em.22087 · PMID: 28485537 · PMCID: PMC5474181

2. A natural mutator allele shapes mutation spectrum variation in mice.

Thomas A Sasani, David G Ashbrook, Annabel C Beichman, Lu Lu, Abraham A Palmer, Robert W Williams, Jonathan K Pritchard, Kelley Harris

Nature (2022-05-11) https://www.ncbi.nlm.nih.gov/pubmed/35545679

DOI: <u>10.1038/s41586-022-04701-5</u> · PMID: <u>35545679</u> · PMCID: <u>PMC9272728</u>

3. A platform for experimental precision medicine: The extended BXD mouse family.

David G Ashbrook, Danny Arends, Pjotr Prins, Megan K Mulligan, Suheeta Roy, Evan G Williams, Cathleen M Lutz, Alicia Valenzuela, Casey J Bohl, Jesse F Ingels, ... Robert W Williams *Cell systems* (2021-01-19) https://www.ncbi.nlm.nih.gov/pubmed/33472028

DOI: 10.1016/j.cels.2020.12.002 · PMID: 33472028 · PMCID: PMC7979527

4. Base-excision repair of oxidative DNA damage.

Sheila S David, Valerie L O'Shea, Sucharita Kundu

Nature (2007-06-21) https://www.ncbi.nlm.nih.gov/pubmed/17581577

DOI: <u>10.1038/nature05978</u> · PMID: <u>17581577</u> · PMCID: <u>PMC2896554</u>

5. Inferring evolutionary dynamics of mutation rates through the lens of mutation spectrum variation.

Jedidiah Carlson, William S DeWitt, Kelley Harris

Current opinion in genetics & development (2020-06-30)

https://www.ncbi.nlm.nih.gov/pubmed/32619789

DOI: 10.1016/j.gde.2020.05.024 · PMID: 32619789 · PMCID: PMC7646088

6. Parental influence on human germline de novo mutations in 1,548 trios from Iceland.

Hákon Jónsson, Patrick Sulem, Birte Kehr, Snaedis Kristmundsdottir, Florian Zink, Eirikur Hjartarson, Marteinn T Hardarson, Kristjan E Hjorleifsson, Hannes P Eggertsson, Sigurjon Axel Gudjonsson, ... Kari Stefansson

Nature (2017-09-20) https://www.ncbi.nlm.nih.gov/pubmed/28959963

DOI: 10.1038/nature24018 · PMID: 28959963

7. Large, three-generation human families reveal post-zygotic mosaicism and variability in germline mutation accumulation.

Thomas A Sasani, Brent S Pedersen, Ziyue Gao, Lisa Baird, Molly Przeworski, Lynn B Jorde, Aaron R Quinlan

eLife (2019-09-24) https://www.ncbi.nlm.nih.gov/pubmed/31549960

DOI: 10.7554/elife.46922 · PMID: 31549960 · PMCID: PMC6759356

8. Similarities and differences in patterns of germline mutation between mice and humans.

Sarah J Lindsay, Raheleh Rahbari, Joanna Kaplanis, Thomas Keane, Matthew E Hurles *Nature communications* (2019-09-06) https://www.ncbi.nlm.nih.gov/pubmed/31492841
DOI: 10.1038/s41467-019-12023-w · PMID: 31492841 · PMCID: PMC6731245

9. Genetic drift, selection and the evolution of the mutation rate.

Michael Lynch, Matthew S Ackerman, Jean-Francois Gout, Hongan Long, Way Sung, WKelley Thomas, Patricia L Foster

Nature reviews. Genetics (2016-10-14) https://www.ncbi.nlm.nih.gov/pubmed/27739533
DOI: 10.1038/nrg.2016.104 · PMID: 27739533

10. Inference of Candidate Germline Mutator Loci in Humans from Genome-Wide Haplotype Data.

Cathal Seoighe, Aylwyn Scally

PLoS genetics (2017-01-17) https://www.ncbi.nlm.nih.gov/pubmed/28095480

DOI: <u>10.1371/journal.pgen.1006549</u> · PMID: <u>28095480</u> · PMCID: <u>PMC5283766</u>

11. Significant Strain Variation in the Mutation Spectra of Inbred Laboratory Mice.

Beth L Dumont

Molecular biology and evolution (2019-05-01) https://www.ncbi.nlm.nih.gov/pubmed/30753674

DOI: <u>10.1093/molbev/msz026</u> · PMID: <u>30753674</u> · PMCID: <u>PMC6501876</u>

12. Spontaneous Mutation Accumulation Studies in Evolutionary Genetics

Daniel L Halligan, Peter D Keightley

Annual Review of Ecology, Evolution, and Systematics (2009-12-01) https://doi.org/dvrjz8

DOI: 10.1146/annurev.ecolsys.39.110707.173437

13. R/qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations.

Karl W Broman, Daniel M Gatti, Petr Simecek, Nicholas A Furlotte, Pjotr Prins, Śaunak Sen, Brian S Yandell, Gary A Churchill

Genetics (2018-12-27) https://www.ncbi.nlm.nih.gov/pubmed/30591514

DOI: 10.1534/genetics.118.301595 · PMID: 30591514 · PMCID: PMC6366910

14. Array programming with NumPy

Charles R Harris, KJarrod Millman, Stéfan J van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian Berg, Nathaniel J Smith, ... Travis E Oliphant *Nature* (2020-09-16) https://doi.org/ghbzf2

DOI: 10.1038/s41586-020-2649-2 · PMID: 32939066 · PMCID: PMC7759461

15. pandas-dev/pandas: Pandas

The Pandas Development Team

Zenodo (2023-02-20) https://doi.org/ggt8bh

DOI: 10.5281/zenodo.3509134

16. Matplotlib: A 2D Graphics Environment

John D Hunter

Computing in Science & Engineering (2007) https://doi.org/drbjhg

DOI: 10.1109/mcse.2007.55

17. Scikit-learn: Machine Learning in Python

Fabian Pedregosa, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter Prettenhofer, Ron Weiss, Vincent Dubourg, ... Édouard Duchesnay

Journal of Machine Learning Research (2011) http://jmlr.org/papers/v12/pedregosa11a.html

18. pandera: Statistical Data Validation of Pandas Dataframes

Niels Bantilan

Proceedings of the Python in Science Conference (2020) https://doi.org/grr54q

DOI: 10.25080/majora-342d178e-010

19. **seaborn: statistical data visualization**

Michael Waskom

Journal of Open Source Software (2021-04-06) https://doi.org/gjqn3g

DOI: 10.21105/joss.03021

20. Numba

Siu Kwan Lam, Antoine Pitrou, Stanley Seibert

Proceedings of the Second Workshop on the LLVM Compiler Infrastructure in HPC (2015-11-15)

https://doi.org/gf3nks

DOI: <u>10.1145/2833157.2833162</u>

21. https://github.com/quinlan-lab/proj-mutator-mapping

22. Sustainable data analysis with Snakemake

Felix Mölder, Kim Philipp Jablonski, Brice Letcher, Michael B Hall, Christopher H Tomkins-Tinch, Vanessa Sochat, Jan Forster, Soohyun Lee, Sven O Twardziok, Alexander Kanitz, ... Johannes Köster

F1000Research (2021-01-18) https://doi.org/gjjkwv

DOI: 10.12688/f1000research.29032.1 · PMID: 34035898 · PMCID: PMC8114187

23. A natural mutator allele shapes mutation spectrum variation in mice

Tom Sasani

(2023-01-24) https://github.com/tomsasani/bxd mutator manuscript

24. tomsasani/bxd_mutator_manuscript: Final figure generation updates prior to publication

Tom Sasani

Zenodo (2022-02-01) https://doi.org/grrwv8

DOI: 10.5281/zenodo.5941048

25. **BXD Genotype / WebQTL** https://gn1.genenetwork.org/dbdoc/BXDGeno.html

26. GeneNetwork: A Toolbox for Systems Genetics.

Megan K Mulligan, Khyobeni Mozhui, Pjotr Prins, Robert W Williams

Methods in molecular biology (Clifton, N.J.) (2017)

https://www.ncbi.nlm.nih.gov/pubmed/27933521

DOI: <u>10.1007/978-1-4939-6427-7_4</u> · PMID: <u>27933521</u> · PMCID: <u>PMC7495243</u>

27. Genetics and Probability in Animal Breeding Experiments

https://link.springer.com/book/10.1007/978-1-349-04904-2

28. Sustainable data analysis with Snakemake.

Felix Mölder, Kim Philipp Jablonski, Brice Letcher, Michael B Hall, Christopher H Tomkins-Tinch, Vanessa Sochat, Jan Forster, Soohyun Lee, Sven O Twardziok, Alexander Kanitz, ... Johannes Köster

F1000Research (2021-01-18) https://www.ncbi.nlm.nih.gov/pubmed/34035898

DOI: 10.12688/f1000research.29032.2 · PMID: 34035898 · PMCID: PMC8114187

29. **ENA Browser** https://www.ebi.ac.uk/ena/browser/view/PRJEB45429

30. Twelve years of SAMtools and BCFtools.

Petr Danecek, James K Bonfield, Jennifer Liddle, John Marshall, Valeriu Ohan, Martin O Pollard, Andrew Whitwham, Thomas Keane, Shane A McCarthy, Robert M Davies, Heng Li GigaScience (2021-02-16) https://www.ncbi.nlm.nih.gov/pubmed/33590861

DOI: 10.1093/gigascience/giab008 · PMID: 33590861 · PMCID: PMC7931819

31. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3.

Pablo Cingolani, Adrian Platts, Le Lily Wang, Melissa Coon, Tung Nguyen, Luan Wang, Susan J Land, Xiangyi Lu, Douglas M Ruden *Fly* (2012) https://www.ncbi.nlm.nih.gov/pubmed/22728672
DOI: 10.4161/fly.19695 · PMID: 22728672 · PMCID: PMCID: PMC3679285

32. cyvcf2: fast, flexible variant analysis with Python.

Brent S Pedersen, Aaron R Quinlan

Bioinformatics (Oxford, England) (2017-06-15) https://www.ncbi.nlm.nih.gov/pubmed/28165109

DOI: 10.1093/bioinformatics/btx057 · PMID: 28165109 · PMCID: PMC5870853

33. Mouse genomic variation and its effect on phenotypes and gene regulation.

Thomas M Keane, Leo Goodstadt, Petr Danecek, Michael A White, Kim Wong, Binnaz Yalcin, Andreas Heger, Avigail Agam, Guy Slater, Martin Goodson, ... David J Adams *Nature* (2011-09-14) https://www.ncbi.nlm.nih.gov/pubmed/21921910

DOI: <u>10.1038/nature10413</u> · PMID: <u>21921910</u> · PMCID: <u>PMC3276836</u>

34. A simple regression method for mapping quantitative trait loci in line crosses using flanking markers.

CS Haley, SA Knott

Heredity (1992-10) https://www.ncbi.nlm.nih.gov/pubmed/16718932

DOI: 10.1038/hdy.1992.131 · PMID: 16718932

35. Genomic resources for wild populations of the house mouse, Mus musculus and its close relative Mus spretus.

36. A Specific Mutational Signature Associated with DNA 8-Oxoguanine Persistence in MUTYH-defective Colorectal Cancer.

Alessandra Viel, Alessandro Bruselles, Ettore Meccia, Mara Fornasarig, Michele Quaia, Vincenzo Canzonieri, Eleonora Policicchio, Emanuele Damiano Urso, Marco Agostini, Maurizio Genuardi, ... Margherita Bignami

EBioMedicine (2017-04-13) https://www.ncbi.nlm.nih.gov/pubmed/28551381
DOI: 10.1016/j.ebiom.2017.04.022 · PMID: 28551381 · PMCID: PMCID: PMC5478212

37. Mutational signature analysis identifies MUTYH deficiency in colorectal cancers and adrenocortical carcinomas.

Camilla Pilati, Jayendra Shinde, Ludmil B Alexandrov, Guillaume Assié, Thierry André, Zofia Hélias-Rodzewicz, Romain Ducoudray, Delphine Le Corre, Jessica Zucman-Rossi, Jean-François Emile, ... Pierre Laurent-Puig

The Journal of pathology (2017-03-29) https://www.ncbi.nlm.nih.gov/pubmed/28127763
DOI: 10.1002/path.4880 · PMID: 28127763

38. Mapping the Effects of Genetic Variation on Chromatin State and Gene Expression Reveals Loci That Control Ground State Pluripotency.

Daniel A Skelly, Anne Czechanski, Candice Byers, Selcan Aydin, Catrina Spruce, Chris Olivier, Kwangbom Choi, Daniel M Gatti, Narayanan Raghupathy, Gregory R Keele, ... Laura G Reinholdt *Cell stem cell* (2020-08-13) https://www.ncbi.nlm.nih.gov/pubmed/32795400
DOI: 10.1016/j.stem.2020.07.005 · PMID: 32795400 · PMCID: PMC7484384

39. On the subspecific origin of the laboratory mouse.

Hyuna Yang, Timothy A Bell, Gary A Churchill, Fernando Pardo-Manuel de Villena *Nature genetics* (2007-07-29) https://www.ncbi.nlm.nih.gov/pubmed/17660819
DOI: 10.1038/ng2087 · PMID: 17660819

40. Rapid evolution of the human mutation spectrum.

Kelley Harris, Jonathan K Pritchard

eLife (2017-04-25) https://www.ncbi.nlm.nih.gov/pubmed/28440220

DOI: 10.7554/elife.24284 · PMID: 28440220 · PMCID: PMC5435464

41. Nonparametric coalescent inference of mutation spectrum history and demography.

William S DeWitt, Kameron Decker Harris, Aaron P Ragsdale, Kelley Harris

Proceedings of the National Academy of Sciences of the United States of America (2021-05-25)

https://www.ncbi.nlm.nih.gov/pubmed/34016747

DOI: 10.1073/pnas.2013798118 · PMID: 34016747 · PMCID: PMC8166128

42. Signatures of mutational processes in human cancer.

Ludmil B Alexandrov, Serena Nik-Zainal, David C Wedge, Samuel AJR Aparicio, Sam Behjati, Andrew V Biankin, Graham R Bignell, Niccolò Bolli, Ake Borg, Anne-Lise Børresen-Dale, ... Michael R Stratton

Nature (2013-08-14) https://www.ncbi.nlm.nih.gov/pubmed/23945592

DOI: <u>10.1038/nature12477</u> · PMID: <u>23945592</u> · PMCID: <u>PMC3776390</u>

43. Uncovering novel mutational signatures by

SMAshiqul Islam, Marcos Díaz-Gay, Yang Wu, Mark Barnes, Raviteja Vangara, Erik N Bergstrom, Yudou He, Mike Vella, Jingwei Wang, Jon W Teague, ... Ludmil B Alexandrov

Cell genomics (2022-11-09) https://www.ncbi.nlm.nih.gov/pubmed/36388765

DOI: 10.1016/j.xgen.2022.100179 · PMID: 36388765 · PMCID: PMC9646490

44. The mutational footprints of cancer therapies.

Oriol Pich, Ferran Muiños, Martijn Paul Lolkema, Neeltje Steeghs, Abel Gonzalez-Perez, Nuria Lopez-Bigas

Nature genetics (2019-11-18) https://www.ncbi.nlm.nih.gov/pubmed/31740835

DOI: <u>10.1038/s41588-019-0525-5</u> · PMID: <u>31740835</u> · PMCID: <u>PMC6887544</u>

45. Mutational signatures associated with tobacco smoking in human cancer.

Ludmil B Alexandrov, Young Seok Ju, Kerstin Haase, Peter Van Loo, Iñigo Martincorena, Serena Nik-Zainal, Yasushi Totoki, Akihiro Fujimoto, Hidewaki Nakagawa, Tatsuhiro Shibata, ... Michael R Stratton

Science (New York, N.Y.) (2016-11-04) https://www.ncbi.nlm.nih.gov/pubmed/27811275

DOI: 10.1126/science.aag0299 · PMID: 27811275 · PMCID: PMC6141049

46. Novel mutations of OGG1 base excision repair pathway gene in laryngeal cancer patients.

Ishrat Mahjabeen, Nosheen Masood, Ruqia Mehmood Baig, Maimoona Sabir, Uzma Inayat, Faraz Arshad Malik, Mahmood Akhtar Kayani

Familial cancer (2012-12) https://www.ncbi.nlm.nih.gov/pubmed/22829015

DOI: <u>10.1007/s10689-012-9554-2</u> · PMID: <u>22829015</u>

47. Alterations of the DNA repair gene OGG1 in human clear cell carcinomas of the kidney.

M Audebert, S Chevillard, C Levalois, G Gyapay, A Vieillefond, J Klijanienko, P Vielh, AK El Naggar, S Oudard, S Boiteux, JP Radicella

Cancer research (2000-09-01) https://www.ncbi.nlm.nih.gov/pubmed/10987279

PMID: 10987279

48. Mutations in OGG1, a gene involved in the repair of oxidative DNA damage, are found in human lung and kidney tumours.

S Chevillard, JP Radicella, C Levalois, J Lebeau, MF Poupon, S Oudard, B Dutrillaux, S Boiteux *Oncogene* (1998-06-11) https://www.ncbi.nlm.nih.gov/pubmed/9662341

DOI: 10.1038/sj.onc.1202096 · PMID: 9662341

49. Defects in 8-oxo-guanine repair pathway cause high frequency of C > A substitutions in neuroblastoma

Marlinde L van den Boogaard, Rurika Oka, Anne Hakkert, Linda Schild, Marli E Ebus, Michael R van Gerven, Danny A Zwijnenburg, Piet Molenaar, Lieke L Hoyng, MEmmy M Dolman, ... Jan J Molenaar

Proceedings of the National Academy of Sciences (2021-09-03) https://doi.org/grtcs9
DOI: 10.1073/pnas.2007898118 · PMID: 34479993 · PMCID: PMCID: PMC8433536

50. Resolution of structural variation in diverse mouse genomes reveals chromatin remodeling due to transposable elements

Ardian Ferraj, Peter A Audano, Parithi Balachandran, Anne Czechanski, Jacob I Flores, Alexander A Radecki, Varun Mosur, David S Gordon, Isha A Walawalkar, Evan E Eichler, ... Christine R Beck *Cold Spring Harbor Laboratory* (2022-09-27) https://doi.org/grtctb

DOI: <u>10.1101/2022.09.26.509577</u>

51. A modified fluctuation assay reveals a natural mutator phenotype that drives mutation spectrum variation within

Pengyao Jiang, Anja R Ollodart, Vidha Sudhesh, Alan J Herr, Maitreya J Dunham, Kelley Harris *eLife* (2021-09-15) https://www.ncbi.nlm.nih.gov/pubmed/34523420
DOI: 10.7554/elife.68285 · PMID: 34523420 · PMCID: PMCID: PMC8497059

52. Limited role of generation time changes in driving the evolution of mutation spectrum in humans

Ziyue Gao, Yulin Zhang, Nathan Cramer, Molly Przeworski, Priya Moorjani *bioRxiv* (2023-01-13) https://doi.org/10.1101/2022.06.17.496622

53. Mutational Signatures of Replication Timing and Epigenetic Modification Persist through the Global Divergence of Mutation Spectra across the Great Ape Phylogeny.

Michael E Goldberg, Kelley Harris

Genome biology and evolution (2022-01-04) https://www.ncbi.nlm.nih.gov/pubmed/33983415
DOI: 10.1093/gbe/evab104 · PMID: 33983415 · PMCID: PMCID: PMC8743035

54. Heritability of de novo germline mutation reveals a contribution from paternal but not maternal genetic factors

Seongwon Hwang, Matthew DC Neville, Genomics England Research Consortium, Felix R Day, Aylwyn Scally

bioRxiv (2022-12-17) https://doi.org/grr526
DOI: https://doi.org/10.1101/2022.12.17.520885

55. The impact of genetic modifiers on variation in germline mutation rates within and among human populations.

William R Milligan, Guy Amster, Guy Sella

Genetics (2022-07-30) https://www.ncbi.nlm.nih.gov/pubmed/35666194
DOI: 10.1093/genetics/iyac087 · PMID: 35666194 · PMCID: PMC9339295