

Ch 7.5 Integrating Rational Function by Partial Fraction

Anton's
Calculus
10th Ed.

Decompositions

$$\frac{3x-1}{(x-3)(x+4)} = \frac{A}{x-3} + \frac{B}{x+4}$$

$$\frac{5}{x(x^2-4)} = \frac{A}{x} + \frac{B}{x+2} + \frac{C}{x-2}$$

$$\frac{2x-3}{x^3-x^2} = \frac{2x-3}{x^2(x-1)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x-1}$$

$$\frac{x^2}{(x+2)^3} = \frac{A}{x+2} + \frac{B}{(x+2)^2} + \frac{C}{(x+2)^3}$$

$$\frac{1-x^2}{x^3(x^2+2)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x^3} + \frac{Dx+E}{x^2+2}$$

$$\frac{3x}{(x-1)(x^2+6)} = \frac{A}{x-1} + \frac{Bx+D}{x^2+6}$$

$$\frac{4x^3-x}{(x^2+5)^2} = \frac{Ax+B}{x^2+5} + \frac{Dx+E}{(x^2+5)^2}$$

$$\frac{1-3x^4}{(x-2)(x^2+1)^2} = \frac{A}{x-2} + \frac{Bx+C}{(x^2+1)} + \frac{Dx+E}{(x^2+1)^2}$$

Improper ~~Integral~~ fraction

$$\int \frac{3x^4 + 3x^3 - 5x^2 + x - 1}{x^2 + x - 2} dx$$

$$x^2 \times \frac{3x^2}{x^2} = 3x^4$$

$$x^2 \times \frac{1}{x^2} = x^2$$

$$\begin{array}{r} 3x^2 \quad +1 \\ x^2+x-2 \overline{) 3x^4+3x^3-5x^2+x-1} \\ \underline{3x^4+3x^3-6x^2} \\ x^2+x-1 \\ \underline{x^2+x-2} \\ 1 \end{array}$$

$$x^2+x-2 = x^2-2x+x-2 = (x-2)(x+1)$$

$$\int \left[3x^2 + 1 + \frac{1}{x^2+x-2} \right] dx$$

$$= \frac{3x^3}{3} + x + \int \frac{1}{x^2+x-2} dx$$

$$\frac{1}{x^2+x-2} = \frac{1}{(x+1)(x-2)}$$

$$= x^3 + x + \int \left(\frac{A}{x+1} + \frac{B}{x-2} \right) dx$$

$$\frac{1}{(x+1)(x-2)} = \frac{A}{x+1} + \frac{B}{x-2}$$

$$1 = A(x-2) + B(x+1)$$

$$= x^3 + x + \frac{1}{3} \int \left(\frac{1}{x+1} - \frac{1}{x-2} \right) dx$$

$$1 = Ax - 2A + Bx + B$$

$$= x^3 + x + \frac{1}{3} \left[\ln|x+1| - \ln|x-2| \right] + C$$

x

Constant

$$0 = A + B$$

$$1 = -2A + B$$

$$= x^3 + x + \frac{1}{3} \ln \left| \frac{x-1}{x+2} \right| + C$$

$$A = \frac{1}{3}, B = -\frac{1}{3}$$

Integration by substitution

$$\begin{aligned}\int \frac{\sec^2(\sqrt{x})}{\sqrt{x}} dx & \quad \text{let} \\ & \sqrt{x} = z \\ & \frac{1}{2\sqrt{x}} dx = dz \\ & \frac{1}{\sqrt{x}} dx = 2dz \\ & = 2 \int \sec^2 z dz \\ & = 2 \tan z + C \\ & = 2 \tan \sqrt{x} + C\end{aligned}$$

$$\begin{aligned}\int x \sqrt{1-x} dx & \quad \text{let} \\ & 1-x = z \\ & -dx = dz \\ & dx = -dz \\ & x = 1-z \\ & = \int (1-z) \sqrt{z} (-dz) \\ & = -\int (\sqrt{z} - z\sqrt{z}) dz \\ & = -\int (z^{1/2} - z^{3/2}) dz \\ & = -\frac{z^{3/2}}{\frac{3}{2}} + \frac{z^{5/2}}{\frac{5}{2}} + C \\ & = -\frac{2}{3} z^{3/2} + \frac{2}{5} z^{5/2} + C \\ & = -\frac{2}{3} (1-x)^{3/2} + \frac{2}{5} (1-x)^{5/2} + C\end{aligned}$$

$$\begin{aligned}\int x^2 e^{-2x^3} dx & \quad 2x^3 = z \\ & 6x^2 dx = dz \\ & x^2 dx = \frac{1}{6} dz \\ & = \int \frac{1}{6} e^{-z} dz \\ & = \frac{1}{6} \left[\frac{e^{-z}}{-1} \right] + C \\ & = -\frac{1}{6} e^{-2x^3} + C\end{aligned}$$

Ch 7.2

Reduction Formula

$$\begin{aligned}\int \cos^5 x dx &= \frac{1}{5} \cos^4 x \sin x + \frac{4}{5} \int \cos^3 x dx \\&= \frac{1}{5} \cos^4 x \sin x + \frac{4}{5} \left[\frac{1}{3} \cos^2 x \sin x + \frac{2}{3} \int \cos x dx \right] \\&= \frac{1}{5} \cos^4 x \sin x + \frac{4}{15} \cos^2 x \sin x + \frac{8}{15} \sin x + C\end{aligned}$$

$$\begin{aligned}\int \sin^3 2x \cos^2 2x \cancel{\cos 2x} dx \\&= \int \sin^2 2x \cos^2 2x \sin 2x dx \\&= \int (1 - \cos^2 2x) \cos^2 2x \sin 2x dx \\&= \int (\cos^2 2x - \cos^4 2x) \sin 2x dx \quad \rightarrow \begin{array}{l} \text{let} \\ \cos 2x = u \\ -\sin 2x (2) dx = du \\ -2 \sin 2x dx = du \\ \sin 2x dx = -\frac{1}{2} du \end{array} \\&= - \int (u^2 - u^4) \frac{1}{2} du \\&= -\frac{1}{2} \int (u^2 - u^4) du \\&= -\frac{1}{2} \left[\frac{u^3}{3} - \frac{u^5}{5} \right] + C \\&= -\frac{1}{2} \left[\frac{\cos^3 2x}{3} - \frac{\cos^5 2x}{5} \right] + C\end{aligned}$$

$$\int \tan^5 x \sec^1 x dx$$

$$= \int \tan^5 x \sec^2 x \sec^2 x dx$$

$$= \int \tan^5 x (1 + \tan^2 x) \sec^2 x dx$$

$$u = \tan x$$

$$du = \sec^2 x dx$$

$$= \int u^5 (1 + u^2) du$$

$$= \int (u^5 + u^7) du$$

$$= \frac{u^6}{6} + \frac{u^8}{8} + C$$

$$= \frac{\tan^6 x}{6} + \frac{\tan^8 x}{8} + C$$

Ch 7.2 Integration by Parts

$$\int x \tan^2 x dx$$

$$= x \int \tan^2 x dx - \int \left(\frac{d}{dx} x \int \tan^2 x dx \right) dx$$

$$= x \int (\sec^2 x - 1) dx - \int (1 \int (\sec^2 x - 1) dx) dx$$

$$= x (\tan x - x) - \int (\tan x - x) dx$$

$$= x \tan x - x^2 - \left[\ln |\sec x| - \frac{x^2}{2} \right] + C$$

$$= x \tan x - x^2 - \ln |\sec x| + \frac{x^2}{2} + C$$

$$= x \tan x - \frac{x^2}{2} - \ln |\sec x| + C$$

$$\int \sec^2 x dx = \tan x$$

$$\int \tan x = -\ln |\cos x|$$

$$= \ln |\sec x|$$

$$\int uv = u \int v - \int (u' \int v)$$

Integration by Parts

$$\int \cos(\ln x) dx$$

$$\ln x = z$$

$$\frac{1}{x} dx = dz$$

$$= \int e^z \cos z dz$$

$$dx = x dz$$

$$\because \ln x = z \\ x = e^z$$

$$= \int e^z \cos z dz - \int (e^z \int \cos z) dz = e^z dz$$

$$= e^z \sin z - \int e^z \sin z dz \quad \int uv = u \int v - \int (u' \int v)$$

$$= e^z \sin z - \left[e^z \int \sin z dz - \int \frac{d}{dz} e^z \int \sin z dz \right]$$

$$= e^z \sin z - \left[e^z \cos z - \int e^z (-\cos z) dz \right]$$

$$\int e^z \cos z dz = e^z \sin z + e^z \cos z - \int e^z \cos z dz$$

$$2 \int e^z \cos z dz = e^z (\sin z + \cos z)$$

$$\int e^z \cos z dz = \frac{1}{2} e^z (\sin z + \cos z)$$

$$\int \cos(\ln x) dx = \frac{1}{2} e^{\ln x} [\sin(\ln x) + \cos(\ln x)]$$

$$= \frac{1}{2} x [\quad , \quad]$$

Gamma Function

See the
Exercise
Sheet in box

$$\int_0^{\infty} e^{-x} x^{n-1} dx = \Gamma(n)$$

$$\text{or } \int_0^{\infty} e^{-t} t^{x-1} dt = \Gamma(x)$$

$$\int_0^{\infty} \sqrt{x} e^{-x^2} dx$$

$$= \int_0^{\infty} z^{1/4} e^{-z} \frac{1}{2\sqrt{z}} dz$$

$$= \frac{1}{2} \int_0^{\infty} e^{-z} z^{1/4 - 1/2} dz$$

$$= \frac{1}{2} \int_0^{\infty} e^{-z} z^{-1/4} dz$$

$$= \frac{1}{2} \int_0^{\infty} e^{-z} z^{\frac{3}{4}-1} dz = \frac{1}{2} \Gamma\left(\frac{3}{4}\right)$$

$$\begin{aligned} x^2 &= z \\ 2x dx &= dz \\ dx &= \frac{1}{2x} dz \\ dx &= \frac{1}{2\sqrt{z}} dz \end{aligned}$$

limits
are unchanged

$$z^{-1/4} = z^{n-1}$$

$$-\frac{1}{4} = n-1$$

$$n = 1 - \frac{1}{4} = \frac{3}{4}$$

$$\therefore n-1 = \frac{3}{4} - 1$$

$$\int_0^1 \frac{1}{\sqrt{x \ln(\frac{1}{x})}} dx$$

$$= \int_{\infty}^0 \frac{-e^{-z} dz}{\sqrt{e^{-z} z}}$$

$$= - \int_{\infty}^0 \frac{e^{-z} dz}{e^{-z/2} z^{1/2}}$$

$$= - \int_{\infty}^0 e^{-z+z/2} z^{-1/2} dz$$

$$= \int_0^{\infty} e^{-z/2} z^{-1/2} dz$$

$$= \int_0^{\infty} e^{-y} (2y)^{-1/2} 2dy$$

$$= 2 \cdot 2^{-1/2} \int_0^{\infty} e^{-y} y^{-1/2} dy$$

$$= 2^{1/2} \int_0^{\infty} e^{-y} y^{1/2-1} dy$$

$$= \sqrt{2} \Gamma\left(\frac{1}{2}\right)$$

$$\ln\left(\frac{1}{x}\right) = z$$

$$\ln 1 - \ln x = z$$

$$0 - \ln x = z$$

$$\ln x = -z$$

$$x = e^{-z}$$

$$dx = -e^{-z} dz$$

$$x=0 \rightarrow z=\infty$$

$$x=1 \rightarrow z=0$$

$$\frac{z}{2} = y$$

$$z = 2y$$

$$dz = 2dy$$

Limits

$$z=0 \rightarrow y=0$$

$$z=\infty \rightarrow y=\infty$$

$$y^{-1/2} = y^{n-1}$$

$$n-1 = -\frac{1}{2}$$

$$n = -1 - \frac{1}{2} = -\frac{1}{2}$$

$$n-1 = -\frac{1}{2} - 1$$