

20BXM01

HIMADRI MANDAL

November 20, 2021

§1 Solution

Solution. This is equivalent to finding d such that $\exists P \in \mathbb{Z}[X]$ with

$$S(P) = \#_{\mathbb{Z}}(P-1) + \#_{\mathbb{Z}}(P+1) \geq d+1$$

where $\#_{\mathbb{Z}}(P)$ denotes the number of distinct integral roots of P .

Claim — $S(P) \leq 4$ which also means $d \leq 3$.

Proof.

$$P(x) + 1 = Q(x)(x - \alpha_1)^{a_1}(x - \alpha_2)^{a_2} \cdots (x - \alpha_k)^{a_k}$$

$$P(x) - 1 = Q(x)(x - \alpha_1)^{a_1}(x - \alpha_2)^{a_2} \cdots (x - \alpha_k)^{a_k} - 2$$

If $P(x_0) - 1 = 0$ with $x_0 \in \mathbb{Z}$,

$$Q(x_0)(x_0 - \alpha_1)^{a_1}(x_0 - \alpha_2)^{a_2} \cdots (x_0 - \alpha_k)^{a_k} = 2$$

So, either $Q(x_0) = \pm 2, k \leq 2$ or $Q(x_0) = \pm 1, k \leq 3$, translating we get,

Case 1:

$$\pm 2x^{a_1}(x + c)^{a_2}$$

$$S(P) \leq 3$$

Case 2:

$$\pm 1x^{a_1}(x + c_1)^{a_2}(x + c_2)^{a_3}$$

Sorting $x, x + c_1, x + c_2$ and translating accordingly such that $x > x + c_1 > x + c_2$, but then $x = 2$ or 1 which determines the root itself. Thus, $S(P) \leq 4$ \square

Construction:

- $d = 1 : x$
- $d = 2 : 2x^2 - 1$
- $d = 3 : -x(x-1)(x-3) - 1$

\square