

# DENSITIES AND STABILITY VIA FACTORIZATION HOMOLOGY

Q.P. HỒ

**ABSTRACT.** Using factorization homology, we develop a uniform and conceptual approach for treating homological stability, homological densities, and arithmetic densities in algebraic geometry. This categorifies and generalizes the coincidences appearing in the work [FWW16] of Farb-Wolfson-Wood, and in fact, provides a conceptual understanding of these coincidences. Our computation of the stable homological densities also yields rational homotopy types which answer a question posed by Vakil-Wood in [VW15]. Our approach hinges on the study of homological stability of cohomological Chevalley complexes, which is of independent interest.

## CONTENTS

1. Introduction	2
1.1. Motivation	2
1.2. The goal of this paper	4
1.3. Other links to the literature	4
1.4. Prerequisites and guides to the literature	5
1.5. An outline of our strategy and results	5
1.6. An outline of the paper	8
2. Preliminaries	8
2.1. Notation and conventions	9
2.2. Koszul duality	9
2.3. Unital vs non-unital	12
2.4. Prestacks	13
2.5. Sheaves on prestacks	13
2.6. The Ran space/prestack	16
3. $\mathrm{coBar}_{\mathcal{P}}$ as a sequential limit	17
3.1. Co-filtered and graded objects	17
3.2. Stabilizing and decaying sequences	18
3.3. A co-filtration on $\mathrm{coBar}_{\mathcal{P}}$	21
4. Some generalities on homological stability	23
4.1. Filtered and graded objects	23
4.2. Unital and augmented algebras	26
4.3. Graded-unital and graded-augmented	27
4.4. Change of gradings	28
4.5. Stabilization	29
4.6. Relation to homological stability	31
4.7. Homological stability of a limit	33
4.8. Stable homology for objects of the form $W \otimes \Lambda[\mathbb{Z}_{\geq 0}^m]$	34
5. Cohomological Chevalley complex	35

---

*Date:* February 19, 2018.

*2010 Mathematics Subject Classification.* Primary 81R99. Secondary 18G55.

*Key words and phrases.* Chiral algebras, chiral homology, factorization algebras, Koszul duality, Ran space, homological stability, homological densities.

5.1. Homological stability	35
5.2. Stable homology	38
5.3. Decategorification	39
6. Factorization homology	44
6.1. Graded commutative factorization algebras	44
6.2. Koszul duality	47
6.3. Factorization cohomology	48
6.4. A multiplicative trace formula	49
6.5. Homological stability	51
6.6. Variants	52
7. Cohomology of $Z_n^m(X)$ as factorization cohomology	55
7.1. Commutative factorization algebras	55
7.2. Densities	57
7.3. coLie-coalgebras	58
7.4. Homological stability	60
7.5. Stable homological densities	60
7.6. Decategorifications	62
7.7. $L_\infty$ -algebras, the case of Poincaré series, and stable homological densities	63
Acknowledgments	65
References	65

## 1. INTRODUCTION

1.1. **Motivation.** For any connected curve  $X$  over  $\mathbb{F}_q$ , consider the symmetric powers and the (unordered) configuration spaces of  $X$

$$\mathrm{Sym}^d(X) = X^d / \Sigma_d, \quad \mathrm{Conf}_d(X) = \overset{\circ}{X}^d / \Sigma_d \subset^{\mathrm{open}} X^d / \Sigma_d = \mathrm{Sym}^d X,$$

where in the latter, one requires all the points to be distinct and where  $\Sigma_d$  denotes the group of permutations on  $d$  letters. Moreover, let

$$(\mathrm{Sym}^{d_1} X \times \mathrm{Sym}^{d_2} X)_{\mathrm{disj}} \subset^{\mathrm{open}} \mathrm{Sym}^{d_1} X \times \mathrm{Sym}^{d_2} X$$

be an open subscheme such that the collections of points in the two factors are disjoint. Then, it is well-known that we have the following coincidence

$$\lim_{d \rightarrow \infty} \frac{(\mathrm{Conf}_d X)(\mathbb{F}_q)}{(\mathrm{Sym}^d X)(\mathbb{F}_q)} = \lim_{(d_1, d_2) \rightarrow \infty} \frac{((\mathrm{Sym}^{d_1} X \times \mathrm{Sym}^{d_2} X)_{\mathrm{disj}})(\mathbb{F}_q)}{(\mathrm{Sym}^{d_1} X \times \mathrm{Sym}^{d_2} X)(\mathbb{F}_q)} = \zeta_X(2)^{-1}.$$

1.1.1. More generally, for any variety  $X$ , a pair of positive integers  $m, n$ , and an  $m$ -tuple of numbers

$$\mathbf{d} = (d_1, d_2, \dots, d_m) \in \mathbb{Z}_{\geq 0}^m,$$

let

$$\mathrm{Sym}^{\mathbf{d}}(X) = \prod_k \mathrm{Sym}^{d_k}(X),$$

and  $Z_n^{\mathbf{d}}(X) \subset \mathrm{Sym}^{\mathbf{d}}(X)$  consisting of sets  $D$  of  $|\mathbf{d}|$  (not necessarily distinct) points in  $X$  such that:

- (i) precisely  $d_k$  of the points in  $D$  are labeled with the “color”  $k$ , and
- (ii) no point of  $X$  has multiplicity of at least  $n$  for every color.

1.1.2. For example, when  $m = 1$  and  $\mathbf{d} = (d)$ , we have

$$\begin{aligned} Z_\infty^{\mathbf{d}}(X) &= Z_\infty^d(X) = \mathrm{Sym}^d X, \\ Z_2^{\mathbf{d}}(X) &= Z_2^d(X) = \mathrm{Conf}_d X. \end{aligned}$$

Similarly, when  $m = 2$  and  $\mathbf{d} = (d_1, d_2)$ , we have

$$\begin{aligned} Z_\infty^{\mathbf{d}}(X) &= \mathrm{Sym}^{d_1} X \times \mathrm{Sym}^{d_2} X, \\ Z_1^{\mathbf{d}}(X) &= (\mathrm{Sym}^{d_1} X \times \mathrm{Sym}^{d_2} X)_{\mathrm{disj}}. \end{aligned}$$

1.1.3. We have the following classical result (see, for example, [FWW16, Thm. 2.3]).

**Theorem 1.1.4.** *We have the following equality*

$$\lim_{\mathbf{d} \rightarrow \infty} \frac{|Z_n^{\mathbf{d}}(X)(\mathbb{F}_q)|}{|\mathrm{Sym}^{\mathbf{d}}(X)(\mathbb{F}_q)|} = \zeta_X(mn \dim X)^{-1}$$

for any connected variety  $X$  and any positive integers  $m, n$ .

1.1.5. Observing that the limit appearing in Theorem 1.1.4 only depends on the product  $mn$  and  $X$  itself, the authors of [FWW16] initiated the studies of “homological densities” of these spaces in the case where  $X$  is a smooth manifold or a complex variety. More specifically, they asked if the following limit of the quotient between the Poincaré polynomials

$$\lim_{\mathbf{d} \rightarrow \infty} \frac{P_{Z_n^{\mathbf{d}}(X)}(t)}{P_{\mathrm{Sym}^{\mathbf{d}}(X)}(t)} \in \mathbb{Z}[[t]]$$

also has the same kind of coincidence as in the case of  $\zeta$ -functions.

In many cases, the answer is positive, and we have the following result.

**Theorem 1.1.6** ([FWW16, Thm. 1.2]). *The limit*

$$\lim_{\mathbf{d} \rightarrow \infty} \frac{P_{Z_n^{\mathbf{d}}(X)}(t)}{P_{\mathrm{Sym}^{\mathbf{d}}(X)}(t)} \in \mathbb{Z}[[t]]$$

*exists, and depends only on the product  $mn$ , the Betti numbers of  $X$ , and  $\dim X$ , when any of the following assumptions is satisfied:*

- (i)  $X$  is a connected, smooth affine variety over  $\mathbb{C}$  and  $mn > 2$ ; or
- (ii)  $X$  is a connected, open submanifold of  $\mathbb{C}^r$  and  $m, n \geq 1$ ; or
- (iii)  $X$  is a connected open, noncompact submanifold of a smooth, orientable, even-dimensional compact manifold and  $mn > \dim X$ .

1.1.7. The proof of this theorem uses Björner-Wachs theory of lexicographic shellability from algebraic combinatorics as the main computational input. The extra conditions listed above are there to ensure that certain spectral sequences degenerate already at page 2 so we can have a good handle on the Poincaré polynomials, since these are not motivic.

Using essentially the same computational input, [FWW16] went on to show that similar coincidences hold for quotients of Euler characteristics as well as quotients of Hodge-Deligne polynomials (i.e. Euler characteristic densities and Hodge-Deligne densities). Unlike the case of Poincaré polynomials, however, these objects are motivic, which obviates the need for the spectral sequences involved to degenerate. As a result, these coincidences hold true without any of the three assumptions listed above.

**1.2. The goal of this paper.** Since the proofs of these coincidences in [FWW16] proceed by computing the densities using combinatorial means, we still do not know conceptually why these phenomena should occur in the first place. As the main goal of this paper, using factorization homology and Koszul duality as organizing tools, we provide a uniform and conceptual framework for understanding homological stability as well as coincidences in arithmetic and homological densities. This categorifies and in fact generalizes these results in two directions:

- (i)  $X$  is only required to be irreducible (rather than smooth), and in place of cohomology, we work with Borel-Moore homology  $C^*(X, \omega_X[-2 \dim X](-\dim X)) = C^*(X, \vec{\omega}_X)$  (see Notation 7.1.6 for the definition of  $\vec{\omega}$ ), which specializes to usual cohomology when  $X$  is smooth.
- (ii) Enlarge the class of configurations allowed (more on this below).

To reach this final goal, along the way, we prove a general algebraic statement about homological stability and stable homology of the cohomology of coLie-coalgebras,<sup>1</sup> which are of independent interest, and which forms the technical heart of the paper. The geometric statements about stability and coincidences are then obtained by applying these general results to certain coLie-coalgebras produced using techniques from factorization homology.

We would like to remark that most of the results in this paper are of a general nature. Moreover, once the general formalism has been laid down, results about the cohomology of these generalized configuration spaces follow naturally, requiring almost no computation. Indeed, configuration spaces do not appear until the last two sections, §6 and §7, and all the computations, which are simple and elementary in nature, are done in §7.

### 1.3. Other links to the literature.

1.3.1. In [VW15], motivated by the Dold-Thom theorem and its apparent link to homological stability as well as special values of  $\zeta$ -functions, Vakil-Wood ask if the appearance of special values of  $\zeta$ -functions is controlled by a certain rational homotopy type. As a result of our work, we see that all the  $\zeta$ -values that appear as arithmetic densities come from corresponding commutative algebras obtained from taking the quotients (as commutative algebras) of the stable homology of  $Z_n^m(X)$  by that of  $Z_\infty^m(X)$  (see Corollary 7.6.3 and Remark 7.6.5). The resulting commutative algebras, which we call the stable homological densities, could thus be viewed as the rational homotopy types<sup>2</sup> responsible for the appearance of the  $\zeta$ -values  $\zeta_X(mn \dim(X))^{-1}$ , and hence, give an answer to a question asked by Vakil-Wood in [VW15]. In fact, these homotopy types have a simple description which we compute explicitly, in Proposition 7.7.8.

1.3.2. Another interesting feature of our work is its resemblance to the topological factorization homology picture, making the link between the two worlds more precise.

First of all, the construction of the limiting rational homotopy type  $\bar{\mathcal{A}}_{m,n}(X)$  from  $\mathcal{A}_{m,n}(X)$  closely related to the construction of the group completion of an  $E_n$ -algebra via the telescoping construction in [MS76]. This is of special interest since group completion is known to be intimately linked to homological stability (see [May72, Seg73, KM18, KM16]). Note, however, that our procedure is applied to an  $E_\infty$ -algebra (i.e. commutative), rather than  $E_n$ .

Moreover, our result on homological stability shares the same local-to-global principle as the one in the work of Kupers-Miller in [KM18]. More precisely, the finiteness condition on the coLie-coalgebras resembles that which appears in loc. cit (see Remark 6.5.7). Our result, therefore, effectively provides an algebro-geometric analog of their topological result.

<sup>1</sup>Due to technical convenience, we will work most of the time with coLie-coalgebras rather than Lie-algebras. However, since all the concrete coLie-coalgebras that appear in the paper are finite dimensional, the reader should feel free to instead think about their duals, which are Lie-algebras. Note also that since everything is derived in our convention, what we call Lie-algebras are classically called dg-Lie algebras.

<sup>2</sup>Since everything is derived by our convention, by commutative algebra, we mean cdga.

Finally, note also that the commutative algebras appearing in our work is the  $E_n$ -Koszul dual to those in [KM18, Knu14, Ho17]. The fact that these happen to be commutative allows us, via another type of Koszul duality between coLie-coalgebras and commutative algebras, to reduce many computations to the coLie-coalgebra side, which is finite dimensional in nature, greatly simplifying the picture appearing there.

**1.4. Prerequisites and guides to the literature.** For the reader's convenience, we include a quick review of the necessary background as well as pointers to the existing literature in §2. The readers who are unfamiliar with the language and notation used in the introduction are encouraged to take a quick look there before returning to the current section.

**1.5. An outline of our strategy and results.** We will now give a more precise outline of our results. Technically, our work can be divided into two main parts, factorization cohomology and Koszul duality. The first is used to link the geometric world to the world of coLie-coalgebras and cohomological Chevalley complexes. The second serves as the technical tools to process the output produced by the first. The two aspects are technically quite different. The readers can thus read either one first, and refer to the other whenever necessary.

Throughout, we fix an irreducible scheme  $X$  of dimension  $d$  over an algebraically closed field. When we talk about Frobenius actions, we assume that  $X$  is over a field of characteristic  $p$ , and moreover, it is a pullback of a scheme defined over  $\mathbb{F}_q$ . Moreover, when we talk about point counts, we refer to the point counts of the original object over  $\mathbb{F}_q$ .

**1.5.1. Koszul duality and homological stability of cohomological Chevalley complexes.** We start with the Koszul duality aspect of the approach. Let  $\mathfrak{a}$  be a  $\mathbb{Z}_{\geq 0, +}^m$ -graded coLie-coalgebra. We want to ask when  $\mathrm{coChev}^{\mathrm{un}} \mathfrak{a}$ , the unital cohomological Chevalley complex, satisfies homological stability. But of course, the question does not make sense, since to talk about homological stability, we need to first have maps between different graded-degrees so that we can compare them, whereas a priori,  $\mathrm{coChev}^{\mathrm{un}} \mathfrak{a}$  is only a  $\mathbb{Z}_{\geq 0}^m$ -graded object.

Suppose that  $\mathrm{coChev}^{\mathrm{un}} \mathfrak{a}$  admits the structure of an algebra over the graded polynomial algebra  $\Lambda[\mathbb{Z}_{\geq 0}^m]$  (where the grading is the obvious one), then using the algebra action, we obtain maps between the different gradings, which put us in the setting where questions about homological stability make sense. Using Koszul duality, we can translate this condition to a condition on the coLie-coalgebra  $\mathfrak{a}$ . This allows us to single out a class of coLie-coalgebras, which we call strongly unital, where we can establish homological stability. The main result we prove in this setting is the following

**Theorem 1.5.2** (Theorem 5.1.12). *Let  $\mathfrak{a} \in \mathrm{coLie}^{\mathrm{un}}(\mathrm{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  be a strongly unital coLie-coalgebra. Fix  $c_0 \in \mathbb{N}$ ,  $1 \leq k \leq m$  and suppose that there exist  $s, s_1, \dots, s_{k-1}, s_{k+1}, \dots, s_m \in \mathbb{N}$  such that for any non-zero  $v \in H^*(\mathrm{Quot}_{\mathrm{un}}(\mathfrak{a}))^3$  of cohomological degree  $c - 1 \leq c_0$  and graded degree  $\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m$ , we have*

$$\mathbf{d}_k \leq sc + \sum_{i \neq k} s_i \mathbf{d}_i.$$

*Then, for all  $c \leq c_0$  and  $\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m$  such that*

$$\mathbf{d}_k \geq sc + \sum_{i \neq k} s_i \mathbf{d}_i,$$

*the following map is an equivalence*

$$\tau_{\leq c}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow \tau_{\leq c}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d} + \mathbf{1}_k},$$

*and the following map is injective*

$$H^{c+1}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow H^{c+1}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d} + \mathbf{1}_k}.$$

In particular, when  $\mathfrak{a}$  is strongly unital and finite dimensional, we have homological stability (Corollary 5.1.15 and Remark 5.1.16).

<sup>3</sup> $\mathrm{Quot}_{\mathrm{un}}(\mathfrak{a})$  denote the coLie-coalgebra obtained from removing the “unit” of  $\mathfrak{a}$ . See §5.1.3 for the precise definition.

1.5.3. When  $\mathfrak{a}$  is abelian,  $\mathrm{coChev}^{\mathrm{un}} \mathfrak{a} = \mathrm{Sym}(\mathfrak{a}[-1])$ , and the result is immediate. To reduce to this case, in Corollary §3.3.6, we establish a technical fact about the functor  $\mathrm{coChev}$  which says that it can be written as a sequential limit, whose successive fibers form  $\mathrm{Sym}(\mathfrak{a}[-1])$ . In fact, we prove a similar result for  $\mathrm{coBar}_{\mathcal{P}}$  for any co-operad  $\mathcal{P}$  in Proposition 3.3.5, which specializes to the case of  $\mathrm{coChev}$  when  $\mathcal{P} = \mathrm{coLie}$ . In §4.7, we establish the relation between homological stability of a sequential limit and that of the associated fiber, which allows us to make the reduction.

1.5.4. The use of Koszul duality is crucial not only in the understanding of homological stability, but also in extracting information out of an algebra (or more relevant to us, quotients of algebras) via decategorification. Roughly speaking, the functor  $\mathrm{coChev}$  exhibits a multiplicative nature, which, in many cases, manifests itself via products/quotients of the decategorification. This is why it is useful for our purposes.

In §5.3, we establish several results in this direction which will be used later to extract information out of factorization cohomology. We learned this picture from [GL14], of which the results in §5.3 are routine adaptations.

1.5.5. *Factorization cohomology and cohomology of configuration spaces.* We will now discuss the factorization cohomology aspect of the approach.

1.5.6. Let us start with the definition of the generalized configuration spaces associated to  $X$  that we will study. Let  $m$  be a positive integer and  $\mathbf{n}$  an  $m$ -tuple of positive integers such that

$$|\mathbf{n}| = \sum_{k=1}^m \mathbf{n}_k \geq 2.$$

Similarly to §1.1.1, for each  $\mathbf{d} \in \mathbb{Z}_{\geq 0}^m$ , we let

$$Z_{\mathbf{n}}^m(X) \subset \mathrm{Sym}^{\mathbf{d}}(X)$$

consisting of sets  $D$  of  $|\mathbf{d}|$  (not necessarily distinct) points of  $X$  such that

- (i) precisely  $\mathbf{d}_k$  of the points in  $D$  are labeled with the “color”  $k$ , and
- (ii) no point of  $X$  has multiplicities of at least  $\mathbf{n}_k$  for every  $k$ .

We will use the following notation to gather the whole family together

$$Z_{\mathbf{n}}^m = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} Z_{\mathbf{n}}^{\mathbf{d}}.$$

We will also use the following notation

$$Z_{\infty}^m(X) = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} Z_{\infty}^{\mathbf{d}}(X) = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} \mathrm{Sym}^{\mathbf{d}}(X).$$

1.5.7. The starting point is the  $\mathbb{Z}_{\geq 0}^m$ -graded commutative algebra<sup>4</sup>

$$A_{m,\mathbf{n}} = \Lambda[x_1, \dots, x_m] / (\cap_{k=1}^m x_k^{\mathbf{n}_k}),$$

and when  $\mathbf{n} = \infty$ ,

$$A_{m,\infty} = \Lambda[x_1, \dots, x_m].$$

We prove, in Proposition 7.1.8, that the factorization cohomology of  $X$  with coefficients in  $A_{m,\mathbf{n}}$ , which we will denote by  $\mathcal{A}_{m,\mathbf{n}}(X)$ , computes the cohomology of  $Z_{\mathbf{n}}^m(X)$  (including the case of  $\mathbf{n} = \infty$ ).<sup>5</sup> Since the algebras  $A_{m,\mathbf{n}}$  are very simple, their Koszul duals  $a_{m,\mathbf{n}}$  can be easily computed. As a result, we have an expression of  $\mathcal{A}_{m,\mathbf{n}}(X)$  as the  $\mathrm{coChev}^{\mathrm{un}}$  of a simple  $\mathrm{coLie}$ -coalgebra  $\mathfrak{a}_{m,\mathbf{n}}(X)$ . Homological stability of  $Z_{\mathbf{n}}^m(X)$ , or equivalently, of  $\mathcal{A}_{m,\mathbf{n}}(X)$ , is now a direct consequence of Theorem 5.1.12 and we obtain

<sup>4</sup>To keep the introduction simple, we exclude cohomological shifts and the gradings from the notation.

<sup>5</sup>As mentioned earlier, by cohomology of  $Z_{\mathbf{n}}^{\mathbf{d}}(X)$  for example, we really mean  $C^*(Z_{\mathbf{n}}^{\mathbf{d}}, \bar{\omega}_{Z_{\mathbf{n}}^{\mathbf{d}}})$ . See Notation 7.1.6.

**Theorem 1.5.8** (Theorem 7.4.1). *Let  $X$  be an irreducible scheme of dimension  $d \geq 1$ . For each  $1 \leq k \leq m$  and  $c \geq 0$ , there exists a natural map (see Notation 7.1.6)*

$$H^c(Z_{\mathbf{n}}^{\mathbf{d}}(X), \vec{\omega}_{Z_{\mathbf{n}}^{\mathbf{d}}(X)}) \rightarrow H^c(Z_{\mathbf{n}}^{\mathbf{d}+1_k}(X), \vec{\omega}_{Z_{\mathbf{n}}^{\mathbf{d}+1_k}(X)})$$

that is

- (i) an equivalence when  $\mathbf{d}_k \geq 2c$ , and injective when  $\mathbf{d}_k = 2c - 1$ , when  $d = 1, m = 1, n = 2$  (the case of configuration spaces of a curve),
- (ii) an equivalence when  $\mathbf{d}_k \geq c$ , and injective when  $\mathbf{d}_k = c - 1$  otherwise.

More qualitative, and more generally, we prove the following result about homological stability of factorization cohomology

**Theorem 1.5.9** (Theorem 6.5.6). *Let  $X$  be an irreducible scheme of dimension  $d$ . Let  $\mathcal{A} \in \text{ComAlg}(\text{Vect}_{\geq 0, +}^{\mathbb{Z}^m})$  such that  $\mathfrak{a} = \text{coPrim}[1](\mathcal{A})$  is a finite dimensional  $d$ -shifted strongly unital coLie-coalgebra (see Definition 6.5.3). Then,*

$$\pi_{?_*}^{\text{un}} \pi^! \mathcal{A} \simeq \text{coChev}^{\text{un}}(C^*(X, \omega_X) \otimes \mathfrak{a})$$

satisfies homological stability.

This Theorem is closely related to the one in [KM18]. Indeed, the finiteness condition on  $\mathfrak{a}$  (or more generally, the bound on the cohomological amplitudes stated in Theorem 5.1.12) could be viewed as the analog of the bounded generation property of loc. cit. See also Remark 6.5.7.

In what follows, we will use  $\bar{\mathcal{A}}_{m, \mathbf{n}}(X)$  to denote the stable homology of  $\mathcal{A}_{m, \mathbf{n}}(X)$  (or equivalently, of  $Z_{\mathbf{n}}^m(X)$ ), including the case where  $\mathbf{n} = \infty$ . Via a general functorial construction (mimicking the telescoping construction of [MS76]) which works even when there is no homological stability, we show that  $\bar{\mathcal{A}}_{m, \mathbf{n}}(X)$  naturally inherits a structure of a commutative algebra (see §4.5.12 and Corollary 4.5.7).

1.5.10. *Categorified densities.* We have a natural map of commutative algebras

$$\mathcal{A}_{m, \infty} \simeq \Lambda[x_1, \dots, x_m] \rightarrow \Lambda[x_1, \dots, x_m] / (\cap_{k=1}^m x_k^{\mathbf{n}_k}) \simeq \mathcal{A}_{m, \mathbf{n}},$$

which, via taking factorization cohomology, induces a map of commutative algebras

$$\mathcal{A}_{m, \infty}(X) \rightarrow \mathcal{A}_{m, \mathbf{n}}(X),$$

which, in turn, induces a map of commutative algebras

$$\bar{\mathcal{A}}_{m, \infty}(X) \rightarrow \bar{\mathcal{A}}_{m, \mathbf{n}}(X)$$

by taking stable homology.

1.5.11. Now, there are naturally two kinds of “densities” we can examine:

$$\Lambda \otimes_{\mathcal{A}_{m, \infty}(X)} \mathcal{A}_{m, \mathbf{n}}(X) \quad \text{and} \quad \Lambda \otimes_{\bar{\mathcal{A}}_{m, \infty}(X)} \bar{\mathcal{A}}_{m, \mathbf{n}}(X).$$

The first captures the density of the whole family  $Z_{\mathbf{n}}^m(X)$  inside the family  $Z_{\infty}^m(X)$ , whereas the latter captures the stable density, i.e. density between stable homologies. The goal is to see if these densities depend only on  $|\mathbf{n}| = \sum_{k=1}^m \mathbf{n}_k$ .

Using the fact that factorization cohomology preserves relative tensors of algebras, it is now a direct consequence of the following equivalences

$$\Lambda \otimes_{\Lambda[x_1, \dots, x_m]} (\Lambda[x_1, \dots, x_m] / (\cap_{k=1}^m x_k^{\mathbf{n}_k})) \simeq \Lambda \oplus \Lambda[1] \simeq \Lambda \otimes_{\Lambda[x]} (\Lambda[x] / (x^{|\mathbf{n}|})).$$

In other words, the a priori complicated densities between the cohomology of  $Z_{\mathbf{n}}^m(X)$  (either stable or the whole family) are controlled by extremely simple quotients involving polynomial algebras, which explains the source of the coincidences. We prove the following result.



**Theorem 1.5.12** (Theorems 7.2.4, 7.5.1, and Proposition 7.7.8). *We have the following equivalence of algebras<sup>6</sup>*

$$\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X) \simeq \text{oblv}_{\text{gr}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)).$$

Moreover,  $\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)$  and  $\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)$  depend only on  $|\mathbf{n}|$ .

We in fact produce canonical maps between these quotients for various  $m, |\mathbf{n}|$ . Moreover, we explicitly compute these two quotients, in Corollary 7.3.8 and Proposition 7.7.8 respectively. We note that the equivalence in the Theorem above is a special feature of the current situation, which is due to the fact that a certain coLie-coalgebra happens to be abelian due to degree reasons.

1.5.13. *Decategorifications.* Applying various decategorification procedures (for example, Euler characteristics, Poincaré polynomials, or when  $X$  is a pullback of a scheme over  $\mathbb{F}_q$ , virtual Poincaré polynomials, and also traces of the Frobenius) to Theorem 1.5.12, we obtain generalizations of the results proved in [FWW16]. This is done in §7.6 and §7.7. In this part of the introduction, let us highlight the two cases of decategorification that are of special interest: Frobenius traces and Poincaré polynomials.

1.5.14. We show, in Corollary 7.6.3, that computing the Frobenius trace on  $\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)$  recovers the  $\zeta$ -value  $\zeta(\dim X|\mathbf{n}|)^{-1}$  and moreover, it is linked to the limit of the quotient of the point counts on  $Z_{\mathbf{n}}^m(X)$  and  $Z_{\infty}^m(X)$  (Remark 7.6.5). As mentioned above, the commutative algebra  $\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)$  thus provides an answer to the question posed in [VW15].

1.5.15. One feature that makes the case of Poincaré polynomials stands out is that it does not behave nicely with respect to cofiber sequences. Because of this reason, quotients of algebras do not translate to quotients of Poincaré polynomials. In Proposition 5.3.12, we formulate a special condition where Poincaré polynomials still behave nicely with respect to relative tensors of commutative algebras. Translated to the geometric setting, it has to do with the coLie-coalgebras  $\mathfrak{a}_{m,\mathbf{n}}(X)$  (the Koszul dual of  $\mathcal{A}_{m,\mathbf{n}}(X)$ ) being abelian. A simple consideration of the graded-vs-cohomological degrees shows why we need the extra conditions in Theorem 1.1.6 of [FWW16], which is generalized in Proposition 7.7.5.

1.6. **An outline of the paper.** We will now give a brief outline of how the paper is organized.

In §2, besides fixing the notations, we give a brief review of the various mathematical tools used throughout the paper as well as pointers to the existing literature. In §3, we establish a technical result concerning the functor of taking cohomological Chevalley complex (or more generally, the functor  $\text{coBar}_{\mathcal{P}}$  for a co-operad  $\mathcal{P}$ ) in a pro-nilpotent category, which will allow us to reduce many arguments to the case where the coLie-coalgebra involved is trivial (a.k.a. abelian). In §4 we lay down the framework where homological stability naturally takes place. In §5 we prove homological stability for cohomological Chevalley complexes for a large class of coLie-coalgebras (along with the explicit stability bounds) and give an expression for the stable homology in special cases. Moreover, we also introduce various decategorification procedures to extract information out of a cohomological Chevalley complex. Geometry makes its first appearance in §6. After a quick review of factorization algebra and homology (in the graded setting), we prove homological stability of factorization cohomology and end the section with a functorial procedure on commutative factorization algebras which will be used later in §7 to link the cohomology of  $Z_{\mathbf{n}}^m(X)$  to factorization cohomology. In §7, we finally study the spaces  $Z_{\mathbf{n}}^m(X)$ . In particular, we prove agreements between categorified homological densities and show that via decategorifications, we obtain generalizations of previous known results.

## 2. PRELIMINARIES

In this section, we collect notation, conventions, as well as provide definitions, results, as well as references to them, that we will use throughout the paper.

<sup>6</sup>Here,  $\text{oblv}_{\text{gr}}$  denotes the functor of forgetting the gradings, i.e. taking direct sum of all the graded components.



## 2.1. Notation and conventions.

**2.1.1. Category theory.** We will use  $\mathrm{DGCat}$  to denote the  $(\infty, 1)$ -category of stable infinity categories,  $\mathrm{DGCat}_{\mathrm{pres}}$  to denote the full subcategory of  $\mathrm{DGCat}$  consisting of presentable categories, and  $\mathrm{DGCat}_{\mathrm{pres}, \mathrm{cont}}$  the (non-full) subcategory of  $\mathrm{DGCat}_{\mathrm{pres}}$  where we only allow continuous functors, i.e. those commuting with colimits.

We will also fix an algebraically closed field  $\Lambda$  of characteristic  $o$  (see also §2.5.1). Unless otherwise specified, all our stable infinity categories and functors between them are in  $\mathrm{DGCat}_{\mathrm{pres}, \mathrm{cont}}$ . In particular, for symmetric monoidal categories, the monoidal/tensor products commute with colimit in each variable. Moreover, we ask that these categories are linear over  $\Lambda$ .

The main reference for the subject are [Lur17a, Lur17b]. For a slightly different point of view, see also [GR17].

**2.1.2. Algebraic geometry.** Throughout this paper,  $k$  will be used to denote an algebraically closed ground field. We will denote by  $\mathrm{Sch}$  the  $\infty$ -category obtained from the ordinary category of separated schemes of finite type over  $k$ . All our schemes will be objects of  $\mathrm{Sch}$ . In most cases, we will use the calligraphic font to denote prestacks, for eg.  $\mathcal{X}, \mathcal{Y}$  etc., and the usual font to denote schemes, for eg.  $X, Y$  etc.

**2.1.3.  $t$ -structure.** Let  $\mathcal{C}$  be a stable infinity category, equipped with a  $t$ -structure. Then we have the following diagram of adjoint functors

$$\mathcal{C}^{\leq 0} \begin{array}{c} \xleftarrow{i_{\leq 0}} \\ \xrightarrow{\mathrm{tr}_{\leq 0}} \end{array} \mathcal{C} \begin{array}{c} \xleftarrow{\mathrm{tr}_{\geq 1}} \\ \xrightarrow{i_{\geq 1}} \end{array} \mathcal{C}^{\geq 1}$$

We use  $\tau_{\leq 0}$  and  $\tau_{\geq 1}$  to denote

$$\tau_{\leq 0} = i_{\leq 0} \circ \mathrm{tr}_{\leq 0} : \mathcal{C} \rightarrow \mathcal{C}$$

and

$$\tau_{\geq 1} = i_{\geq 1} \circ \mathrm{tr}_{\geq 1} : \mathcal{C} \rightarrow \mathcal{C}$$

respectively.

Shifts of these functors, for e.g.  $\tau_{\geq n}$  and  $\tau_{\leq n}$ , are defined in the obvious ways.

**2.2. Koszul duality.** The theory of Koszul duality, initially developed in [Qui69], illuminated the duality between co-commutative co-algebras and Lie algebras. It was vastly generalized to the operadic setting in [GK94]. An application of this theory to the factorizable setting was carried out in [FG11].

In this paper, we will make extensive use of Koszul duality between commutative algebras and  $\mathrm{coLie}$ -coalgebras. We will summarize the relevant features of this theory below. The interested readers could find a more detailed development in [FG11, GR17, LV12]. The presentation here follows closely those of [FG11, GR17].

**2.2.1. Symmetric sequences.** Let  $\mathrm{Vect}^{\Sigma}$  denote the category of symmetric sequences. Namely, it consists, as objects, collections

$$\mathcal{O} = \{\mathcal{O}(n), n \geq 1\},$$

where  $\mathcal{O}(n) \in \mathrm{Vect}$  equipped with an action of  $\Sigma_n$ , the permutation group of  $n$  letters.

This category is equipped with a (non-symmetric) monoidal structure, designed to make the functor

$$\mathrm{Vect}^{\Sigma} \rightarrow \mathrm{Fun}(\mathrm{Vect}, \mathrm{Vect})$$

given by

$$\mathcal{O} \star V = \bigoplus_{n \geq 1} (\mathcal{O}(n) \otimes V^{\otimes n})_{\Sigma_n}$$

monoidal.

2.2.2. *Operads and co-operads.* By an operad (resp. co-operad), we will mean an augmented associative algebra (resp. co-algebra) in  $\text{Vect}^\Sigma$ . We will use  $\text{Op}$  (resp.  $\text{coOp}$ ) to denote the categories of operads (resp. co-operads).

2.2.3. The Bar and coBar construction gives us a pair of adjoint functors

$$\text{Bar} : \text{Op} \rightleftarrows \text{coOp} : \text{coBar}.$$

For an operad  $\mathcal{O}$  (resp. co-operad  $\mathcal{P}$ ), we will also use  $\mathcal{O}^\vee$  (resp.  $\mathcal{P}^\vee$ ) to denote  $\text{Bar}(\mathcal{O})$  (resp.  $\text{coBar}(\mathcal{P})$ ).

In this paper, we will restrict ourselves to the class of operads/co-operads where the augmentation map is an equivalence in degree 1, i.e.  $\mathcal{O}(1) \simeq \Lambda$  and  $\mathcal{P}(1) \simeq \Lambda$  (see 2.5.1). Under this restriction, one can show that the unit map

$$\mathcal{O} \rightarrow \text{coBar} \circ \text{Bar}(\mathcal{O}) = (\mathcal{O}^\vee)^\vee$$

is an equivalence.

2.2.4. *Algebras and co-algebras.* Let  $\mathcal{C}$  be a stable presentable symmetric monoidal  $\infty$ -category compatibly tensored over  $\text{Vect}$ . Then, an operad  $\mathcal{O}$  (resp. co-operad  $\mathcal{P}$ ) naturally defines a monad (resp. comonad) on  $\mathcal{C}$ . Thus, we can talk about the category of algebras  $\mathcal{O}\text{-alg}(\mathcal{C})$  (resp. coalgebras  $\mathcal{P}\text{-coalg}(\mathcal{C})$ ) in  $\mathcal{C}$ .

One has the following pairs of adjoint functors

$$\text{Free}_{\mathcal{O}} : \mathcal{C} \rightleftarrows \mathcal{O}\text{-alg}(\mathcal{C}) : \text{oblv}_{\mathcal{O}} \quad \text{and} \quad \text{Bar}_{\mathcal{O}} : \mathcal{O}\text{-alg}(\mathcal{C}) \rightleftarrows \mathcal{C} : \text{triv}_{\mathcal{O}}$$

for an operad  $\mathcal{O}$ , and similarly, the following pairs of adjoint functors

$$\text{oblv}_{\mathcal{P}} : \mathcal{P}\text{-coalg}(\mathcal{C}) \rightleftarrows \mathcal{C} : \text{coFree}_{\mathcal{P}} \quad \text{and} \quad \text{cotriv}_{\mathcal{P}} : \mathcal{C} \rightleftarrows \mathcal{P}\text{-coalg}(\mathcal{C}) : \text{coBar}_{\mathcal{P}}$$

for a co-operad  $\mathcal{P}$ .

2.2.5. *Koszul duality.* The functors mentioned above in fact lift to the following pair of adjoint functors

$$(2.2.6) \quad \text{Bar}_{\mathcal{O}}^{\text{enh}} : \mathcal{O}\text{-alg}(\mathcal{C}) \rightleftarrows \mathcal{P}\text{-coalg}(\mathcal{C}) : \text{coBar}_{\mathcal{P}}^{\text{enh}},$$

where  $\mathcal{P} = \mathcal{O}^\vee$  and where

$$\text{oblv}_{\mathcal{P}} \circ \text{Bar}_{\mathcal{O}}^{\text{enh}} \simeq \text{Bar}_{\mathcal{O}} \quad \text{and} \quad \text{oblv}_{\mathcal{O}} \circ \text{coBar}_{\mathcal{P}}^{\text{enh}} \simeq \text{coBar}_{\mathcal{P}}.$$

2.2.7. *The pro-nilpotent case.* The pair of adjoint functors  $\text{Bar}_{\mathcal{O}}^{\text{enh}} \dashv \text{coBar}_{\mathcal{P}}^{\text{enh}}$  are not mutually inverses in general. One of the main achievements of [FG11] is to formulate a general condition on  $\mathcal{C}$ , namely, the pro-nilpotence condition, then they are mutually inverses. Let us recall the definition.

**Definition 2.2.8** ([FG11, Defn. 4.1.1]). Let  $\mathcal{C}$  be a (not necessarily unital) stable presentable symmetric monoidal category. We say that  $\mathcal{C}$  is pro-nilpotent if it can be exhibited as a limit

$$\mathcal{C} \simeq \lim(\mathcal{C}_0 \leftarrow \mathcal{C}_1 \leftarrow \mathcal{C}_2 \leftarrow \cdots)$$

such that

- (i)  $\mathcal{C}_0 = 0$ ;
- (ii) For every  $i \geq j$ , the transition functor

$$f_{i,j} : \mathcal{C}_i \rightarrow \mathcal{C}_j$$

commutes with limits;<sup>7</sup>

- (iii) For every  $i$ , the restriction of the tensor product  $\mathcal{C}_i \otimes \mathcal{C}_i \rightarrow \mathcal{C}_i$  to  $\ker(f_{i,i-1}) \otimes \mathcal{C}_i$  is null-homotopic.

We have the following result.

**Proposition 2.2.9** ([FG11, Prop. 4.1.2., Lem. 3.3.4, Lem. 4.1.6 (b)]). *When  $\mathcal{C}$  is pro-nilpotent, then the pair of adjoint functors (2.2.6) are mutually inverses. Moreover, they exchange trivial and free objects on both sides.*

<sup>7</sup>They also commute with colimits as per our convention, see §2.1.1.

*Remark 2.2.10.* In this paper, the pro-nilpotent categories that appear will take the following pattern. Let  $\mathcal{C}$  be a monoidal stable infinity category, and  $\mathbb{Z}_+$  the (discrete) monoidal category whose objects are the positive integers and monoidal structure given by addition. Then, the category of graded objects in  $\mathcal{C}$ ,  $\text{Fun}(\mathbb{Z}_+, \mathcal{C})$ , has a natural monoidal structure, which is pro-nilpotent. See §4 for the precise forms of the categories that appear.

2.2.11. *The case of  $\text{coLie}$  and  $\text{ComAlg}$ .* We have the following equivalence of co-operads (see [FG11]):

$$(2.2.12) \quad \text{Lie}^\vee \simeq \text{ComCoAlg}[1],$$

where

$$\text{ComCoAlg}[1](n) \simeq \Lambda[n-1]$$

is equipped with the sign action of the symmetric group  $\Sigma_n$ . This give rise to the Koszul duality between Lie algebras, and cocommutative coalgebras, originally discovered by Quillen.

2.2.13. In general, for any co-operad  $\mathcal{P}$ ,

$$\mathcal{P}[1](n) \simeq \mathcal{P}(n)[n-1].$$

Equivalently, the functor

$$[1] : \mathcal{C} \rightarrow \mathcal{C}$$

gives rise to an equivalence of categories

$$[1] : \mathcal{P}[1]\text{-coalg}(\mathcal{C}) \simeq \mathcal{P}\text{-coalg}(\mathcal{C}).$$

2.2.14. Dual to (2.2.12) is the following equivalence of operads

$$\text{ComAlg}^\vee \simeq \text{coLie}[1],$$

which we will make use of extensively in this paper.

Note that from what we said above, the functor

$$[1] : \mathcal{C} \rightarrow \mathcal{C}$$

gives rise to an equivalence of categories

$$[1] : \text{coLie}[1](\mathcal{C}) \simeq \text{coLie}(\mathcal{C}).$$

2.2.15. This gives us the following diagram which summarizes the situation for the case of  $\text{ComAlg}$  and  $\text{coLie}$ .

$$\begin{array}{ccc}
 \text{ComAlg}(\mathcal{C}) & \begin{array}{c} \xleftarrow{\text{Bar}_{\text{ComAlg}}} \\ \xrightarrow{\text{coBar}_{\text{coLie}[1]}} \end{array} & \text{coLie}[1](\mathcal{C}) \\
 \begin{array}{c} \uparrow [-1] \\ \downarrow [1] \end{array} & \begin{array}{c} \swarrow \text{coPrim}[1] \\ \searrow \text{coChev} \end{array} & \begin{array}{c} \uparrow [-1] \\ \downarrow [1] \end{array} \\
 \text{ComAlg}[-1](\mathcal{C}) & \begin{array}{c} \xleftarrow{\text{Bar}_{\text{ComAlg}[-1]}} \\ \xrightarrow{\text{coBar}_{\text{coLie}}} \end{array} & \text{coLie}(\mathcal{C})
 \end{array}$$

As indicated in the diagram, we usually use  $\text{coChev}$  to denote

$$[-1] \circ \text{coBar}_{\text{coLie}} \simeq \text{coBar}_{\text{coLie}[1]} \circ [-1]$$

and  $\text{coPrim}[1]$  to denote

$$[1] \circ \text{Bar}_{\text{ComAlg}} \simeq \text{Bar}_{\text{ComAlg}[-1]} \circ [1].$$

2.2.16. Since a trivial Lie-algebra (one with trivial brackets) is usually called an abelian Lie-algebra. In this paper, we will adopt the same convention for  $\text{coLie}$ -coalgebras. Namely, the phrases trivial/abelian  $\text{coLie}$ -coalgebras are used interchangeably.

**2.3. Unital vs non-unital.** Because of the way our operad  $\text{ComAlg}$  is defined,  $\text{ComAlg}(\mathcal{C})$  is the category of *non-unital* commutative algebras. In particular, for any object  $c \in \mathcal{C}$ ,

$$\text{Free}_{\text{ComAlg}} c = \bigoplus_{n>0} \text{Sym}^n c = (\text{Sym } c)_+.$$

Compared to the usual (i.e. unital) commutative algebras, non-unital ones are somewhat less used, and moreover, many constructions are more naturally implemented in the unital world. In this subsection, we will give a quick review of the relations between the two.

**2.3.1.** When  $\mathcal{C}$  is (unital) symmetric monoidal, we use  $\text{ComAlg}^{\text{un,au}}(\mathcal{C})$  to denote the category of augmented unital commutative algebra objects in  $\mathcal{C}$ , and we have an equivalence of categories

$$\text{addUnit} : \text{ComAlg}(\mathcal{C}) \rightleftarrows \text{ComAlg}^{\text{un,au}}(\mathcal{C}) : (-)_+$$

given by formally adjoining a unit in one direction and taking the augmentation ideal the other direction.

**2.3.2.** In general, when  $\mathcal{C}$  is not unital, but it is a full sub-monoidal category of a unital symmetric monoidal stable  $\infty$ -category  $\mathcal{C}^{\text{un}}$ , we let

$$\text{ComAlg}^{\text{un,au}}(\mathcal{C})$$

be the category of unital augmented commutative algebra objects in  $\mathcal{C}^{\text{un}}$  whose augmentation ideal lies in  $\mathcal{C}$ . Then, by the same reasoning as above, we have an equivalence of categories

$$\text{addUnit} : \text{ComAlg}(\mathcal{C}) \rightleftarrows \text{ComAlg}^{\text{un,au}}(\mathcal{C}) : (-)_+$$

given by formally adjoining the unit and removing the unit.

Note that a priori,  $\text{ComAlg}^{\text{un,au}}(\mathcal{C})$  depends on the embedding  $\mathcal{C} \subset \mathcal{C}^{\text{un}}$ . However, the equivalence above implies that this is not the case. This is why we choose to suppress  $\mathcal{C}^{\text{un}}$  from the notation.

**2.3.3.** One of the common procedures we perform with commutative algebras is taking relative tensor products. Let

$$\begin{array}{ccc} \mathcal{A}_1 & \longrightarrow & \mathcal{A}_2 \\ \downarrow & & \\ & & \mathcal{A}_3 \end{array}$$

be a diagram in  $\text{ComAlg}(\mathcal{C})$ . The pushout of this diagram in  $\text{ComAlg}(\mathcal{C})$ , denoted by  $\mathcal{A}_2 \sqcup_{\mathcal{A}_1} \mathcal{A}_3$ , could be computed as follows

$$\mathcal{A}_2 \sqcup_{\mathcal{A}_1} \mathcal{A}_3 \simeq (\text{addUnit}(\mathcal{A}_2) \otimes_{\text{addUnit}(\mathcal{A}_1)} \text{addUnit}(\mathcal{A}_3))_+.$$

Here, the tensor on the right hand side is the usual relative tensor of unital commutative algebras. Note that for unital commutative algebras, relative tensors implement pushouts.

**2.3.4.** Let  $\mathcal{C}$  be a pro-nilpotent category. As mentioned above, we have the following equivalence of categories

$$\text{coPrim}[1] : \text{ComAlg}(\mathcal{C}) \rightleftarrows \text{coLie}(\mathcal{C}) : \text{coChev}.$$

Suppose  $\mathcal{C}$  is a full sub-monoidal-category of  $\mathcal{C}^{\text{un}}$  as above. We will use the notation

$$\text{coChev}^{\text{un}} = \text{addUnit} \circ \text{coChev}$$

to denote the unital cohomological Chevalley functor. In particular, when  $\mathfrak{a}$  is a trivial  $\text{coLie}$ -coalgebra (i.e. trivial co-multiplication maps), we have

$$\text{coChev}^{\text{un}} \mathfrak{a} \simeq \text{addUnit} \circ \text{Free}_{\text{ComAlg}}(\mathfrak{a}[-1]) = \text{Sym}(\mathfrak{a}[-1]).$$

2.4. **Prestacks.** The theory of sheaves on prestacks has been developed in [GL14, Gai15a]. In this subsection and the next, we will give a brief review of this theory. We will state them as facts, without any proof, which, unless otherwise specified, could all be found in [Gai15a].

Prestacks are needed to set up the theory of factorization algebra and factorization homology. We will only make use of this theory in §6 and §7, where it is needed to bridge coLie-cohomology and cohomology of the spaces  $Z_n^d(X)$  mentioned in the introduction. At the first pass, the reader should feel free to treat it as a blackbox.

2.4.1. A prestack is a contravariant functor from  $\text{Sch}$  to  $\text{Spc}$ , the  $\infty$ -category of spaces. In other words, a prestack  $\mathcal{Y}$  is a functor

$$\mathcal{Y} : \text{Sch}^{\text{op}} \rightarrow \text{Spc}.$$

Let  $\text{PreStk}$  be the  $\infty$ -category of prestacks. By Yoneda's lemma, we have a fully-faithful embedding

$$\text{Sch} \hookrightarrow \text{PreStk}.$$

2.4.2. Direct from the definition, any prestack  $\mathcal{Y}$  can be written as a colimit of schemes

$$\mathcal{Y} \simeq \text{colim}_{i \in I} Y_i.$$

2.4.3. A morphism

$$\mathcal{X} \rightarrow \mathcal{Y}$$

between prestacks is said to be *schematic* if its pullback along any scheme

$$S \rightarrow \mathcal{Y}$$

is a scheme.

2.5. **Sheaves on prestacks.** As mentioned above, proofs of all the results in this subsection, unless otherwise specified, could be found in [Gai15a].

2.5.1. *Sheaves on schemes.* For a scheme  $S$ ,

- (i) when the ground field is  $\mathbb{C}$ , and  $\Lambda$  is an arbitrary field of characteristic 0, we take  $\text{Shv}(S)$  to be the ind-completion of the category of constructible sheaves on  $S$  with  $\Lambda$ -coefficients.
- (ii) for any algebraically closed ground field  $k$  in general, and  $\Lambda = \mathbb{Q}_\ell, \overline{\mathbb{Q}}_\ell$  with  $\ell \neq \text{char } k$ , we take  $\text{Shv}(S)$  to be the ind-completion of the category of constructible  $\ell$ -adic sheaves on  $S$  with  $\Lambda$ -coefficients. See also [GL14, §4], [LZ12], and [LZ14].

For any morphism between schemes

$$f : S_1 \rightarrow S_2,$$

we have the following pairs of adjoint functors

$$\begin{aligned} f_! : \text{Shv}(S_1) &\rightleftarrows \text{Shv}(S_2) : f^! \\ f^* : \text{Shv}(S_2) &\rightleftarrows \text{Shv}(S_1) : f_* \end{aligned}$$

Moreover, we also have the box-product  $\boxtimes$ , and hence,  $\otimes$  and  $\overset{!}{\otimes}$ .

2.5.2. When  $X = \text{pt} = \text{Spec } k$ , we use

$$(2.5.3) \quad \text{Vect} = \text{Shv}(\text{pt})$$

to denote the (infinity derived) category of chain complexes in vector spaces over  $\Lambda$ .

2.5.4. *Sheaves on prestacks.* The theory of sheaves on scheme provides us with a functor

$$\mathrm{Shv} : \mathrm{Sch}^{\mathrm{op}} \rightarrow \mathrm{DGCat}_{\mathrm{pres}, \mathrm{cont}},$$

where we use the  $!$ -pullback functor to move between schemes.

We can right Kan extend this functor along the Yoneda embedding

$$\mathrm{Sch}^{\mathrm{op}} \hookrightarrow \mathrm{PreStk}^{\mathrm{op}}$$

to obtain a functor

$$\mathrm{Shv} : \mathrm{PreStk}^{\mathrm{op}} \rightarrow \mathrm{DGCat}_{\mathrm{pres}, \mathrm{cont}}.$$

By formal reason, the functor

$$\mathrm{Shv} : \mathrm{PreStk}^{\mathrm{op}} \rightarrow \mathrm{DGCat}_{\mathrm{pres}, \mathrm{cont}}$$

preserves limits, i.e.

$$\mathrm{Shv}(\mathrm{colim}_i \mathcal{Y}_i) \simeq \lim_i \mathrm{Shv}(\mathcal{Y}_i).$$

In particular, if a prestack

$$\mathcal{Y} \simeq \mathrm{colim}_i Y_i$$

is a colimit of schemes, then

$$\mathrm{Shv}(\mathcal{Y}) \simeq \lim_i \mathrm{Shv}(Y_i).$$

2.5.5. Unwinding the definition of limits, informally, an object  $\mathcal{F} \in \mathrm{Shv}(\mathcal{Y})$  is the same as the following data

(i) A sheaf  $\mathcal{F}_{S, \mathcal{Y}} \in \mathrm{Shv}(S)$  for each  $S \in \mathrm{Sch}$  and  $\mathcal{Y} : S \rightarrow \mathcal{Y}$  (i.e.  $\mathcal{Y} \in \mathcal{Y}(S)$ ).

(ii) An equivalence of sheaves  $\mathcal{F}_{S', f(\mathcal{Y})} \rightarrow f^! \mathcal{F}_{S, \mathcal{Y}}$  for each morphism of schemes  $f : S' \rightarrow S$ .

Moreover, we require that this assignment satisfies a homotopy-cherent system of compatibilities.

2.5.6. Directly from the definition, for any morphism of prestacks

$$f : \mathcal{X} \rightarrow \mathcal{Y},$$

we obtain a functor

$$f^! : \mathrm{Shv}(\mathcal{Y}) \rightarrow \mathrm{Shv}(\mathcal{X})$$

which commutes with both limits and colimits. In particular,  $f^!$  admits a left adjoint  $f_!$ .

2.5.7. The functor  $f_!$  is generally not computable. However, when the target of  $f$  is a scheme, i.e.

$$f : \mathcal{Y} \rightarrow S,$$

where  $S$  is a scheme, and suppose that

$$\mathcal{Y} \simeq \mathrm{colim}_i Y_i.$$

Then,

$$f_! \mathcal{F} \simeq \mathrm{colim}_i f_{i!} \mathcal{F}_{Y_i}$$

where

$$f_i : Y_i \rightarrow \mathcal{Y} \rightarrow S$$

and  $\mathcal{F}_{Y_i}$  is the  $!$ -pullback of  $\mathcal{F}$  to  $Y_i$ .

2.5.8. More generally, suppose  $f : \mathcal{X} \rightarrow \mathcal{Y}$  is a pseudo-proper morphism (briefly reviewed below), then the functor  $f_!$  satisfies base-change with respect to  $g^!$  for any morphism of prestacks  $g : \mathcal{Y}' \rightarrow \mathcal{Y}$ . In particular, we can take  $S$  to be a scheme, and reduce to the situation above. Here, by pseudo-properness, we mean that the base change of  $f$  to any scheme  $S$  is a colimit of schemes proper over  $S$ .

2.5.9. For a general morphism between prestacks

$$f : \mathcal{X} \rightarrow \mathcal{Y},$$

the adjoint pair  $f^* \dashv f_*$  between the associated categories of sheaves is not defined. However, when  $f$  is schematic, then the adjoint pair exists and computable (see [Ho17, §2.6]).

The most important property of  $f_*$  is that it satisfies base change. Namely for any pullback square of prestacks

$$\begin{array}{ccc} \mathcal{X}' & \xrightarrow{g} & \mathcal{X} \\ f \downarrow & & \downarrow f \\ \mathcal{Y}' & \xrightarrow{g} & \mathcal{Y} \end{array}$$

where  $\mathcal{X} \rightarrow \mathcal{Y}$ , and hence, also  $\mathcal{X}' \rightarrow \mathcal{Y}'$ , is schematic, and for any  $\mathcal{F} \in \mathrm{Shv}(\mathcal{X})$ , we have a natural equivalence

$$g^! f_* \mathcal{F} \simeq f_* g^! \mathcal{F}$$

2.5.10. In particular, if

$$\mathcal{Y} = \mathrm{colim}_i \mathcal{Y}_i$$

where  $\mathcal{Y}_i$ 's are schemes, then for each  $i$ , the pullback diagram above becomes

$$\begin{array}{ccc} X_i & \xrightarrow{g_i} & \mathcal{X} \\ f \downarrow & & \downarrow f \\ Y_i & \xrightarrow{g_i} & \mathcal{Y} \end{array}$$

where  $X_i$  is also a scheme. The object  $f_* \mathcal{F} \in \mathrm{Shv}(\mathcal{Y})$  is determined by its  $!$ -pullbacks to the  $\mathcal{Y}_i$ 's, which is computed in terms of the usual  $*$ -pushforward. Namely:

$$g_i^! f_* \mathcal{F} \simeq f_* g_i^! \mathcal{F}$$

and the  $f_*$  on the RHS is the usual pushforward of sheaves between schemes.

2.5.11. The functor  $f^*$  is slightly more complicated to describe. However, when

$$f : \mathcal{Y}_1 \rightarrow \mathcal{Y}_2$$

is étale, which is the case where we need, we have a natural equivalence (see [Ho17, Prop. 2.7.3])

$$(2.5.12) \quad f^! \simeq f^*.$$

2.5.13. *Monoidal structure.* The theory of sheaves on prestacks inherits the monoidal structure (i.e. box tensor) from the theory of sheaves on schemes. Namely, let  $\mathcal{F}_i \in \mathrm{Shv}(\mathcal{Y}_i)$  where  $\mathcal{Y}_i$ 's are prestacks, for  $i = 1, 2$ . Then, the sheaf  $\mathcal{F}_1 \boxtimes \mathcal{F}_2$  is characterized by the condition that for any pair of maps  $f_i : S_i \rightarrow \mathcal{Y}_i$  where the  $S_i$ 's are schemes, we have

$$(f_1 \times f_2)^! (\mathcal{F}_1 \boxtimes \mathcal{F}_2) \simeq f_1^! \mathcal{F}_1 \boxtimes f_2^! \mathcal{F}_2.$$

As in the theory of sheaves for schemes, restricting along the diagonal map lets us define  $\overset{!}{\otimes}$  in general, and  $\otimes^*$  when the diagonal map is schematic.



**2.6. The Ran space/prestack.** The Ran space plays the central role in the theory of factorization algebras and homology. The version of the Ran space used in this paper is that of a colored/graded Ran space. In this section, however, we will review the usual/non-graded Ran space to familiarize the readers with the kinds of structure involved. The colored version will be reviewed in §6.

All results in this section could be found in [FG11, GL14, Gai15a]. The readers are referred to these papers for a full account of the theory. The topologically inclined readers could also find an intuitive introduction in [Ho17].

**2.6.1.** For a scheme  $X \in \text{Sch}$ , we will use  $\text{Ran } X$  to denote the prestack parametrizing non-empty finite subsets of  $X$ . More precisely, for each scheme  $S \in \text{Sch}$ ,

$$(\text{Ran } X)(S) = \{\text{non-empty finite subsets of } X(S)\}.$$

Equivalently, we have

$$\text{Ran } X = \text{colim}_{I \in \text{fSet}^{\text{surj}, \text{op}}} X^I$$

where  $\text{fSet}^{\text{surj}}$  is the category of non-empty finite sets.

**2.6.2.  $\otimes^*$ -monoidal structure.** The Ran space has a natural (non-unital) monoidal structure given by the union map

$$\text{union} : \text{Ran } X \times \text{Ran } X \rightarrow \text{Ran } X.$$

This allows us to define the  $\otimes^*$ -monoidal structure on  $\text{Shv}(\text{Ran } X)$  given by:

$$\mathcal{F} \otimes^* \mathcal{G} = \text{union}_!(\mathcal{F} \boxtimes \mathcal{G}).$$

**2.6.3. Commutative factorizable sheaves.** Using  $\otimes^*$ -monoidal structure on  $\text{Shv}(\text{Ran } X)$ , one can make sense of the categories of algebras and coalgebras on it. The one that is relevant to us is

$$\text{ComAlg}^*(\text{Ran } X) = \text{ComAlg}(\text{Shv}(\text{Ran } X)^{\otimes^*}),$$

the category of commutative algebra objects in  $\text{Shv}(\text{Ran } X)$  with respect to the  $\otimes^*$ -monoidal structure.

**2.6.4.** For any (positive) integer  $n$ , let

$$j : (\text{Ran } X)_{\text{disj}}^n \rightarrow (\text{Ran } X)^n$$

where  $(\text{Ran } X)_{\text{disj}}^n$  is the open sub-prestack of  $(\text{Ran } X)^n$  such that for any scheme  $S$ ,  $(\text{Ran } X)_{\text{disj}}^n(S)$  consists of  $n$  non-empty subsets of  $X(S)$  whose graphs are pair-wise disjoint.

Let  $\mathcal{A} \in \text{ComAlg}^*(\text{Ran } X)$ , then for any  $n$ , we have the multiplication map

$$\text{union}_!(\mathcal{A} \boxtimes \cdots \boxtimes \mathcal{A}) = \mathcal{A} \otimes^* \mathcal{A} \otimes^* \cdots \otimes^* \mathcal{A} \rightarrow \mathcal{A}$$

which is equivalent to a map

$$\mathcal{A} \boxtimes \cdots \boxtimes \mathcal{A} \rightarrow \text{union}^! \mathcal{A}.$$

Applying  $j^!$  to both sides, we get a map of sheaves on  $(\text{Ran } X)_{\text{disj}}^n$

$$(2.6.5) \quad j^!(\mathcal{A} \boxtimes \cdots \boxtimes \mathcal{A}) \rightarrow (j \circ \text{union})^! \mathcal{A}.$$

**Definition 2.6.6.** Let  $\mathcal{A} \in \text{ComAlg}^*(\text{Ran } X)$ . We say that  $\mathcal{A}$  is a commutative factorization algebra if the map (2.6.5) is an equivalence for each  $n$ . We use  $\text{Fact}^*(X)$  to denote the full subcategory of  $\text{ComAlg}^*(\text{Ran } X)$  consisting of commutative factorization algebras.

### 3. $\text{coBar}_{\mathcal{P}}$ AS A SEQUENTIAL LIMIT

The key step in the proof of the homological stability of cohomological Chevalley complexes is the reduction to the case where the  $\text{coLie}$ -coalgebra is trivial. Since the functor  $\text{coChev}$ , or more generally, the functor  $\text{coBar}_{\mathcal{P}}$  for a co-operad  $\mathcal{P}$ , is defined abstractly using operadic Koszul duality, it is too opaque to study directly conveniently. In the dual situation, where one wants to understand the functor  $\text{Bar}_{\mathcal{O}}$  for an operad  $\mathcal{O}$ , a general strategy was given in [GR17], which goes by the name “the  $\text{addFil}$  trick”, and which allows us, in many cases, to reduce the study of  $\text{Bar}_{\mathcal{O}} A$  for any  $\mathcal{O}$ -algebra  $A$  to the case where  $A$  has trivial multiplication maps (i.e. trivial algebra structure).

Unfortunately, the strategy presented in [GR17, §IV.2.1, §IV.2.2] cannot be used directly. The main difficulty in the co-operad case lies mainly in the fact that the functor  $\text{coBar}_{\mathcal{P}}$  is defined using limits, and most of our functors do not commute with limits.

A similar difficulty appears in [Ho16]. There, it was overcome by imposing a certain homological stability condition with respect to a certain  $t$ -structure to make sure that the limits are well-behaved. In this section, we will give a variation of the strategy there. In our case, the situation is better, since our (co-)algebras live in a pro-nilpotent category, and so, our stability condition does not involve any  $t$ -structure, but instead, is expressed in terms of the limit defining the pro-nilpotent category. As a result, the final statement works without any restriction on the object.

The main result of this section is Proposition 3.3.5, which says that inside a pro-nilpotent category,  $\text{coBar}_{\mathcal{P}} A$  could be written as a sequential limit, whose successive fibers are naturally identified with the free  $\mathcal{P}^{\vee}$  object generated  $A$  itself. In the case where  $\mathcal{P} = \text{coLie}$ , the free object is a symmetric algebra, which is easy to investigate.

In §3.1, we set up the general frame work of co-filtered and graded objects, which helps formalizing the notion of taking limits and successive fibers (which we will call associated-graded below) mentioned above. In §3.2, we will introduce the stability and decay conditions on co-filtered and graded objects respectively. These technical conditions enable us to have control over infinite limits. We finally prove our main results in §3.3.

**3.1. Co-filtered and graded objects.** In this subsection, we will introduce the notations for co-filtered and graded objects as well as basic facts about them. These are used to formulate the idea of taking limit and associated-graded. The discussion here is parallel to that of [GR17, §IV.2.1].

**Notation 3.1.1.** Let  $\mathcal{C}$  be a stable  $\infty$ -category. Then the category of co-filtered objects in  $\mathcal{C}$  is defined to be the category of functors

$$\mathcal{C}^{\text{coFil}} = \text{Fun}(\mathbb{Z}^{\rightarrow, \text{op}}, \mathcal{C}),$$

and the category of graded objects

$$\mathcal{C}^{\text{gr}} = \text{Fun}(\mathbb{Z}, \mathcal{C}).$$

Here, the notation  $\mathbb{Z}^{\rightarrow}$  is used to denote the category obtained from considering the usual ordering of  $\mathbb{Z}$ .

**Notation 3.1.2.** We will use  $\mathcal{C}^{\text{coFil}^{>0}}$  and  $\mathcal{C}^{\text{gr}^{>0}}$  to denote the full subcategories of  $\mathcal{C}^{\text{coFil}}$  and  $\mathcal{C}^{\text{gr}}$  respectively of co-filtered and graded objects concentrated in only positive degrees.

3.1.3. We define

$$\text{ass-gr} : \mathcal{C}^{\text{coFil}} \rightarrow \mathcal{C}^{\text{gr}}$$

to be a functor of taking the associated graded object

$$\text{ass-gr}(V)_n = \text{Fib}(V_n \rightarrow V_{n-1}).$$

*Remark 3.1.4.* Later on, we will consider another functor of taking associated graded object of a filtered (rather than co-filtered) object. That functor will be defined as a  $\text{coFib}$  rather than  $\text{Fib}$ . It should be clear from the context which functor we are using.

3.1.5. We define

$$\text{oblv}_{\text{coFil}} : \mathcal{C}^{\text{coFil}} \rightarrow \mathcal{C}$$

to be the functor of taking limit

$$\text{oblv}_{\text{coFil}}(V) = \lim_{n \in \mathbb{Z}^{\rightarrow, \text{op}}} V_n,$$

where

$$V = (\cdots V_{n+1} \rightarrow V_n \rightarrow V_{n-1} \rightarrow \cdots) \in \mathcal{C}^{\text{coFil}}.$$

Similarly, we define

$$\prod : \mathcal{C}^{\text{gr}} \rightarrow \mathcal{C}$$

to be the functor of taking product

$$\prod(V) = \prod_{n \in \mathbb{Z}} V_n$$

where  $V = (V_n)_{n \in \mathbb{Z}} \in \mathcal{C}^{\text{gr}}$ .

Note that both functors can be implemented as the right Kan extensions along  $\mathbb{Z}^{\rightarrow, \text{op}} \rightarrow \text{pt}$  and  $\mathbb{Z} \rightarrow \text{pt}$  respectively, where  $\text{pt}$  is the trivial category.

3.1.6. When  $\mathcal{C}$  is a symmetric monoidal category, then so are  $\mathcal{C}^{\text{coFil}}$  and  $\mathcal{C}^{\text{gr}}$  and the functor  $\text{ass-gr}$  is symmetric monoidal. Note that here, we use the right Kan extension to construct these monoidal structures, i.e.

$$(3.1.7) \quad (V \hat{\otimes} W)_n \simeq \lim_{i+j \leq n} V_i \otimes W_j \in \mathcal{C}^{\text{coFil}}$$

when  $V, W \in \mathcal{C}^{\text{coFil}}$  and

$$(3.1.8) \quad (V \hat{\otimes} W)_n \simeq \prod_{i+j=n} V_i \otimes W_j.$$

We use the notation  $\hat{\otimes}$  to distinguish the monoidal structures obtained by taking limits from the usual ones.

3.1.9. The limits appearing in (3.1.7) and (3.1.8) are infinite in general. However, restricted to  $\mathcal{C}^{\text{coFil}^{>0}}$  and  $\mathcal{C}^{\text{gr}^{>0}}$ , these are finite limits. The monoidal products in these cases, therefore, commute with colimits in each variable. Moreover, in the case of  $\mathcal{C}^{\text{gr}^{>0}}$ , the product is equivalent to a finite direct sum, which is how the usual  $\otimes$ -monoidal structure on  $\mathcal{C}^{\text{gr}^{>0}}$  is defined. In other words,  $\hat{\otimes} \simeq \otimes$  for  $\mathcal{C}^{\text{gr}^{>0}}$ .

3.1.10. The functor  $\text{oblv}_{\text{coFil}}$  is not symmetric monoidal in general, even when we restrict ourselves to the case of  $\mathcal{C}^{\text{coFil}^{>0}}$ , due to the fact that

$$\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$$

does not commute with infinite limits in each variable in general. Indeed,

$$(3.1.11) \quad \text{oblv}_{\text{coFil}}(V \hat{\otimes} W) \simeq \lim_{n \in \mathbb{Z}^{\rightarrow, \text{op}}} (V \hat{\otimes} W)_n \simeq \lim_{n \in \mathbb{Z}^{\rightarrow, \text{op}}} \lim_{i+j \leq n} V_i \otimes W_j \simeq \lim_{i,j \in \mathbb{Z}^{\rightarrow, \text{op}}} V_i \otimes W_j,$$

but the last term is not equivalent to

$$(\lim_{i \in \mathbb{Z}^{\rightarrow, \text{op}}} V_i) \otimes (\lim_{j \in \mathbb{Z}^{\rightarrow, \text{op}}} W_j) \simeq (\text{oblv}_{\text{coFil}} V) \otimes (\text{oblv}_{\text{coFil}} W)$$

in general.

3.1.12. Using the same reasoning, we see that the functor  $\prod$  is not symmetric monoidal in general.

3.1.13. These are the main reasons why we cannot directly apply the  $\text{addFil}$  trick of [GR17] in this setting. The fix is to consider certain full sub-categories of  $\mathcal{C}^{\text{coFil}}$  and  $\mathcal{C}^{\text{gr}}$  which satisfy some stability/finiteness condition to ensure that the limits are well-behaved. This is exactly what we are doing next.

**3.2. Stabilizing and decaying sequences.** We will now introduce the stability conditions required to make the limits involved well-behaved.

3.2.1. Let  $\mathcal{C}$  be a pro-nilpotent category (see Definition 2.2.8). In particular, it can be exhibited as a limit

$$(3.2.2) \quad \mathcal{C} \simeq \lim_{p \in \mathbb{N}^{\text{op}}} \mathcal{C}_p,$$

where all the functors involved in the limit are symmetric monoidal and commute with both limits and colimits.

For each  $p \in \mathbb{N}$ , we will use

$$\text{ev}_p : \mathcal{C} \rightarrow \mathcal{C}_p$$

to denote the canonical projection/evaluation functor. Note that this functor is symmetric monoidal, and moreover, it commutes with both limits and colimits.

For an object  $V$  in  $\mathcal{C}^{\text{coFil}}$  or  $\mathcal{C}^{\text{gr}}$ , we use  $V_{n,p}$  to denote  $\text{ev}_p(V_n)$ .

**Definition 3.2.3.** Let  $\mathcal{C}$  be as above. Then, a co-filtered object  $V \in \mathcal{C}^{\text{coFil}^{>0}}$  is said to stabilize if for all  $p$ , the map

$$(\text{ev}_p(V_{n+1}) \rightarrow \text{ev}_p(V_n)) = \text{ev}_p(V_{n+1} \rightarrow V_n)$$

is an equivalence when  $n \gg 0$ .

A graded object  $V \in \mathcal{C}^{\text{gr}^{>0}}$  is said to be decaying if for all  $p$ ,

$$\text{ev}_p(V_n) \simeq 0$$

when  $n \gg 0$ .

**Notation 3.2.4.** We use  $\mathcal{C}^{\text{coFil}^{>0}, \text{stab}}$  and  $\mathcal{C}^{\text{gr}^{>0}, \text{decay}}$  to denote the full-subcategories of  $\mathcal{C}^{\text{coFil}^{>0}}$  and  $\mathcal{C}^{\text{gr}^{>0}}$  consisting of stabilizing and decaying objects respectively.

*Remark 3.2.5.* Note that the notions of stabilizing and decaying objects depend on the presentation of  $\mathcal{C}$  as a limit as in (3.2.2). We will elide this dependence from the notation unless confusion is likely to happen.

The following result is tautological.

**Lemma 3.2.6.** Let  $\mathcal{C}$  be as above and  $V \in \mathcal{C}^{\text{coFil}^{>0}}$ . Then

$$V \in \mathcal{C}^{\text{coFil}^{>0}, \text{stab}}$$

if and only if

$$\text{ass-gr}(V) \in \mathcal{C}^{\text{gr}^{>0}, \text{decay}}.$$

As a consequence, we have the following

**Corollary 3.2.7.** The categories  $\mathcal{C}^{\text{gr}^{>0}, \text{decay}}$  and  $\mathcal{C}^{\text{coFil}^{>0}, \text{stab}}$  are closed under tensor products in  $\mathcal{C}^{\text{gr}^{>0}}$  and  $\mathcal{C}^{\text{coFil}^{>0}}$  respectively. In other words, they are symmetric monoidal categories.

*Proof.* By Lemma 3.2.6 above and the fact that  $\text{ass-gr}$  is symmetric monoidal, it suffices to show the statement for  $\mathcal{C}^{\text{gr}^{>0}, \text{decay}}$ . However, this is direct from the way tensors are computed in  $\mathcal{C}^{\text{gr}^{>0}}$ . Indeed, let  $V, W \in \mathcal{C}^{\text{gr}^{>0}, \text{decay}}$ . We want to show that for any fixed  $p$ ,

$$(V \hat{\otimes} W)_{n,p} \simeq \prod_{n_1+n_2=n} V_{n_1,p} \otimes W_{n_2,p} \simeq 0$$

when  $n \gg 0$ . But this is clear since when  $n \gg 0$ , either  $n_1 \gg 0$  or  $n_2 \gg 0$ , and we are done.  $\square$

Finally, using the computation at (3.1.11), we immediately get the following result promised earlier.

**Lemma 3.2.8.** The functor

$$\text{oblv}_{\text{coFil}} : \mathcal{C}^{\text{coFil}^{>0}, \text{stab}} \rightarrow \mathcal{C}$$

is symmetric monoidal.

*Proof.* We want to show that for each  $V, W \in \mathcal{C}^{\text{coFil}^{>0}, \text{stab}}$ , the natural map

$$\text{oblv}_{\text{coFil}} V \otimes \text{oblv}_{\text{coFil}} W \rightarrow \text{oblv}_{\text{coFil}}(V \hat{\otimes} W)$$

is an equivalence. It suffices to show that this is an equivalence after evaluating on  $\mathcal{C}_p$  for each  $p$ . But now, the stability condition allows us to turn the limit in the last term of (3.1.11) inside the tensor, since we can just take  $i$  and  $j$  big enough there and remove the limit altogether.  $\square$

3.2.9. The decay condition on  $\mathcal{C}^{\text{gr}^{>0}, \text{decay}}$  also makes the functor

$$\prod : \mathcal{C}^{\text{gr}^{>0}, \text{decay}} \rightarrow \mathcal{C}$$

well-behave.

**Lemma 3.2.10.** *We have the following natural equivalence of functors*

$$\bigoplus \simeq \prod : \mathcal{C}^{\text{gr}^{>0}, \text{decay}} \rightarrow \mathcal{C}.$$

*In particular, the functor  $\prod$  commutes with colimits and is symmetric monoidal*

*Proof.* It suffices to show that we have the following natural equivalence

$$\text{ev}_p \circ \bigoplus \simeq \text{ev}_p \circ \prod$$

for each  $p$ . However, this is clear since within each  $\mathcal{C}_p$ , the products/direct sums that appear are finite.  $\square$

*Remark 3.2.11.* Note that  $\prod$  automatically commutes with limits since it's defined as a limit itself.

3.2.12. To end this subsection, we will record the following piece of notation: we will use

$$\text{addCoFil} : \mathcal{C} \rightarrow \mathcal{C}^{\text{coFil}^{>0}, \text{stab}}$$

to denote the following functor which sends  $V \in \mathcal{C}$  to the co-filtered object

$$\cdots \leftarrow 0 \leftarrow V \leftarrow V \leftarrow \cdots$$

where  $V$ 's first appearance is at degree 1 and all the maps between them are identities.

3.2.13. As in [GR17, §IV.2.1.4], for any co-operad  $\mathcal{P}$ , the functor  $\text{addCoFil}$  upgrades to a functor

$$\text{addCoFil} : \mathcal{P}\text{-coalg}(\mathcal{C}) \rightarrow \mathcal{P}\text{-coalg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}}),$$

and moreover, as in [GR17, Prop. IV.2.1.4.6] the following diagram commutes

$$(3.2.14) \quad \begin{array}{ccccc} \mathcal{P}\text{-coalg}(\mathcal{C}) & \xrightarrow{\text{addCoFil}} & \mathcal{P}\text{-coalg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}}) & \xrightarrow{\text{ass-gr}} & \mathcal{P}\text{-coalg}(\mathcal{C}^{\text{gr}^{>0}, \text{decay}}) \\ \text{oblv}_{\mathcal{P}} \downarrow & & & & \parallel \\ \mathcal{C} & \xrightarrow{\text{deg}=1} & \mathcal{C}^{\text{gr}^{>0}, \text{decay}} & \xrightarrow{\text{cotriv}_{\mathcal{P}}} & \mathcal{P}\text{-coalg}(\mathcal{C}^{\text{gr}^{>0}, \text{decay}}) \end{array}$$

Here,  $(\text{deg} = 1) = \text{ass-gr} \circ \text{addCoFil}$  denotes the functor of placing an object of  $\mathcal{C}$  in graded degree 1.

3.3. **A co-filtration on  $\text{coBar}_{\mathcal{P}}$ .** Using the framework setup above, in this subsection, we will show that for any  $\text{coLie-coalgebra } \mathfrak{a}$  in a pro-nilpotent category  $\mathcal{C}$ ,  $\text{coChev } \mathfrak{a}$  could be computed as a sequential limit

$$\text{coChev } \mathfrak{a} = \lim(\text{coChev}^1 \mathfrak{a} \leftarrow \text{coChev}^2 \mathfrak{a} \leftarrow \cdots) = \text{oblv}_{\text{coFil}}((\text{coChev}^n \mathfrak{a})_{n \in \mathbb{Z}_{>0}})$$

for some

$$(\text{coChev}^n \mathfrak{a})_{n \in \mathbb{Z}_{>0}} \in \mathcal{C}^{\text{coFil}^{>0}, \text{stab}}$$

such that

$$\text{ass-gr}((\text{coChev}^n \mathfrak{a})_{n \in \mathbb{Z}_{>0}}) \simeq (\text{Sym}^n \mathfrak{a}[-1])_{n \in \mathbb{Z}_{>0}} \in \mathcal{C}^{\text{gr}^{>0}, \text{decay}}.$$

In fact, we will show a general statement, Proposition 3.3.5, about  $\text{coBar}_{\mathcal{P}}$  for any co-operad  $\mathcal{P}$ , which specializes to the case of  $\text{coChev}$  described above, Corollary 3.3.6. The following Proposition is the technical statement from which our desired result follows.

**Proposition 3.3.1.** *Let  $\mathcal{C}$  be a pro-nilpotent category,  $\mathcal{P}$  a co-operad with Koszul dual  $\mathcal{O} = \mathcal{P}^\vee$ . Then, the following diagram commutes*

(3.3.2)

$$\begin{array}{ccc} \mathcal{P}\text{-coalg}(\mathcal{C}) & \xrightarrow{\text{coBar}_{\mathcal{P}}} & \mathcal{O}\text{-alg}(\mathcal{C}) \\ \text{addCoFil} \downarrow & & \uparrow \text{oblv}_{\text{coFil}} \\ \mathcal{P}\text{-coalg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}}) & \xrightarrow{\text{coBar}_{\mathcal{P}}} & \mathcal{O}\text{-alg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}}) \\ \text{ass-gr} \downarrow & & \downarrow \text{ass-gr} \\ \mathcal{P}\text{-coalg}(\mathcal{C}^{\text{gr}^{>0}, \text{decay}}) & \xrightarrow{\text{coBar}_{\mathcal{P}}} & \mathcal{O}\text{-alg}(\mathcal{C}^{\text{gr}^{>0}, \text{decay}}) \\ \oplus \simeq \Pi \downarrow & & \downarrow \oplus \simeq \Pi \\ \mathcal{P}\text{-coalg}(\mathcal{C}) & \xrightarrow{\text{coBar}_{\mathcal{P}}} & \mathcal{O}\text{-alg}(\mathcal{C}) \end{array}$$

*Proof.* The first thing we need to check is that this diagram actually makes sense. Namely, the functors involved land in the correct categories. By inspection, we see that the only cases where it is not immediate are the middle two instances of  $\text{coBar}_{\mathcal{P}}$ . By Lemma 3.2.6, it suffices to check the statement for the third instance of  $\text{coBar}_{\mathcal{P}}$  (from top down). For that, it suffices to check that for each

$$A \in \mathcal{P}\text{-coalg}(\mathcal{C}^{\text{gr}^{>0}, \text{decay}})$$

and each  $p \in \mathbb{Z}_+$ ,

$$\text{ev}_p(\text{coBar}_{\mathcal{P}}(A)_n) = \text{coBar}_{\mathcal{P}}(A)_{n,p} \simeq 0$$

when  $n \gg 0$ .

Let  $\text{coBar}_{\mathcal{P}}^\bullet(A)$  be the co-simplicial object defining  $\text{coBar}_{\mathcal{P}}(A)$ . Then since the monoidal structure on  $\mathcal{C}_p$  is nilpotent by assumption, only finite numbers of tensor powers of  $A$  appear in

$$\text{ev}_p(\text{coBar}_{\mathcal{P}}^\bullet(A)).$$

The decay condition on  $A$  then implies that

$$\text{coBar}_{\mathcal{P}}^\bullet(A)_{n,p} \simeq 0$$

when  $n \gg 0$ , and we are done.

Now, observe that the bottom two squares of (3.3.2) commute since the vertical arrows are all symmetric monoidal and co-continuous (i.e. commute with limits). It remains to show that the top square commutes. This follows from the equivalence

$$\text{id} \simeq \text{oblv}_{\text{coFil}} \circ \text{addCoFil}$$

and the following commutative diagram

$$\begin{array}{ccc}
 \mathcal{P}\text{-coalg}(\mathcal{C}) & \xrightarrow{\text{coBar}_{\mathcal{P}}} & \mathcal{O}\text{-alg}(\mathcal{C}) \\
 \text{oblv}_{\text{coFil}} \uparrow & & \uparrow \text{oblv}_{\text{coFil}} \\
 \mathcal{P}\text{-coalg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}}) & \xrightarrow{\text{coBar}_{\mathcal{P}}} & \mathcal{O}\text{-alg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}})
 \end{array}$$

which is due to the fact that  $\text{oblv}_{\text{coFil}}$  is also co-continuous, being a right adjoint, and symmetric monoidal, by Lemma 3.2.8.

This concludes the proof of Proposition 3.3.1.  $\square$

*Remark 3.3.3.* If we remove the stability conditions, the “dual” diagram of (3.3.2) (i.e. replacing co-operad by operad, Bar by coBar etc.) is the one appearing in [GR17], which makes the trick of adding a filtration work. Here, the stability conditions are what needed to make the same diagram commute in the co-operad setting.

3.3.4. We will now come to the main result of this section.

**Proposition 3.3.5.** *Let  $\mathcal{C}$  be a pro-nilpotent category,  $\mathcal{P}$  a co-operad with Koszul dual  $\mathcal{O}$ , and  $A \in \mathcal{P}\text{-coalg}(\mathcal{C})$ . Then,  $\text{coBar}_{\mathcal{P}} A$  could be canonically written as a limit*

$$\text{coBar}_{\mathcal{P}} A \simeq \lim_{n \in \mathbb{Z}_{>0}^{\rightarrow, \text{op}}} \text{coBar}_{\mathcal{P}}^n A$$

for some

$$(\text{coBar}_{\mathcal{P}}^n A)_{n \in \mathbb{Z}_{>0}} \in \mathcal{O}\text{-alg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}})$$

such that

$$\text{ass-gr}((\text{coBar}_{\mathcal{P}}^n A)_{n \in \mathbb{Z}_{>0}}) \simeq \text{Free}_{\mathcal{O}}(\text{oblv}_{\mathcal{P}}(A)^{\deg=1}) \in \mathcal{O}\text{-alg}(\mathcal{C}^{\text{gr}^{>0}, \text{decay}}).$$

*Proof.* The top square of (3.3.2) implies that for any  $A \in \mathcal{P}\text{-coalg}(\mathcal{C})$ ,  $\text{coBar}_{\mathcal{P}} A$  could be written as a limit

$$\text{coBar}_{\mathcal{P}} A \simeq \text{oblv}_{\text{coFil}} \circ \text{coBar}_{\mathcal{P}} \circ \text{addCoFil}(A) \simeq \lim_{n \in \mathbb{Z}_{>0}^{\rightarrow, \text{op}}} (\text{coBar}_{\mathcal{P}}^n A)$$

for some

$$(\text{coBar}_{\mathcal{P}}^n A)_{n \in \mathbb{Z}_{>0}} \in \mathcal{O}\text{-alg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}}).$$

Moreover, the bottom two squares of 3.3.2, coupled with (3.2.14) imply that

$$\text{ass-gr}((\text{coBar}_{\mathcal{P}}^n A)_{n \in \mathbb{Z}_{>0}}) \simeq \text{coBar}_{\mathcal{P}}(\text{cotriv}_{\mathcal{P}} \circ \text{oblv}_{\mathcal{P}}(A)^{\deg=1}) \simeq \text{Free}_{\mathcal{O}}(\text{oblv}_{\mathcal{P}}(A)^{\deg=1}).$$

$\square$

Specializing to the case where  $\mathcal{P} = \text{coLie}$  we get the desired result stated at the beginning of this subsection.

**Corollary 3.3.6.** *Let  $\mathcal{C}$  be a pro-nilpotent category and  $\mathfrak{a} \in \text{coLie}(\mathcal{C})$ . Then  $\text{coChev } \mathfrak{a}$  could be canonically written as a limit*

$$\text{coChev } \mathfrak{a} \simeq \lim_{n \in \mathbb{Z}_{>0}^{\rightarrow, \text{op}}} \text{coChev}^n \mathfrak{a}$$

for some

$$(\text{coChev}^n \mathfrak{a})_{n \in \mathbb{Z}_{>0}} \in \text{ComAlg}(\mathcal{C}^{\text{coFil}^{>0}, \text{stab}})$$

such that

$$\text{ass-gr}((\text{coChev}^n \mathfrak{a})_{n \in \mathbb{Z}_{>0}}) \simeq \text{Sym}(\mathfrak{a}^{\deg=1}[-1])_+ \in \text{ComAlg}(\mathcal{C}^{\text{gr}^{>0}, \text{decay}}).$$



#### 4. SOME GENERALITIES ON HOMOLOGICAL STABILITY

In this section, we set up the general formalism and prove general results needed for the discussion of homological stability. Subsections §4.1–§4.5 set up the language, whereas the remaining subsections §4.6–§4.8 prove various general results about homological stability. The upshot of this section is that stable homology (when homological stability does occur) can be captured by taking the colimit of the filtered object, a process we call stabilization. This colimit is, however, completely general and enjoys nice categorical properties. We should, therefore, think of it as the generalization of stable homology, and study it carefully, even if the ultimate aim is to understand homological stability and stable homology.

We will now give a detailed overview of the section. We start, in §4.1 and §4.2, with the fundamental building block, the notion of a (multi-)graded/filtered object/algebra in a given category. Here, the grading is given by  $\mathbb{Z}_{\geq 0}^m$  rather than just  $\mathbb{Z}$ . In the geometric applications that we have in mind, these gradings come from the natural gradings on the spaces  $(Z_n^d(X))_{d \in \mathbb{Z}_{\geq 0}^m}$ . The discussion here is only a slight extension of [GR17, §IV.2.1.3], where the case of  $\mathbb{Z}$ -graded and filtered case is treated.

In §4.3, we introduce the notion of a graded-unital algebra, a special kind of graded algebras which upgrades to a filtered algebra so that questions about homological stability can be discussed. §4.4 discusses the formalism needed to compare objects filtered/graded by different gradings. Finally, in §4.5, the notion of stabilization of a filtered object is introduced. In particular, in §4.5.12, we explain how one stabilizes graded-unital algebras introduced in §4.3.

The highlight of this section comes in §4.6, which proves that this stabilization captures stable homology (when homological stability does occur). In §4.7, we provide a general strategy for attacking homological stability questions for objects presented as a sequential limit, whose successive fibers are known. Coupled with the result in §3, this will be use in §5 to prove homological stability for the cohomological Chevalley complexes of a large class of coLie-coalgebras by reducing to the case of trivial coLie-coalgebras. In the last subsection §4.8, we provide a calculation of the stable homology in a particularly simple case.

**4.1. Filtered and graded objects.** In this subsection, we introduce the categories of (multi-)filtered and (multi-)graded objects in a given category. These categories provide the natural setting where one can discuss homological stability. The readers might find the discussion here parallel to that of §3.1. Indeed, this subsection deals with a slight generalization of the dual of the picture presented there, and is, in turn, a slight generalization of [GR17, §IV.2.1.3].

4.1.1. We start with the indexing categories. Let

$$\mathbb{Z}_{\geq 0, +}^m = \mathbb{Z}_{\geq 0}^m - \{0\}$$

be the sub-semigroup of  $\mathbb{Z}_{\geq 0}^m$ . It can be viewed as a (discrete) symmetric monoidal category, where the monoidal structure is given by addition.

Consider also the symmetric monoidal category  $\mathbb{Z}_{\geq 0}^{m, \rightarrow}$  whose objects are the same as for  $\mathbb{Z}_{\geq 0}^m$ , morphisms given by the natural partially ordered structure, and the monoidal structure given by the monoid structure on  $\mathbb{Z}_{\geq 0}^m$ . Similarly, consider the full sub-(non-unital) monoidal category

$$\mathbb{Z}_{\geq 0, +}^{m, \rightarrow} = \mathbb{Z}_{\geq 0}^{m, \rightarrow} - \{0\}.$$

4.1.2. We will employ the following notation: for each  $1 \leq k \leq m$ , we will use  $\mathbf{1}_k$  to denote

$$(0, \dots, 0, 1, 0, \dots, 0) \in \mathbb{Z}_{\geq 0}^m$$

where the only “1” appears at the  $k$ -th position.

4.1.3. Recall the following general construction: for any symmetric monoidal categories  $\mathcal{C}$  and  $\Gamma$  with  $\mathcal{C}$  being cocomplete, we can associate to them a new symmetric monoidal category

$$\mathcal{C}^\Gamma = \text{Fun}(\Gamma, \mathcal{C})$$

of all functors from  $\Gamma$  to  $\mathcal{C}$ , where the monoidal structure on  $\mathcal{C}^\Gamma$  is given by Day convolution (see [Lur17a, §2.2.6]). More explicitly, for  $V, W \in \text{Fun}(\Gamma, \mathcal{C})$ , we have following diagram

$$\begin{array}{ccc} \Gamma \times \Gamma & \xrightarrow{(V,W)} & \mathcal{C} \times \mathcal{C} \\ \otimes \downarrow & \searrow & \downarrow \otimes \\ \Gamma & \xrightarrow{V \otimes W} & \mathcal{C} \end{array}$$

where  $V \otimes W$  is defined to be the left Kan extension of the diagonal arrow along the vertical arrow on the left.

*Example 4.1.4.* When  $\Gamma = \mathbb{Z}$  is the (discrete) monoidal category,  $\mathcal{C}^{\mathbb{Z}}$  is the category of graded objects in  $\mathcal{C}$ . Similarly, when  $\Gamma = \mathbb{Z}^{\rightarrow}$  is the monoidal category with morphism given by the ordering on  $\mathbb{Z}$ ,  $\mathcal{C}^{\mathbb{Z}^{\rightarrow}}$  is the category of filtered objects in  $\mathcal{C}$ . These categories are symmetric monoidal categories in the usual way.

4.1.5. In this paper, we will deal with categories of the form

$$(4.1.6) \quad \mathcal{C}_{\geq 0}^m, \quad \mathcal{C}_{\geq 0,+}^m, \quad \mathcal{C}_{\geq 0}^{m,\rightarrow}, \quad \mathcal{C}_{\geq 0,+}^{m,\rightarrow}$$

where  $\mathcal{C}$  is a stable symmetric monoidal category. We will call objects in the first two and the last two categories multi-graded and multi-filtered respectively. To keep the terminology simple, we will simply refer to them as graded and filtered objects, unless confusion might arise.

Note that we have the following inclusion (i.e. fully-faithful embeddings) of categories

$$\mathcal{C}_{\geq 0,+}^m \subset \mathcal{C}_{\geq 0}^m \quad \text{and} \quad \mathcal{C}_{\geq 0,+}^{m,\rightarrow} \subset \mathcal{C}_{\geq 0}^{m,\rightarrow}.$$

For later use, we record the following straightforward lemma.

**Lemma 4.1.7.** *Let  $\mathcal{C}$  be a stable symmetric monoidal category. Then, the symmetric monoidal categories*

$$\mathcal{C}_{\geq 0,+}^m \quad \text{and} \quad \mathcal{C}_{\geq 0,+}^{m,\rightarrow}$$

*are pro-nilpotent.*

4.1.8. When  $\mathcal{C}$  is a unital symmetric monoidal category with monoidal unit  $\mathbb{1}_{\mathcal{C}}$ , then so are  $\mathcal{C}_{\geq 0}^m$  and  $\mathcal{C}_{\geq 0}^{m,\rightarrow}$ . The units are  $\mathbb{1}_{\mathcal{C}}$  in graded degrees  $\mathbf{0}$  in the first case and the constant object  $\mathbb{1}_{\mathcal{C}}$  living in each graded-degree in the second case. The symmetric monoidal categories  $\mathcal{C}_{\geq 0,+}^m$  and  $\mathcal{C}_{\geq 0,+}^{m,\rightarrow}$  are, however, not unital.

4.1.9. *Rees construction.* We have a natural embedding of categories

$$\mathbb{Z}_{\geq 0}^m \hookrightarrow \mathbb{Z}_{\geq 0}^{m,\rightarrow}.$$

Restricting and left-Kan extending along this functor give us a pair of adjoint functors

$$\text{Rees} : \mathcal{C}_{\geq 0}^m \rightleftarrows \mathcal{C}_{\geq 0}^{m,\rightarrow} : \text{Fil} \rightarrow \text{gr}.$$

4.1.10. When  $m = 1$ , the functor Rees is indeed the usual Rees construction: for any object  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^m}$

$$\text{Rees}(V)_d \simeq \bigoplus_{d' \leq d} V_{d'}.$$

**Lemma 4.1.11.** *The functor Rees is symmetric monoidal. As a result,  $\text{Fil} \rightarrow \text{gr}$  is right lax monoidal.*

*Proof.* Since the monoidal structures on both sides are defined by left-Kan extension, Rees is symmetric monoidal, and hence, its right adjoint,  $\text{Fil} \rightarrow \text{gr}$  is right-lax monoidal.  $\square$

The same discussion applies equally well to  $\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m}$  and  $\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^{m, \rightarrow}}$ .

4.1.12. *Associated-graded.* One common way to extract information out of a  $\mathbb{Z}_{\geq 0}^{\rightarrow}$ -filtered objects in a stable  $\infty$ -category is to take its associated-graded. This functor commutes with both limits, colimits, and moreover, it is conservative. In addition, when  $\mathcal{C}$  is symmetric monoidal, so is  $\text{ass-gr}$  (see [GR17, §IV.2.1.3]).

4.1.13. We will now construct a generalization of this functor in the case of multi-filtered objects. First, consider the following functor

$$(\text{gr} \xrightarrow{0} \text{Fil})_{\mathcal{C}, m} : \mathcal{C}^{\mathbb{Z}_{\geq 0}^m} \rightarrow \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$$

which sends an object  $V = (V_d)_{d \in \mathbb{Z}_{\geq 0}^m}$  to a filtered object  $(V_d)_{d \in \mathbb{Z}_{\geq 0}^m}$  where all the maps are 0. Clearly, this functors commutes with both limits and colimits, and hence, it admits a left adjoint, which we will denote by

$$\text{ass-gr}_{\mathcal{C}, m} : \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}} \rightarrow \mathcal{C}^{\mathbb{Z}_{\geq 0}^m}.$$

Unless confusion is likely to occur, we will omit the subscript  $\mathcal{C}$ , or  $m$ , or both, from our notation.

4.1.14. The functor  $\text{ass-gr}_{\mathcal{C}, m}$  we just defined is indeed closely related to the usual functor of taking associated graded. In fact, as we will soon see, it is the functor of taking successive associated graded with respect to all the gradings. This will help us establish nice formal properties of  $\text{ass-gr}_{\mathcal{C}, m}$ .

First, observe that in the case where  $m = 1$ ,

$$\text{ass-gr}_{\mathcal{C}} : \mathcal{C}^{\mathbb{Z}_{\geq 0}^{\rightarrow}} \rightarrow \mathcal{C}^{\mathbb{Z}_{\geq 0}}$$

is indeed the functor of taking the associated-graded object of a filtered object. Namely,

$$\text{ass-gr}(V)_d = \text{coFib}(V_{d-1} \rightarrow V_d)$$

where  $V = (V_d)_{d \in \mathbb{Z}_{\geq 0}} \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{\rightarrow}}$ .

Next, observe that for any  $m$ ,

$$\mathcal{C}^{\mathbb{Z}_{\geq 0}^m} \simeq (\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m-1}})^{\mathbb{Z}_{\geq 0}} \simeq (\mathcal{C}^{\mathbb{Z}_{\geq 0}})^{\mathbb{Z}_{\geq 0}^{m-1}} \quad \text{and} \quad \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}} \simeq (\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m-1, \rightarrow}})^{\mathbb{Z}_{\geq 0}^{\rightarrow}} \simeq (\mathcal{C}^{\mathbb{Z}_{\geq 0}^{\rightarrow}})^{\mathbb{Z}_{\geq 0}^{m-1, \rightarrow}}.$$

Since compositions of left adjoints are left adjoints, we have the following decomposition of  $\text{ass-gr}_{\mathcal{C}, m}$

$$\text{ass-gr}_{\mathcal{C}, m} \simeq \text{ass-gr}_{\mathcal{C}^{\mathbb{Z}_{\geq 0}, m-1}} \circ \text{ass-gr}_{\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m-1, \rightarrow}}},$$

which means that  $\text{ass-gr}_{\mathcal{C}, m}$  could be computed by taking successive associated-graded in the usual sense.

By induction, this immediately gives us the following results.

**Lemma 4.1.15.** *The functor  $\text{ass-gr}_{\mathcal{C}, m}$  commutes with limits, colimits, and is conservative.*

**Lemma 4.1.16.** *When  $\mathcal{C}$  is symmetric monoidal, so is the functor  $\text{ass-gr}_{\mathcal{C}, m}$ .*

4.1.17. The discussion above applies equally well for  $\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m}$  and  $\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^{m, \rightarrow}}$ . We will use the same notation  $\text{ass-gr}_{\mathcal{C}, m}$  to denote the functor of taking associated-graded in this case.

**4.2. Unital and augmented algebras.** For any symmetric monoidal category  $\mathcal{C}$  as above, we let

$$\text{ComAlg}(\mathcal{C}_{\geq 0}^m), \quad \text{ComAlg}(\mathcal{C}_{\geq 0,+}^m), \quad \text{ComAlg}(\mathcal{C}_{\geq 0}^{m,\rightarrow}), \quad \text{ComAlg}(\mathcal{C}_{\geq 0,+}^{m,\rightarrow})$$

be the categories of graded and filtered commutative algebras respectively. Moreover, when the monoidal structure on  $\mathcal{C}$  is unital, we can consider the categories of unital and augmented algebras, denoted by

$$\text{ComAlg}^{\text{au}}(-), \quad \text{ComAlg}^{\text{un}}(-), \quad \text{ComAlg}^{\text{un,au}}(-),$$

where “ $-$ ” is the place-holder for  $\mathcal{C}_{\geq 0}^m$  or  $\mathcal{C}_{\geq 0}^{m,\rightarrow}$ .

**4.2.1.** As we have seen above, the monoidal structures on  $\mathcal{C}_{\geq 0}^m$  is not unital. However, the fully-faithful embedding

$$\mathcal{C}_{\geq 0,+}^m \subset \mathcal{C}_{\geq 0}^m$$

brings us to the setting of §2.3.2, which allows us to talk about the category  $\text{ComAlg}^{\text{un,au}}(\mathcal{C}_{\geq 0,+}^m)$  of unital, augmented commutative algebra objects in  $\mathcal{C}_{\geq 0}^m$  whose augmentation ideal lives in  $\mathcal{C}_{\geq 0,+}^m$ .

**4.2.2.** The various categories of algebras mentioned above naturally contain a specific class of polynomial algebras of interest to us:  $\Lambda[\mathbb{Z}_{\geq 0}^m]$  and its associated non-unital algebra  $\Lambda[\mathbb{Z}_{\geq 0}^m]_+ = \Lambda[\mathbb{Z}_{\geq 0,+}^m]$ . Let us explain what we mean by these notations. Let  $\mathbb{1}_{\mathcal{C}}$  denote the monoidal unit of  $\mathcal{C}$ . By  $\Lambda[\mathbb{Z}_{\geq 0}^m]$ , we mean the free unital commutative algebra in  $\text{ComAlg}(\mathcal{C}_{\geq 0}^m)$  generated by

$$\mathbb{1}_{\mathbf{1}_1} \oplus \mathbb{1}_{\mathbf{1}_2} \oplus \cdots \oplus \mathbb{1}_{\mathbf{1}_m} \in \mathcal{C}_{\geq 0}^m$$

i.e.  $m$  copies of  $\mathbb{1}_{\mathcal{C}}$ , sitting in graded-degrees

$$\mathbf{1}_1, \mathbf{1}_2, \dots, \mathbf{1}_m \in \mathbb{Z}_{\geq 0}^m.$$

Note that  $\Lambda[\mathbb{Z}_{\geq 0}^m] \in \text{ComAlg}^{\text{un,au}}(\mathcal{C}_{\geq 0,+}^m)$ , and its augmentation ideal  $\Lambda[\mathbb{Z}_{\geq 0}^m]_+ \simeq \Lambda[\mathbb{Z}_{\geq 0,+}^m]$  is an object of  $\text{ComAlg}(\mathcal{C}_{\geq 0,+}^m)$ .

**4.2.3.** Similarly, we consider the algebras  $\Lambda[\mathbb{Z}_{\geq 0}^{m,\rightarrow}]$  and  $\Lambda[\mathbb{Z}_{\geq 0,+}^{m,\rightarrow}]$ . At each graded degree  $\mathbf{d}$ ,

$$\Lambda[\mathbb{Z}_{\geq 0}^{m,\rightarrow}]_{\mathbf{d}} \simeq \mathbb{1}_{\mathcal{C},\mathbf{d}}$$

and all the maps (from the filtration) are equivalences.

We see at once from the construction that these algebras go to  $\Lambda[\mathbb{Z}_{\geq 0}^m]$  and  $\Lambda[\mathbb{Z}_{\geq 0,+}^m]$  respectively under  $\text{Fil} \rightarrow \text{gr}$ . Because of this reason, unless confusion is likely to occur, we will use  $\Lambda[\mathbb{Z}_{\geq 0}^m]$  and  $\Lambda[\mathbb{Z}_{\geq 0,+}^m]$  in place of  $\Lambda[\mathbb{Z}_{\geq 0}^{m,\rightarrow}]$  and  $\Lambda[\mathbb{Z}_{\geq 0,+}^{m,\rightarrow}]$  respectively.

*Remark 4.2.4.* Note that  $\Lambda[\mathbb{Z}_{\geq 0}^m] \in \text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0}^{m,\rightarrow}})$  is the monoidal unit of the category  $\mathcal{C}_{\geq 0}^{m,\rightarrow}$ , and is, therefore, a commutative algebra object. We can thus alternatively define  $\Lambda[\mathbb{Z}_{\geq 0}^m] \in \text{ComAlg}(\mathbb{Z}_{\geq 0}^m)$  as  $(\text{Fil} \rightarrow \text{gr})(\mathbb{1}_{\mathcal{C}_{\geq 0}^{m,\rightarrow}})$ , which is also a commutative algebra object, since  $\text{Fil} \rightarrow \text{gr}$  is right-lax monoidal.

*Remark 4.2.5.* In the case where  $\mathcal{C} = \text{Vect}$  (see (2.5.3) for the conventions),  $\Lambda[\mathbb{Z}_{\geq 0}^m] \in \text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0}^m})$  is precisely the polynomial algebra generated by  $m$  variables sitting in appropriate graded-degrees. For the applications of this paper, it suffices to consider the case where  $\mathcal{C} = \text{Vect}$ . The readers who prefer to think about  $\text{Vect}$  rather than a general category  $\mathcal{C}$  are welcome to do so without losing any content. In fact, the general story goes exactly the same for  $\text{Vect}$  as for a general  $\mathcal{C}$ .

4.2.6. *Another take on Rees and  $\text{Fil} \rightarrow \text{gr}$ .* The language of graded algebras and graded modules over a graded algebra allows us to have a convenient interpretation of the functors Rees and  $\text{Fil} \rightarrow \text{gr}$ . Indeed, we have the following well-known identification

$$(4.2.7) \quad \text{Mod}_{\Lambda[\mathbb{Z}_{\geq 0}^m]}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m, \rightarrow}) \simeq \mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m, \rightarrow},$$

where the multiplication by  $\Lambda[\mathbb{Z}_{\geq 0}^m]$  gives rise precisely to the maps in the filtration. Under this identification,  $\text{Fil} \rightarrow \text{gr}$  is given by forgetting the module structure, and Rees is given by tensoring up  $- \otimes \Lambda[\mathbb{Z}_{\geq 0}^m]$ .

4.2.8. *Another take on  $\text{ass-gr}_{\mathcal{C}, m}$ .* The description of  $\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m, \rightarrow}$  above also gives us a convenient way of interpreting  $\text{ass-gr}$ . Indeed, consider the following map of algebras in  $\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m}$

$$\Lambda[\mathbb{Z}_{\geq 0}^m] \rightarrow \mathbb{1}_{\mathcal{C}, 0}$$

by modding out the augmentation ideal. The functor  $\text{gr} \xrightarrow{0} \text{Fil}$  is given by restricting along this map, and hence, its left adjoint is given by tensoring up, i.e.

$$(4.2.9) \quad \text{ass-gr}_{\mathcal{C}, m} \simeq - \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \mathbb{1}_{\mathcal{C}}.$$

Note that here, we have implicitly used the identification

$$\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m} \simeq \text{Mod}_{\mathbb{1}_{\mathcal{C}, 0}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m})$$

since  $\mathbb{1}_{\mathcal{C}, 0}$  is the monoidal unit of  $\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m}$ .

4.2.10. The same discussion carries over for  $\mathcal{C}_{\geq 0, +}^{\mathbb{Z}_{\geq 0}^m}$  and  $\mathcal{C}_{\geq 0, +}^{\mathbb{Z}_{\geq 0}^m, \rightarrow}$ . Indeed, we still have

$$\text{Mod}_{\Lambda[\mathbb{Z}_{\geq 0}^m]}(\mathcal{C}_{\geq 0, +}^{\mathbb{Z}_{\geq 0}^m, \rightarrow}) \simeq \mathcal{C}_{\geq 0, +}^{\mathbb{Z}_{\geq 0}^m, \rightarrow}$$

which is a full-subcategory of  $\text{Mod}_{\Lambda[\mathbb{Z}_{\geq 0}^m]}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m})$ , and the category  $\mathcal{C}_{\geq 0, +}^{\mathbb{Z}_{\geq 0}^m}$  is a full-subcategory of  $\text{Mod}_{\mathbb{1}_{\mathcal{C}}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m})$ .

These two categories are preserved under  $\text{gr} \xrightarrow{0} \text{Fil}$  and  $\text{ass-gr}_{\mathcal{C}, m'}$ , and so, we have the same formula in this case as in (4.2.9). In particular, it is important to note that we still tensor over  $\Lambda[\mathbb{Z}_{\geq 0}^m]$ .

4.3. **Graded-unital and graded-augmented.** In this paper, we usual start with a graded object. But to be able to talk about homological stability, we need to have maps between the various graded-degrees, i.e. we need a filtered object. The equivalence (4.2.7) gives us the following convenient way of doing so.

4.3.1. Suppose we have a morphism in  $\text{ComAlg}^{\text{un}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m})$

$$\Lambda[\mathbb{Z}_{\geq 0}^m] \rightarrow \mathcal{A}.$$

Then  $\mathcal{A}$  acquires the structure of a  $\Lambda[\mathbb{Z}_{\geq 0}^m]$ -module and hence, the equivalence (4.2.7) implies that  $\mathcal{A}$  comes from an object in  $\text{ComAlg}^{\text{un}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m, \rightarrow})$ . We call such an object *graded-unital*, and let  $\text{ComAlg}^{\text{un}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m})$  denote the category of all such objects. Since  $\Lambda[\mathbb{Z}_{\geq 0}^m]$  is the monoidal unit of  $\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m, \rightarrow}$ , we have an equivalence

$$\text{ComAlg}^{\text{un}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m}) \simeq \text{ComAlg}^{\text{un}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m, \rightarrow}).$$

4.3.2. Similarly, we will use  $\text{ComAlg}^{\text{un, au}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m})$  to denote the category of *graded-unital augmented* algebras, where these objects are defined in the obvious way. As above, we have the following equivalence

$$\text{ComAlg}^{\text{un, au}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m}) \simeq \text{ComAlg}^{\text{un, au}}(\mathcal{C}_{\geq 0}^{\mathbb{Z}_{\geq 0}^m, \rightarrow}).$$

4.3.3. We have a natural forgetful functor

$$\mathrm{ComAlg}^{\underline{\mathrm{un}}, \underline{\mathrm{au}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}) \rightarrow \mathrm{ComAlg}^{\underline{\mathrm{un}}, \underline{\mathrm{au}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}).$$

We can thus define the category  $\mathrm{ComAlg}^{\underline{\mathrm{un}}, \underline{\mathrm{au}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m})$  to be the full-subcategory of  $\mathrm{ComAlg}^{\underline{\mathrm{un}}, \underline{\mathrm{au}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$  spanned by objects whose augmentation ideals live in  $\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m}$ . Namely, we have the following pullback square of categories

$$\begin{array}{ccc} \mathrm{ComAlg}^{\underline{\mathrm{un}}, \underline{\mathrm{au}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m}) & \longrightarrow & \mathrm{ComAlg}^{\underline{\mathrm{un}}, \underline{\mathrm{au}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}) \\ \downarrow & & \downarrow \\ \mathrm{ComAlg}^{\underline{\mathrm{un}}, \underline{\mathrm{au}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m}) & \longrightarrow & \mathrm{ComAlg}^{\underline{\mathrm{un}}, \underline{\mathrm{au}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}) \end{array}$$

4.3.4. Let  $\mathcal{A} \in \mathrm{ComAlg}^{\underline{\mathrm{un}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$ , one can “quotient out” the graded-unit by taking the tensor product, or equivalently, by taking the associated graded (see 4.2.8)

$$\mathrm{ass}\text{-}\mathrm{gr} \mathcal{A} \simeq \mathcal{A} \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \mathbb{1}_{\mathcal{C}} \in \mathrm{ComAlg}^{\underline{\mathrm{un}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$$

where  $\mathbb{1}_{\mathcal{C}}$  on the right sits in graded-degree  $(0, 0, \dots, 0)$ , i.e. it is the monoidal unit  $\mathbb{1}_{\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}}$ .

When we want to emphasize the perspective that we are taking the quotient by the unit rather than just taking the associated graded, we will use  $\mathrm{Quot}_{\underline{\mathrm{un}}}$  to denote this functor

$$\mathrm{Quot}_{\underline{\mathrm{un}}} : \mathrm{ComAlg}^{\underline{\mathrm{un}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}) \rightarrow \mathrm{ComAlg}^{\underline{\mathrm{un}}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}).$$

4.4. **Change of gradings.** One of the goals of the paper is to compare relative tensors of graded algebras. Strictly speaking, however, these algebras live in different categories, since they are graded by different gradings (i.e. different  $m$ ’s, in our notation).<sup>8</sup> We thus need a functorial way to tie them together.

4.4.1. We have the following monoidal functors

$$\mathrm{add} : \mathbb{Z}_{\geq 0}^m \rightarrow \mathbb{Z}_{\geq 0} \quad \text{and} \quad \mathrm{add}^{\rightarrow} : \mathbb{Z}_{\geq 0}^{m, \rightarrow} \rightarrow \mathbb{Z}_{\geq 0}^{\rightarrow}$$

obtained by adding all the components together. This gives rise to the following pairs of adjoint functors

$$\mathrm{add}_! : \mathcal{C}^{\mathbb{Z}_{\geq 0}^m} \rightleftarrows \mathcal{C}^{\mathbb{Z}_{\geq 0}} : \mathrm{add}^!$$

and

$$\mathrm{add}_!^{\rightarrow} : \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}} \rightleftarrows \mathcal{C}^{\mathbb{Z}_{\geq 0}^{\rightarrow}} : \mathrm{add}^{\rightarrow, !},$$

obtained by left Kan extending and restricting along  $\mathrm{add}/\mathrm{add}^{\rightarrow}$  respectively.

4.4.2. Since the monoidal structures on these graded and filtered categories are formed by Day’s convolutions, which is a left Kan extension, it is clear that  $\mathrm{add}_!$  and  $\mathrm{add}_!^{\rightarrow}$  are monoidal, which implies that  $\mathrm{add}^!$  and  $\mathrm{add}^{\rightarrow, !}$  are right-lax monoidal.

**Lemma 4.4.3.** *In the following diagram,*

$$\begin{array}{ccc} \mathcal{C}^{\mathbb{Z}_{\geq 0}^m} & \begin{array}{c} \xrightarrow{\mathrm{add}_!} \\ \xleftarrow{\mathrm{add}^!} \end{array} & \mathcal{C}^{\mathbb{Z}_{\geq 0}} \\ \mathrm{ass}\text{-}\mathrm{gr} \downarrow \mathrm{gr}^0 \mathrm{Fil} & & \mathrm{ass}\text{-}\mathrm{gr} \downarrow \mathrm{gr}^0 \mathrm{Fil} \\ \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}} & \begin{array}{c} \xrightarrow{\mathrm{add}_!^{\rightarrow}} \\ \xleftarrow{\mathrm{add}^{\rightarrow, !}} \end{array} & \mathcal{C}^{\mathbb{Z}_{\geq 0}^{\rightarrow}} \end{array}$$

<sup>8</sup>The algebras we are talking about are the  $\mathcal{A}_{m,n}(X)$  in the introduction.

the right adjoints commute, and hence, so do the left adjoints. In particular,

$$\mathrm{add}_! \circ \mathrm{ass}\text{-}\mathrm{gr} \simeq \mathrm{ass}\text{-}\mathrm{gr} \circ \mathrm{add}_!^{\rightarrow}.$$

4.4.4. The functor  $\mathrm{add}_!^{\rightarrow}$  has the following convenient interpretation in a way that is similar to the case of  $\mathrm{ass}\text{-}\mathrm{gr}$  considered above. Indeed, consider the following diagram

$$\begin{array}{ccccc} \mathrm{Mod}_{\Lambda[\mathbb{Z}_{\geq 0}^m]}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}) & \xrightleftharpoons[\mathrm{add}_!]{\mathrm{add}_!} & \mathrm{Mod}_{\Lambda[\mathbb{Z}_{\geq 0}^m]}(\mathcal{C}^{\mathbb{Z}_{\geq 0}}) & \xrightleftharpoons[\mathrm{res}_{\Lambda[\mathbb{Z}_{\geq 0}^m] \rightarrow \Lambda[\mathbb{Z}_{\geq 0}]}]{-\otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \Lambda[\mathbb{Z}_{\geq 0}]} & \mathrm{Mod}_{\Lambda[\mathbb{Z}_{\geq 0}]}(\mathcal{C}^{\mathbb{Z}_{\geq 0}}) \\ \parallel & & & & \parallel \\ \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}} & \xrightleftharpoons[\mathrm{add}^{\rightarrow, !}]{\mathrm{add}_!^{\rightarrow}} & & & \mathcal{C}^{\mathbb{Z}_{\geq 0}^{\rightarrow}} \end{array}$$

where, in the second term on the top, by  $\Lambda[\mathbb{Z}_{\geq 0}^m]$ , we mean  $\mathrm{add}_! \Lambda[\mathbb{Z}_{\geq 0}^m]$ , but this is only a mild abuse of notation since the two are really “the same” ring: only the gradings differ. Moreover, the adjunction pair  $\mathrm{add}_! \dashv \mathrm{add}_!^{\rightarrow}$  upgrades to one between module categories because  $\mathrm{add}_!$  is monoidal and hence,  $\mathrm{add}_!^{\rightarrow}$  is right-lax monoidal. Finally, we see easily that the right adjoints commute, i.e.

$$\mathrm{add}^{\rightarrow, !} \simeq \mathrm{add}_!^{\rightarrow} \circ \mathrm{res}$$

and hence, so do the left adjoints. Thus, we get the following convenient interpretation of  $\mathrm{add}_!^{\rightarrow}$

**Lemma 4.4.5.** *For each  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$ , we have a natural equivalence*

$$\mathrm{add}_!^{\rightarrow} V \simeq (\mathrm{add}_! V) \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \Lambda[\mathbb{Z}_{\geq 0}].$$

*Remark 4.4.6.* This lemma, combined with §4.2.8, gives us an alternative proof of Lemma 4.4.3. Namely,

$$\mathrm{ass}\text{-}\mathrm{gr}(\mathrm{add}_!^{\rightarrow}(V)) \simeq \Lambda \otimes_{\Lambda[\mathbb{Z}_{\geq 0}]} \Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} V \simeq \mathrm{add}_!(\Lambda \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} V).$$

4.5. **Stabilization.** We will now come to the concept of stabilization. In the simplest case, where  $m = 1$ , given an object

$$V = (\cdots \rightarrow V_1 \rightarrow V_2 \rightarrow \cdots) \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{\rightarrow}},$$

where the maps are not necessarily equivalences, we want to construct an object  $\bar{V} \in \mathcal{C}$ , which, in some sense, captures the stable part of the sequence  $V$ . From the categorical point of view, the obvious candidate is to take

$$\bar{V} = \mathrm{colim}_{d \in \mathbb{Z}_{\geq 0}^{\rightarrow}} V_d,$$

which happily turns out to capture precisely the stable homology of  $V$ , even though this construction makes sense in general. In this subsection, we will discuss various formal properties of this construction. The link to homological stability will be discussed in the next subsection.

4.5.1. We have the following pair of adjoint functors

$$\mathrm{colim}_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^{m, \rightarrow}} : \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}} \rightleftarrows \mathcal{C} : \mathrm{const}_{\mathbb{Z}_{\geq 0}^{m, \rightarrow}},$$

where the left adjoint is taking colimit and the right adjoint is the constant functor, i.e. sending  $V$  to the constant filtered object  $V$ .

**Notation 4.5.2.** For any  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$ , we will use

$$\bar{V} = \mathrm{colim}_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^{m, \rightarrow}} V_{\mathbf{d}}$$

to denote the colimit. Moreover, unless confusion might occur, we will write  $\mathrm{const}$  instead of  $\mathrm{const}_{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$ .

A couple of remarks are in order.



*Remark 4.5.3.* Since the diagonal embedding

$$\mathbb{Z}_{\geq 0}^{\rightarrow} \rightarrow \mathbb{Z}_{\geq 0}^{m, \rightarrow}$$

is co-final, one reduces the computation of  $\overline{V}$  to that of a sequential colimit.

*Remark 4.5.4.* Using the same reasoning, we see that  $\overline{V}$  only depends the behavior of  $V$  on  $\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^{m, \rightarrow}}$ . Because of this, we will also use the notation  $\overline{V}$  even for the case where  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0, +}^{m, \rightarrow}}$ .

4.5.5. We have the following lemma, which is direct from the fact that colimits commute with colimits and that the definition of the monoidal structures on  $\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$  and  $\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^{m, \rightarrow}}$  is defined using colimits.

**Lemma 4.5.6.** *The functor*

$$\overline{(-)} : \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}} \rightarrow \mathcal{C}$$

*is symmetric monoidal. As a consequence, its right adjoint  $\text{const}$  is right-lax monoidal.*

**Corollary 4.5.7.** *We have the following pairs of adjoint functors*

$$\begin{aligned} \overline{(-)} : \text{ComAlg}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}) &\rightleftarrows \text{ComAlg}(\mathcal{C}) : \text{const}, \\ \overline{(-)} : \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}) &\rightleftarrows \text{ComAlg}^{\text{un}}(\mathcal{C}) : \text{const}, \\ \overline{(-)} : \text{ComAlg}^{\text{au}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}) &\rightleftarrows \text{ComAlg}^{\text{au}}(\mathcal{C}) : \text{const}, \\ \overline{(-)} : \text{ComAlg}^{\text{un, au}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}) &\rightleftarrows \text{ComAlg}^{\text{un, au}}(\mathcal{C}) : \text{const}. \end{aligned}$$

*Remark 4.5.8.* Let  $\mathcal{A}$  be an algebra object in any of the categories appearing on the left side of Corollary 4.5.7, the algebra  $\overline{\mathcal{A}}$  should be thought of as the stable part of  $\mathcal{A}$ . Roughly speaking, this is because when  $\mathcal{A}$  exhibits a certain homological stability property, then  $\overline{\mathcal{A}}$  captures precisely this stable homology. This is the content of §4.6.

4.5.9. Since the functor  $\overline{(-)}$  is defined as a colimit, it behaves nicely with respect to the functor  $\text{add}_!^{\rightarrow}$  considered in §4.4, and we have the following

**Lemma 4.5.10.** *We have the following commutative diagram*

$$\begin{array}{ccc} \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}} & \xrightarrow{\text{add}_!^{\rightarrow}} & \mathcal{C}^{\mathbb{Z}_{\geq 0}^{\rightarrow}} \\ & \searrow \overline{(-)} & \downarrow \overline{(-)} \\ & & \mathcal{C} \end{array}$$

4.5.11. From the formal point of view, the functor  $\overline{(-)}$  above has an analog in the graded case, which we now record for later use. Namely, we have the following pair of adjoint functors

$$\text{oblv}_{\text{gr}} : \mathcal{C}^{\mathbb{Z}_{\geq 0}^m} \rightleftarrows \mathcal{C} : \text{const}_{\mathbb{Z}_{\geq 0}^m}$$

where we see easily that  $\text{oblv}_{\text{gr}}$  is the functor of taking direct sum. As above, we will write  $\text{const}$  instead of  $\text{const}_{\mathbb{Z}_{\geq 0}^m}$  unless confusion is likely to occur.

By the same token as for the functor  $\overline{(-)}$ ,  $\text{oblv}_{\text{gr}}$  commutes with colimits, and is symmetric monoidal. And thus, the adjoint pair  $\text{oblv}_{\text{gr}} \dashv \text{const}$  upgrades to adjoint pairs between various categories of commutative algebras over  $\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}$  and  $\mathcal{C}$  as in Corollary 4.5.7.

4.5.12. *Telescope construction.* Let  $\mathcal{A} \in \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$ , i.e. a unital graded commutative algebra equipped with the structure of a  $\Lambda[\mathbb{Z}_{\geq 0}^m]$ -algebra. Then, by §4.3,  $\mathcal{A}$  can be viewed as an object in  $\text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}})$ , and hence, we can talk about its stabilization  $\bar{\mathcal{A}} \in \text{ComAlg}^{\text{un}}(\mathcal{C})$ . Similarly, if  $\mathcal{A} \in \text{ComAlg}^{\text{un, au}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$ , we get  $\bar{\mathcal{A}} \in \text{ComAlg}^{\text{un, au}}(\mathcal{C})$ .

*Remark 4.5.13.* At the level of spaces, we learned this construction from [MS76]. In their setting, the authors want to find a model for the group completion of an  $E_n$ -algebra (naturally graded by the monoid of connected components). Roughly speaking, by choosing a special point on a connected component and using it to act on the original  $E_n$ -algebra, we can form a sequential diagram. The group completion is then computed as the homotopy colimit of this diagram, which has an explicit model as the telescope construction.

The passage from a graded space to a filtered one using the action of a point is similar to our passage from  $\text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$  to  $\text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}})$  using the action of  $\Lambda[\mathbb{Z}_{\geq 0}^m]$ . Moreover, in both cases, the final object is constructed as the homotopy colimit of the resulting diagram. This is the reason why we call this the telescope construction.

4.6. **Relation to homological stability.** We will now turn to the link between the stabilization construction via colimit above and stable homology. The main result of this subsection, Proposition 4.6.3, states that the stabilization  $\bar{V}$  of an object  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$  captures the stable homology of  $V$  when  $V$  has homological stability.

To start, we need to make sense of what homological stability means. For that, we assume that  $\mathcal{C}$  is equipped with a  $t$ -structure such that taking infinite direct sums in  $\mathcal{C}$  is  $t$ -exact. The word cohomology will be with respect to this  $t$ -structure.<sup>9</sup>

**Definition 4.6.1** (Homological stability). Let  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$ .

We say that  $V$  satisfies weak homological stability if for any integer  $c$ , there exists an integer  $D$  such that the map

$$\tau_{\leq c} V_{(d, d, \dots, d)} \rightarrow \tau_{\leq c} V_{(d+1, d+1, \dots, d+1)}$$

is an equivalence for all  $d \geq D$ .

We say that  $V$  satisfies (strong) homological stability if for any integer  $c$ , there exists an integer  $D$  such that the maps

$$\tau_{\leq c} V_{(d_1, d_2, \dots, d_k, \dots, d_m)} \rightarrow \tau_{\leq c} V_{(d_1, d_2, \dots, d_k+1, \dots, d_m)}$$

are equivalences for all  $1 \leq k \leq m$  and for all  $(d_1, \dots, d_m)$  with  $d_k \geq D$ .

It is straightforward to see that (strong) homological stability implies weak homological stability.

**Definition 4.6.2.** Let  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$  satisfying weak homological stability. Then for each integer  $c$ , we use  $\tau_{\leq c}^{\text{stab}} V$  to denote the stable homology of  $V$  in cohomological degrees  $\leq c$ . More explicitly,

$$\tau_{\leq c}^{\text{stab}} V = \tau_{\leq c} V_{(d, \dots, d)}$$

when  $d \gg 0$ .

We now comes the main result of this subsection.

**Proposition 4.6.3.** Let  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$  satisfying weak homological stability. Then, for any integer  $c$ , the natural map

$$\tau_{\leq c}^{\text{stab}} V \rightarrow \tau_{\leq c} \bar{V}$$

is an equivalence.

<sup>9</sup>As before, for the applications of this paper, it suffices to consider the case where  $\mathcal{C} = \text{Vect}$ . We choose to state the results in more generality since the proof is the same for  $\text{Vect}$  as for the more general case. The readers who prefer to think about  $\text{Vect}$  will not miss anything.

*Proof.* Because of Remark 4.5.3, we can reduce to the case where  $m = 1$ . By throwing away elements at the beginning of the sequence if necessary, we can further assume that for all  $d \in \mathbb{Z}_{\geq 0}^{\rightarrow}$ , the map

$$\tau_{\leq c+1} V_d \rightarrow \tau_{\leq c+1} V_{d+1}$$

is an equivalence. The desired conclusion comes from the following lemma.  $\square$

**Lemma 4.6.4.** *Let  $V \in \mathcal{C}_{\geq 0}^{\mathbb{Z}^{\rightarrow}}$  and  $c$  an integer such that for all  $d \in \mathbb{Z}_{\geq 0}^{\rightarrow}$ , the natural map*

$$\tau_{\leq c+1} V_d \rightarrow \tau_{\leq c+1} V_{d+1}$$

*is an equivalence. Then, the natural map*

$$\tau_{\leq c} V_d \rightarrow \tau_{\leq c} \bar{V}$$

*is an equivalence for all  $d$ . Equivalently, the natural map*

$$\tau_{\leq c}^{\text{stab}} V \rightarrow \tau_{\leq c} \bar{V}$$

*is an equivalence.*

*Proof.* Taking the colimit of the following fiber sequence

$$\tau_{\leq c+1} V_d \rightarrow V_d \rightarrow \tau_{> c+1} V_d \rightarrow \cdots,$$

we get the following fiber sequence

$$\tau_{\leq c+1}^{\text{stab}} V \rightarrow \bar{V} \rightarrow \text{colim}_d (\tau_{> c+1} V_d) \rightarrow \cdots.$$

Now, it suffices to show that

$$\text{colim}_d (\tau_{> c+1} V_d)$$

lives in cohomological degrees  $\geq c+1$ . The fact that this colimit fits into the following cofiber sequence

$$\bigoplus_d \tau_{> c+1} V_d \rightarrow \bigoplus_d \tau_{> c+1} V_d \rightarrow \text{colim}_d (\tau_{> c+1} V_d) \rightarrow \cdots$$

gives us the desired conclusion.  $\square$

*Remark 4.6.5.* Note that when  $V \in \mathcal{C}_{\geq 0}^{\mathbb{Z}^{m, \rightarrow}}$  satisfying weak homological stability, then the  $\tau_{\leq c}^{\text{stab}} V$ 's naturally map into each other and form a tower

$$\cdots \rightarrow \tau_{\leq c-1}^{\text{stab}} V \rightarrow \tau_{\leq c}^{\text{stab}} V \rightarrow \tau_{\leq c+1}^{\text{stab}} V \rightarrow \cdots$$

The discussion above implies that if we let

$$V^{\text{stab}} = \text{colim}_c \tau_{\leq c}^{\text{stab}} V,$$

then for any  $c$ ,

$$\tau_{\leq c}^{\text{stab}} V \simeq \tau_{\leq c} V^{\text{stab}}$$

and moreover, the natural map

$$V^{\text{stab}} \rightarrow \bar{V}$$

is an equivalence.

Being independent from homological stability, however, the general construction of  $\bar{V}$  is more robust, and is generally easier to manipulate. For example, when  $V$  is an algebra object,  $\bar{V}$  canonically inherits this algebra structure (see Corollary 4.5.7). We will see more instances of this phenomenon later.

4.6.6. Using the result above and Corollary 4.5.7, we see that if we start with a graded-unital algebra  $\mathcal{A}$  satisfying weak homological stability. Then, then the “stable homology”  $\bar{\mathcal{A}} \simeq \mathcal{A}^{\text{stab}}$  carries a natural unital algebra structure.

*Remark 4.6.7.* Even if an object  $V \in \mathcal{C}_{\geq 0}^{\mathbb{Z}^{m, \rightarrow}}$  exhibits no homological stability phenomenon, by abuse of terminology, we will still refer to the object  $\bar{V}$  as the stable (co)homology of  $V$ .

**4.7. Homological stability of a limit.** In this subsection, we record a simple lemma which provides a method to study the homological stability of a filtered-object that is itself a sequential limit. It allows us to reduce the study of homological stability of the original object to that of the successive fibers of the terms in the limit.

**Lemma 4.7.1.** *Let  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$  such that*

$$V \simeq \lim(\cdots \leftarrow V_{-1} \leftarrow V_0 \leftarrow V_1 \leftarrow V_2 \leftarrow \cdots)$$

*where  $V_1, V_2, \dots \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$ , and where for each  $\mathbf{d}$ ,  $V_{i, \mathbf{d}} \simeq 0$  when  $i \ll 0$ . Let*

$$F_i = \text{Fib}(V_i \rightarrow V_{i-1}) \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}.$$

*Let  $c \in \mathbb{Z}$ ,  $1 \leq k \leq m$ , and  $\mathbf{d} \in \mathbb{Z}_{\geq 0}^m$  be such that the following map is an equivalence*

$$(4.7.2) \quad \tau_{\leq c+1} F_{i, \mathbf{d}} \rightarrow \tau_{\leq c+1} F_{i, \mathbf{d} + \mathbf{1}_k}, \forall i.$$

*Then, the same is true for*

$$(4.7.3) \quad \tau_{\leq c} V_{\mathbf{d}} \rightarrow \tau_{\leq c} V_{\mathbf{d} + \mathbf{1}_k}.$$

*Proof.* Fix  $\mathbf{d} \geq \mathbf{D}$ . It suffices to show that for each  $i$ , the following map is an equivalence

$$\tau_{\leq c} V_{i, \mathbf{d}} \rightarrow \tau_{\leq c} V_{i, \mathbf{d} + \mathbf{1}_k}.$$

Indeed, this is because the functor  $\text{tr}_{\leq c}$ , being a right adjoint, commutes with limits. But now, it is a simple induction on  $i$ . Note that the condition that for each  $\mathbf{d}$ ,  $V_{i, \mathbf{d}} \simeq 0$  when  $i \ll 0$  provides us with the base case.  $\square$

**4.7.4.** In practice, the maps (4.7.2), and hence also (4.7.3), are equivalences for a range  $\mathbf{d} \geq \mathbf{D}$  for some  $\mathbf{D}$ . This Lemma is thus really about homological stability of a sequential limit as we mentioned above. The bound is where homological stability, with respect to homological degrees  $\leq c$ , occurs.

In other words, for the applications that we will encounter, the Lemma above says that if  $V \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^m}$  is a sequential limit, whose associated graded satisfies homological stability, then so does  $V$ . Moreover, the homological degree at which we have homological stability (i.e. the number  $c$  in the above lemma) for  $V$  is only 1 worse than that of the associated graded (i.e.  $c+1$  vs  $c$ ).

We have the following variant of the Lemma above

**Lemma 4.7.5.** *In the situation of Lemma 4.7.1, suppose further that*

$${}^t\text{H}^{c+2}(F_{i, \mathbf{d}}) \rightarrow {}^t\text{H}^{c+2}(F_{i, \mathbf{d}})$$

*is injective. Then we can improve the homological degree at which stability occurs. Namely, the following map is an equivalence*

$$\tau_{\leq c+1} V_{\mathbf{d}} \rightarrow \tau_{\leq c+1} V_{\mathbf{d} + \mathbf{1}_k}.$$

*Proof.* The proof goes the same way as in Lemma 4.7.1. The improvement comes from using the four-lemma for injective maps.  $\square$

**4.7.6.** In the case where taking infinite products in  $\mathcal{C}$  is  $t$ -exact, such as the case of  $\text{Vect}$ , we can push it one step further.

**Lemma 4.7.7.** *When taking infinite products in  $\mathcal{C}$  is  $t$ -exact, in the situation of Lemma 4.7.5, the following map is an equivalence (as in Lemma 4.7.5)*

$$\tau_{\leq c+1} V_{\mathbf{d}} \rightarrow \tau_{\leq c+1} V_{\mathbf{d} + \mathbf{1}_k}$$

*and moreover, the map*

$${}^t\text{H}^{c+2}(V_{\mathbf{d}}) \rightarrow {}^t\text{H}^{c+2}(V_{\mathbf{d} + \mathbf{1}_k})$$

*is injective.*

*Proof.* By induction, and repeated application of the four-lemma for injective morphisms, we get that the following map is an equivalence

$$\tau_{\leq c+1} V_{i,\mathbf{d}} \rightarrow \tau_{\leq c+1} V_{i,\mathbf{d}+\mathbf{1}_k}$$

and that

$${}^t\mathrm{H}^{c+2}(V_{i,\mathbf{d}}) \rightarrow {}^t\mathrm{H}^{c+2}(V_{i,\mathbf{d}+\mathbf{1}_k})$$

is injectively. Now, it suffices to show that taking sequential limit preserves injectivity at cohomological degree  $c+2$ .

Indeed, consider the following diagram

$$\begin{array}{ccccc} \lim_i V_{i,\mathbf{d}} & \longrightarrow & \prod V_{i,\mathbf{d}} & \longrightarrow & \prod V_{i-1,\mathbf{d}} \\ \downarrow & & \downarrow & & \downarrow \\ \lim_i V_{i,\mathbf{d}+\mathbf{1}_k} & \longrightarrow & \prod V_{i,\mathbf{d}+\mathbf{1}_k} & \longrightarrow & \prod V_{i-1,\mathbf{d}+\mathbf{1}_k} \end{array}$$

where the rows are exact sequences. Now, taking cohomology and applying four-lemma, we get the desired conclusion.  $\square$

*Remark 4.7.8.* As we will see later, the conditions of Lemma 4.7.7 are always satisfied in cases that we are interested in. The injectivity appearing in this Lemma is the ultimate reason why we have an extra surjectivity at the edge of homological stability first observed in [Knu14]. The difference injectivity (here) vs. surjectivity (there) is due to the fact that the two pictures are linear dual to each other: homology vs. cohomology.

**4.8. Stable homology for objects of the form  $W \otimes \Lambda[\mathbb{Z}_{\geq 0}^m]$ .** In this subsection, we give a description for  $\bar{V}$  when  $V$  is of the form  $W \otimes \Lambda[\mathbb{Z}_{\geq 0}^m]$ . In this case  $\bar{V}$  has a particularly simple description.

**Proposition 4.8.1.** *Let  $W \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^m}$  and  $V = W \otimes \Lambda[\mathbb{Z}_{\geq 0}^m] \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m,\rightarrow}}$ . Then, we have a natural equivalence*

$$\bar{V} \simeq \mathrm{oblv}_{\mathrm{gr}}(W) \simeq \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} W_{\mathbf{d}}.$$

*Proof.* Consider the following diagram

$$(4.8.2) \quad \begin{array}{ccc} \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m,\rightarrow}} & \xleftarrow[\mathrm{const}]{\overline{(-)}} & \mathcal{C} \\ \mathrm{Rees} \downarrow & \mathrm{oblv}_{\mathrm{gr}} \nearrow & \uparrow \mathrm{const} \\ \mathcal{C}^{\mathbb{Z}_{\geq 0}^m} & \mathrm{Fil} \rightarrow \mathrm{gr} & \end{array}$$

where parallel arrows are adjoint pairs. Moreover, the right adjoints commute, and hence, so do the left adjoints.

Recall that by §4.2.6, we have

$$(4.8.3) \quad \mathrm{Rees} \simeq - \otimes \Lambda[\mathbb{Z}_{\geq 0}^m].$$

But now, the commutativity of the left adjoints in the diagram above gives us the desired conclusion

$$\bar{V} \simeq \overline{W \otimes [\mathbb{Z}_{\geq 0}^m]} \simeq \overline{\mathrm{Rees}(W)} \simeq \mathrm{oblv}_{\mathrm{gr}} W \simeq \bigoplus_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} W_{\mathbf{d}}.$$

$\square$

*Remark 4.8.4.* It is easy to see that the same result holds for  $\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}$ . Note, however, that  $V$  is still of the form  $W \otimes \Lambda[\mathbb{Z}_{\geq 0}^m]$  rather than  $W \otimes \Lambda[\mathbb{Z}_{\geq 0,+}^m]$ . The only difference is that  $W$  is now in  $\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}$  instead, which is a property, rather than an extra piece of data. See also §4.2.8 for a similar discussion.

*Remark 4.8.5.* Note that the object  $V = W \otimes \Lambda[\mathbb{Z}_{\geq 0}^m] \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^{m,\rightarrow}}$  where  $W \in \mathcal{C}^{\mathbb{Z}_{\geq 0}^m}$  does not necessarily have homological stability. However, as in Remark 4.6.7, we still call  $\bar{V} \simeq W$  the stable homology of  $V$ .

**Corollary 4.8.6.** *Let  $\mathcal{A} = \mathcal{B} \otimes \Lambda[\mathbb{Z}_{\geq 0}^m] \in \text{ComAlg}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m,\rightarrow}})$  where  $\mathcal{B} \in \text{ComAlg}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$ . Then, we have a natural equivalence of algebras in  $\mathcal{C}$*

$$\bar{\mathcal{A}} \simeq \mathcal{B} \in \text{ComAlg}(\mathcal{C}).$$

*Remark 4.8.7.* Note that if  $\mathcal{B} \in \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$ , then  $\mathcal{A} = \mathcal{B} \otimes \Lambda[\mathbb{Z}_{\geq 0}^m] \in \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^{m,\rightarrow}})$  (and similarly for  $\text{ComAlg}^{\text{au}}$  and  $\text{ComAlg}^{\text{un,au}}$ ).  $\bar{\mathcal{A}}$  then also carries the relevant unital/augmented structure, which is compatible with that of  $\mathcal{B}$  via the equivalence in Corollary 4.8.6 above.

## 5. COHOMOLOGICAL CHEVALLEY COMPLEX

Using the general framework set up above, we will now study various aspects of the cohomological Chevalley complex associated to a coLie-coalgebra. We will start with a homological stability result in §5.1, whose proof, as indicated earlier, reduces to the case of trivial (aka. abelian) coLie-coalgebras. In §5.2, we will provide a mean to study the limiting cohomology when we have homological stability. Finally, we will, in §5.3, indicate how various decategorification processes could be used to extract information from the stable cohomology.

**5.1. Homological stability.** We will now prove homological stability for the cohomological Chevalley complexes of a large class of coLie-coalgebras. The actual work needed at this stage is in fact small: the results here are essentially formal consequences of the machinery developed above.

To start, we want to put ourselves in the situation of §4.5.12. Namely, we want a condition on the coLie-coalgebra  $\mathfrak{a} \in \text{coLie}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m})$  such that

$$\text{coChev } \mathfrak{a} \in \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}),$$

or equivalently

$$\text{coChev}^{\text{un}} \mathfrak{a} \in \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0}^m})$$

which automatically upgrades to an object in

$$\text{coChev}^{\text{un}} \mathfrak{a} \in \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^{m,\rightarrow}}),$$

which, in turn, allows us to talk about homological stability. Here, as before,  $\mathcal{C}$  is a symmetric monoidal stable  $\infty$ -category.

Via Koszul duality between  $\text{ComAlg}$  and  $\text{coLie}$ , the desired condition is provided in the following

**Definition 5.1.1.** A coLie-coalgebra  $\mathfrak{a} \in \text{coLie}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m})$  is said to be unital if it is equipped with a coLie-coalgebra map

$$(5.1.2) \quad \bigoplus_{k=1}^m \mathbb{1}_{\mathcal{C}}[1]_{1_k} \rightarrow \mathfrak{a},$$

where the object on the LHS is equipped with the trivial coLie-structure.<sup>10</sup>

We will use  $\text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m})$  to denote the category of unital coLie-coalgebras.

In what follows, to simplify the notation, we will write  $\mathbb{1}_{\mathcal{C}}[1]^{\oplus m}$  to denote the coLie-coalgebra on the LHS of (5.1.2). For example, when  $\mathcal{C} = \text{Vect}$ , we will write  $\Lambda[1]^{\oplus m}$ . More generally, we will drop the gradings from the notation, unless it is not clear from the context.

<sup>10</sup>Recall that the subscript  $1_k$  in  $\mathbb{1}_{\mathcal{C}}[1]_{1_k}$  says that  $\mathbb{1}_{\mathcal{C}}[1]$  is placed in graded degree  $1_k$ . See also §4.1.2.

5.1.3. Since  $\text{oblv}_{\text{coLie}}$  is a left adjoint, it commutes with colimits. In other words, at the level of the underlying objects, colimits of  $\text{coLie}$ -coalgebras could be computed inside the underlying category.

Let

$$\text{Quot}_{\text{un}} : \text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}) \rightarrow \text{coLie}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m})$$

denote the functor of removing (or more accurately, quotient-ing out) the unit of the  $\text{coLie}$ -coalgebra by taking the cofiber of the map (5.1.2).

5.1.4. By Koszul duality, we have the following equivalence of categories

$$\text{coPrim}[1] : \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}) \simeq \text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}) : \text{coChev}.$$

Indeed, this is because

$$\text{coChev}(\mathbb{1}_{\mathcal{C}}^{\oplus m}[1]) \simeq \text{Sym}(\mathbb{1}_{\mathcal{C}}^{\oplus m})_+ \simeq \Lambda[\mathbb{Z}_{\geq 0,+}^m].$$

Moreover, the following diagram commutes since  $\text{coChev}$  is an equivalence of categories, and hence, in particular, preserves colimits.

$$\begin{array}{ccc} \text{ComAlg}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightleftharpoons[\text{coChev}]{} & \text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}) \\ \text{Quot}_{\text{un}} \downarrow & & \downarrow \text{Quot}_{\text{un}} \\ \text{ComAlg}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightleftharpoons[\text{coChev}]{} & \text{coLie}(\mathcal{C}^{\mathbb{Z}_{\geq 0,+}^m}) \end{array}$$

Here,  $\text{Quot}_{\text{un}}$  is discussed above, and  $\text{Quot}_{\text{un}}$  is discussed in §4.3.4.

Note that we have a similar commutative diagram for  $\text{coChev}^{\text{un}}$  (see also §2.3.3).

5.1.5. From now to the end of this subsection, to keep the notations and conditions simple, we will restrict ourselves to the case where  $\mathcal{C} = \text{Vect}$ .<sup>11</sup>

We are interested in the following special class of  $\text{coLie}$ -coalgebra.

**Definition 5.1.6.** An object  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m})$  is said to be *strongly unital* if it lives in cohomological degrees  $\geq -1$ , and moreover, the unit map induces an equivalence at cohomological degree  $-1$ , namely, we have an equivalence

$$\Lambda^{\oplus m} \simeq H^{-1}(\mathfrak{a})$$

i.e.  $\text{Quot}_{\text{un}}(\mathfrak{a})$  lives in cohomological degrees  $\geq 0$ .

As mentioned before, the proof of homological stability will be reduced to the case of trivial  $\text{coLie}$ -coalgebras. In this case, we record the following lemma which is immediate by inspection.

**Lemma 5.1.7.** Let  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m})$  be a strongly unital trivial  $\text{coLie}$ -coalgebra. Fix  $c_0 \in \mathbb{N}, 1 \leq k \leq m$  and suppose that there exist  $s, s_1, \dots, s_{k-1}, s_{k+1}, \dots, s_m \in \mathbb{N}$  such that for any  $v \in H^*(\text{Quot}_{\text{un}}(\mathfrak{a}))$  of cohomological degree  $c - 1 \leq c_0$  and graded degree  $\mathbf{d} \in \mathbb{Z}_{\geq 0,+}^m$ , we have

$$(5.1.8) \quad \mathbf{d}_k \leq sc + \sum_{i \neq k} s_i \mathbf{d}_i.$$

Then, for all  $c \leq c_0$  and  $\mathbf{d} \in \mathbb{Z}_{\geq 0,+}^m$  such that

$$\mathbf{d}_k \geq sc + \sum_{i \neq k} s_i \mathbf{d}_i,$$

the following map is an equivalence

$$\tau_{\leq c}(\text{coChev}^{\text{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow \tau_{\leq c}(\text{coChev}^{\text{un}} \mathfrak{a})_{\mathbf{d} + \mathbf{1}_k}.$$

<sup>11</sup>The readers are invited to formulate the general results for a general symmetric monoidal stable  $\infty$ -category  $\mathcal{C}$  equipped with a  $t$ -structure. Essentially, the conditions one has to spell out are about the interaction between the monoidal structure on  $\mathcal{C}$  and the given  $t$ -structure.



More generally, for any  $c$ , the following map is injective

$$H^c(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow H^c(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}+\mathbf{1}_k}.$$

*Proof (Sketch).* Note that the statement is really just a classical statement about homological stability of symmetric algebras, since the  $\mathrm{coLie}$ -coalgebra we are dealing with is abelian, which implies that

$$\mathrm{coChev} \mathfrak{a} \simeq \mathrm{Sym}(\mathfrak{a}[-1])_+.$$

For symmetric algebras, this can be seen easy by carefully examining which monomials can appear in which cohomological and graded-degrees.<sup>12</sup> The injectivity statement at the end is due to the fact that multiplication is injective in a symmetric algebra.  $\square$

*Example 5.1.9.* When  $\mathfrak{a}$  is such that  $\mathrm{coChev}^{\mathrm{un}} \mathfrak{a} = \Lambda[\mathbb{Z}_{\geq 0}^m]$ ,  $\mathrm{Quot}_{\mathrm{un}}(\mathfrak{a}) = 0$ , and hence, for any  $c_0$ , we can take the  $s$  and  $s_i$ 's to be 0. Lemma 5.1.7 then implies that we have homological stability for all homological degrees for all graded-degrees. This is, of course, an obvious statement by looking directly at  $\Lambda[\mathbb{Z}_{\geq 0}^m]_+$ .

*Example 5.1.10.* Suppose that  $\mathfrak{a}$  is a strongly unital trivial  $\mathrm{coLie}$ -coalgebra supported only at graded degrees  $\mathbf{1}_1, \dots, \mathbf{1}_m$ . Then, for any  $c_0$  and  $k$ , we can take  $s = 1, s_i = 0, \forall i \neq k$ . It follows that the map below is an equivalence

$$\tau_{\leq c}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow \tau_{\leq c}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}+\mathbf{1}_k}$$

whenever  $\mathbf{d}_k \geq c$ .

5.1.11. Recall that the result of §3.3 says that  $\mathrm{coChev}$  (and hence,  $\mathrm{coChev}^{\mathrm{un}}$ ) could be computed as a sequential limit, whose successive fibers form the  $\mathrm{coChev}^{\mathrm{un}}$  of the trivial  $\mathrm{coLie}$ -coalgebra with the same underlying object. Recall also that Lemma 4.7.7 says that homological stability of a sequential limit could be deduced from that of the successive fibers. Lemma 5.1.7 above establishes the case of trivial  $\mathrm{coLie}$ -coalgebra. And hence, combining all these results together, we obtain the following result.

**Theorem 5.1.12.** *Let  $\mathfrak{a} \in \mathrm{coLie}^{\mathrm{un}}(\mathrm{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  be a strongly unital  $\mathrm{coLie}$ -coalgebra. Fix  $c_0 \in \mathbb{N}$ ,  $1 \leq k \leq m$  and suppose that there exist  $s, s_1, \dots, s_{k-1}, s_{k+1}, \dots, s_m \in \mathbb{N}$  such that for any non-zero  $v \in H^*(\mathrm{Quot}_{\mathrm{un}}(\mathfrak{a}))$  of cohomological degree  $c - 1 \leq c_0$  and graded degree  $\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m$ , we have*

$$(5.1.13) \quad \mathbf{d}_k \leq sc + \sum_{i \neq k} s_i \mathbf{d}_i.$$

*Then, for all  $c \leq c_0$  and  $\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m$  such that*

$$\mathbf{d}_k \geq sc + \sum_{i \neq k} s_i \mathbf{d}_i,$$

*the following map is an equivalence*

$$\tau_{\leq c}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow \tau_{\leq c}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}+\mathbf{1}_k},$$

*and the following map is injective*

$$H^{c+1}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow H^{c+1}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}+\mathbf{1}_k}.$$

*Example 5.1.14.* Let  $\mathfrak{a} \in \mathrm{coLie}^{\mathrm{un}}(\mathrm{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  be a strongly unital  $\mathrm{coLie}$ -coalgebra supported at graded degrees  $\mathbf{1}_1, \dots, \mathbf{1}_m$ . Then, as in Example 5.1.10, for any  $c_0$  and  $k$ , we can take  $s = 1, s_i = 0, \forall i \neq k$ . It follows that whenever  $\mathbf{d}_k \geq c$  the map below is an equivalence

$$\tau_{\leq c}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow \tau_{\leq c}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}+\mathbf{1}_k}$$

and the map below is injective

$$H^{c+1}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}} \rightarrow H^{c+1}(\mathrm{coChev}^{\mathrm{un}} \mathfrak{a})_{\mathbf{d}+\mathbf{1}_k}.$$

<sup>12</sup>One thing to be careful about is the  $-1$  appearing in  $c - 1$ . This is there because we eventually shift  $\mathfrak{a}$  to the right by 1, in  $\mathrm{coChev} \mathfrak{a} \simeq \mathrm{Sym}(\mathfrak{a}[-1])_+$ .

We have the following easy Corollary of the Theorem above.

**Corollary 5.1.15.** *Let  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  be a strongly unital coLie-coalgebra such that  $H^*(\mathfrak{a})$  is finite dimensional. Then,  $\mathcal{A} = \text{coChev}^{\text{un}} \mathfrak{a}$  satisfies homological stability and moreover, the stable homology  $\overline{\mathcal{A}}$  is a commutative algebra with finite dimensional  $H^i(\overline{\mathcal{A}})$  is for each  $i$ .*

*Remark 5.1.16.* It is obvious how Theorem 5.1.12 allows us to extend beyond the finite dimensional case of Corollary 5.1.15. Indeed, there is no finite dimensionality restriction in the theorem. For the applications in this paper, however, the finite dimensional case suffices.

*Remark 5.1.17.* Suppose that our unital coLie-coalgebra  $\mathfrak{a}$  carries the action of some operator  $T$  (which will be the Frobenius in our application), then by functoriality,  $\text{coChev}^{\text{un}} \mathfrak{a}$  also carries this structure. Suppose the action of  $T$  on the unit of  $\mathfrak{a}$  is the identity, then we see that  $\text{coChev}^{\text{un}} \mathfrak{a}$ , as an object of  $\text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}})$ , carries the action of  $T$  as well. In other words, the action of  $T$  is compatible with maps in the filtration. As a result,  $T$  acts on the stabilization  $\overline{\text{coChev}^{\text{un}} \mathfrak{a}}$ . In particular, when homological stability does occur,  $T$  acts naturally on the stable homology.

In what follows, unless otherwise specified, whenever we have an action of an operator  $T$  on a coLie-coalgebra  $\mathfrak{a}$ , we will always assume that the action on the unit is the identity.

**5.2. Stable homology.** Let  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m})$  and  $\mathcal{A} = \text{coChev} \mathfrak{a}$ . The goal of this subsection is to understand  $\overline{\mathcal{A}}$  in terms of  $\mathfrak{a}$  itself. When we are in the situation where  $\overline{\mathcal{A}}$  satisfies homological stability, Lemma 4.6.4 says that the homology of  $\overline{\mathcal{A}}$  is the stable homology (see also Remark 4.6.5). It is thus crucial to understand  $\overline{\mathcal{A}}$ .

Note that in this section, we do not concern ourselves with questions about homological stability. Thus, we will work with a general symmetric monoidal stable  $\infty$ -category  $\mathcal{C}$ , since that does not bring in any extra complexity.

5.2.1. We start with the simplest case, where  $\mathfrak{a}$  is abelian, or more generally, when the unit of  $\mathfrak{a}$  splits.

**Proposition 5.2.2.** *Let  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m})$  such that the unit of  $\mathfrak{a}$  splits (as objects in  $\text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m})$ ), i.e. we have the following equivalence of coLie-coalgebras*

$$\mathfrak{a} \simeq \Lambda^{\oplus m}[1] \oplus \mathfrak{b}.$$

*Then, we have the following equivalence*

$$\overline{\text{coChev} \mathfrak{a}} \simeq \overline{\text{coChev}^{\text{un}} \mathfrak{a}} \simeq \text{oblv}_{\text{gr}}(\text{coChev}^{\text{un}} \mathfrak{b}) \in \text{ComAlg}^{\text{un}}(\mathcal{C})$$

*Proof.* Since  $\text{coChev} \mathfrak{a}$  and  $\text{coChev}^{\text{un}} \mathfrak{a}$  only differ at  $\mathbf{0}$ , their colimits over  $\mathbb{Z}_{\geq 0}^{m, \rightarrow}$  are the same, which gives the first equivalence.

For the second equivalence, note that for any coLie-coalgebras  $\mathfrak{u}, \mathfrak{v} \in \text{coLie}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m})$ , we have

$$\text{coChev}(\mathfrak{u} \oplus \mathfrak{v}) \simeq \text{coChev}(\mathfrak{u}) \sqcup \text{coChev}(\mathfrak{v}),$$

where the coproduct is taken inside  $\text{ComAlg}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m})$ . Indeed,  $\text{coChev}$  exhibits an equivalence of categories, and hence, preserves colimits. Thus, we have (see §2.3)

$$\text{coChev}^{\text{un}}(\mathfrak{u} \oplus \mathfrak{v}) \simeq \text{coChev}^{\text{un}}(\mathfrak{u}) \otimes \text{coChev}^{\text{un}}(\mathfrak{v}).$$

Applied to the case at hand, we have

$$\text{coChev}^{\text{un}}(\mathfrak{a}) \simeq \Lambda[\mathbb{Z}_{\geq 0}^m] \otimes \text{coChev}^{\text{un}}(\mathfrak{b}).$$

Corollary 4.8.6 then gives us the desired conclusion.  $\square$

5.2.3. In the case where the unit of  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m})$  does not split, we do not have a simple expression for the stable piece. However, we have the following result.

**Lemma 5.2.4.** *Let  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\mathcal{C}^{\mathbb{Z}_{\geq 0, +}^m})$ . Then, we have the following natural equivalence*

$$\text{ass-gr}_{\mathcal{C}, m}(\text{coChev}^{\text{un}}(\mathfrak{a})) \simeq \text{coChev}^{\text{un}}(\text{Quot}_{\text{un}} \mathfrak{a}).$$

*Proof.* The lemma follows directly from §4.2.8. Indeed, we have

$$(5.2.5) \quad \text{ass-gr}_{\mathcal{C}, m}(\text{coChev}^{\text{un}}(\mathfrak{a})) \simeq \text{coChev}^{\text{un}}(\mathfrak{a}) \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \mathbb{1}_{\mathcal{C}} \simeq \text{Quot}_{\text{un}}(\text{coChev}^{\text{un}}(\mathfrak{a})) \simeq \text{coChev}^{\text{un}}(\text{Quot}_{\text{un}} \mathfrak{a}),$$

where the first and last equivalences are due to §4.2.8 and §5.1.4 respectively. The middle equivalence is by definition.  $\square$

*Remark 5.2.6.* Note that the Lemma above is a special feature in the unital setting, i.e.  $\text{coChev}^{\text{un}}$  rather than  $\text{coChev}$ . Indeed, due to §4.2.8,

$$\text{ass-gr}_{\mathcal{C}, m}(\text{coChev} \mathfrak{a}) \simeq \text{coChev}(\mathfrak{a}) \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \mathbb{1}_{\mathcal{C}}.$$

However,

$$\text{Quot}_{\text{un}}(\text{coChev} \mathfrak{a}) \simeq (\text{coChev} \mathfrak{a}) \sqcup_{\Lambda[\mathbb{Z}_{\geq 0, +}^m]} 0 \simeq (\text{coChev}^{\text{un}} \mathfrak{a} \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \mathbb{1}_{\mathcal{C}})_+.$$

And these two do not agree, in fact, already in the case where  $m = 1$  and  $\mathfrak{a} = \Lambda[1]$  sitting in graded-degree 1. Of course, this does not cause us any problem, since we can always use  $\text{coChev}^{\text{un}}$  instead and since, as we mentioned earlier, as far as homological stability and stable homology are concerned, there is no difference.

**5.3. Decategorification.** In this section, we will provide various decategorification processes which can be used to extract information out of the functor  $\text{coChev}$ . More precisely, let  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  be a strongly unital  $\text{coLie}$ -coalgebra. Given a ring  $R$  and a decategorification procedure

$$\text{DeCat} : \mathcal{C} \rightarrow R$$

where  $\mathcal{C}$  is  $\text{Vect}$ ,  $\text{Vect}^{\mathbb{Z}_{\geq 0}^m}$  or  $\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m}$  (plus some appropriate finiteness conditions), we want to understand  $\text{DeCat}(\text{coChev}^{\text{un}} \mathfrak{a})$  and  $\overline{\text{DeCat}(\text{coChev}^{\text{un}} \mathfrak{a})}$  (when  $\mathfrak{a}$  satisfies some finiteness condition) in relation to various push-out operations. Moreover, we want to understand how  $\text{DeCat}$  of  $\text{coChev}^{\text{un}}(-)$  and of  $\text{coChev}^{\text{un}}(-)$  behave with respect to taking quotients of  $\text{coLie}$ -coalgebras. For the geometric applications in this paper, this part is used as a link between quotients of algebras and quotients (in the ring  $R$ ) of their decategorifications, i.e. what we called densities in the introduction.

5.3.1.  $\text{DeCat}$  is said to be additive, if for an exact triangle

$$V' \rightarrow V \rightarrow V'' \rightarrow V[1],$$

we have

$$\text{DeCat}(V) = \text{DeCat}(V') + \text{DeCat}(V'').$$

5.3.2. In this paper, instances of  $\text{DeCat}$  we are interested in are as follows

- (i) *Graded Euler characteristics.* Let  $V \in \text{Vect}^{\mathbb{Z}_{\geq 0}^m}$  such that each graded piece is perfect (i.e. bounded). Then, the graded Euler characteristics of  $V$  is defined to be

$$\chi^{\text{gr}}(V) = \sum_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} \chi(V_{\mathbf{d}}) t^{|\mathbf{d}|} \in \mathbb{Z}[[t]].$$

Here, the  $\chi$  on the RHS is the usual Euler characteristics of a perfect complex.

- (ii) *Graded traces.* More generally, we let  $V \in \text{Vect}^{\mathbb{Z}_{\geq 0}^m}$  as above, now, also equipped with an action of an operator  $T$ . Then, the graded trace of  $T$  on  $V$  is defined to be

$$\chi_T^{\text{gr}}(V) = \sum_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} \chi_T(V_{\mathbf{d}}) t^{|\mathbf{d}|} \in \mathbb{Z}[[t]].$$

As above, the  $\chi_T$  on the RHS is the usual trace. For the applications that we have in mind,  $V$  is equipped with the action of the Frobenius, and we are interested in computing the Frobenius trace.

Clearly, when  $T = \text{id}$ , we recover graded Euler characteristics above  $\chi_{\text{id}} = \chi$ .

- (iii) *Poincaré polynomials.* Let  $V \in \text{Vect}$  such that  $H^i(V)$  is finite dimensional for each  $i$ . Then, we define the Poincaré polynomial of  $V$  to be

$$\text{Poinc}(V) = \sum_{i \in \mathbb{Z}} \dim H^i(V) t^i \in \mathbb{N}[[t^{\pm 1}]].$$

- (iv) *Virtual Poincaré polynomials.* For the geometric applications we have in mind,  $V$  is equipped with the weight filtration  $W$  (from the action of the Frobenius). Let  $\text{Gr}_W^i(V)$  denote the sub-quotient of  $H^*(V)$  with weight  $i$ . We define the virtual Poincaré polynomial of  $V$  to be

$$\text{Poinc}^{\text{vir}}(V) = \sum_{i \in \mathbb{Z}} \chi(\text{Gr}_W^i(V)) t^i \in \mathbb{Z}[[t^{\pm 1}]].$$

- (v) *Trace.* Let  $V \in \text{Vect}$  equipped with an action of an operator  $T$  such that

- $H^c(V)$  is finite dimensional for each  $c$ .
- For each  $\lambda \in \mathbb{C}$  and every integer  $c$ , let  $d_{\lambda,c}$  denote the dimension of the generalized  $\lambda$ -eigenspace of  $T$ .<sup>13</sup> Then the sum

$$|V|_T = \sum_{\lambda, c} d_{\lambda, c} |\lambda|$$

converges.

Then the trace of  $T$  on  $V$  is defined in the usual way, i.e.

$$\chi_T(V) = \sum_{\lambda, c} (-1)^c d_{\lambda, c} \lambda.$$

We call such a graded vector space  $V$  satisfying the two conditions above  $T$ -summable.<sup>14</sup>

Note that in the above, everything except for Poincaré polynomials is additive. In this paper, the first two instances will be used to decategorify  $\text{coChev}^{\text{un}} \mathfrak{a}$  whereas the three two are for  $\text{coChev}^{\text{un}} \mathfrak{a}$ .

*Remark 5.3.3.* We will also use the notation,  $\text{DeCat}(-, t)$  to emphasize that the formal variable is  $t$ . For example,  $\chi^{\text{gr}}(V, t), \chi_T^{\text{gr}}(V, t), \text{Poinc}(V, t)$  etc.

5.3.4. As we mentioned in the introduction, in many cases, the process of taking quotients of  $\text{coLie}$ -coalgebras, which is equivalent to taking pushouts (i.e. relative tensor) of the corresponding commutative algebras via  $\text{coChev}^{\text{un}}$ , transforms to taking quotients (i.e. dividing, literally) of power series at the decategorified level. In other words, pushouts of commutative algebras categorify quotients of power series.

5.3.5. Observe that in all of the above, the expressions defining the decategorification processes make sense as soon as each of the coefficients make sense. Suppose we are interested in understanding  $\text{DeCat}(\text{coChev}^{\text{un}}(\mathfrak{a}))$  and  $\text{DeCat}(\text{coChev}^{\text{un}}(\bar{\mathfrak{a}}))$ , then we will need some finiteness conditions on  $\mathfrak{a}$ . For example, it is easy to see, using §3.3, that when  $\mathfrak{a} \in \text{coLie}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  is strongly unital, and  $H^*(\mathfrak{a})$  is finite dimensional then all the sums defining  $\chi^{\text{gr}}, \chi_T^{\text{gr}}$ , and  $\text{Poinc}$  are well-defined. In the case of  $\text{Poinc}^{\text{vir}}$ , since the coefficients gather multiple cohomological degrees, it suffices to make sure that each weight can only appear in finitely many cohomological degrees. When  $H^*(\mathfrak{a})$  is finite dimensional, it is easy to see that this is satisfied when  $\text{Quot}_{\text{un}} \mathfrak{a}$  has weight of at least 1, since tensor is additive with respect to weight.

<sup>13</sup>Here, we implicitly choose an identification  $\Lambda \simeq \mathbb{C}$ .

<sup>14</sup>We took this notion from [GL14, Defn. 6.3.1].

5.3.6. The common theme is that if  $\text{DeCat}$  is additive,  $\text{DeCat}(\text{coChev}^{\text{un}} -)$  is, in some sense, “multiplicative.” The main point is that by §3.3, one can reduce the study of  $\text{coChev}$  to that of  $\text{Sym}$ , which is multiplicative.

We learned this approach from [GL14, §6.3], which we will refer to for the proofs. The main difference is that our decategorification processes, with the exception of trace (the last item on the list above), have milder finiteness conditions. This is simply because we have this variable  $t$  to “spread” things out, so we do not have to worry about convergence, since each coefficient will be a finite sum. Another difference is that we consider the  $\text{coLie}$ -coalgebra structure instead of just the underlying complex (see also [GL14, Prop. 6.1.10.]). However, for additive categorification processes, this makes no difference, since §3.3 allows us to reduce to the trivial case. We thus have the following result, whose proof is the same as that of [GR17, Prop. 6.3.4].

**Proposition 5.3.7.** *Let  $\mathfrak{a} \in \text{coLie}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  be a  $\text{coLie}$ -coalgebra such that  $H^*(\mathfrak{a})$  is finite dimensional. Assume that  $\mathfrak{a}$  is equipped with an action of an operator  $T$ . Then, we have*

$$\chi_T^{\text{gr}}(\text{coChev}^{\text{un}} \mathfrak{a}, t) = \chi_T^{\text{gr}}(\text{Sym}(\mathfrak{a}[-1]), t) = \exp \left( - \sum_{n>0} \frac{1}{n} \chi_{T^n}^{\text{gr}}(\mathfrak{a}, t^n) \right).$$

*Remark 5.3.8.* Note that compared to [GL14, Prop. 6.1.10.], we have the extra minus sign because our  $\mathfrak{a}$  and their  $V$  differ by a cohomological shift.

5.3.9. Note that since the exponent is additive in  $\mathfrak{a}$ , and since  $\exp$  is multiplicative,  $\chi_T^{\text{gr}}(\text{coChev}^{\text{un}}, \mathfrak{a})$  is also multiplicative. Namely, we have the following

**Corollary 5.3.10.** *Consider the following push-out diagram of  $\text{coLie}$ -coalgebras equipped with the action of some operator  $T$*

$$\begin{array}{ccc} \mathfrak{a} & \longrightarrow & \mathfrak{b} \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathfrak{c} \end{array}$$

where  $\mathfrak{a}$  and  $\mathfrak{b}$  are with finite dimensional  $H^*(\mathfrak{a})$  and  $H^*(\mathfrak{b})$  (which implies the same for  $\mathfrak{c}$ ). Then, we have

$$\frac{\chi_T^{\text{gr}}(\text{coChev}^{\text{un}} \mathfrak{b})}{\chi_T^{\text{gr}}(\text{coChev}^{\text{un}} \mathfrak{a})} = \chi_T^{\text{gr}}(\text{coChev}^{\text{un}} \mathfrak{b} \otimes_{\text{coChev}^{\text{un}} \mathfrak{a}} \Lambda) = \chi_T^{\text{gr}}(\text{coChev}^{\text{un}} \mathfrak{c}).$$

5.3.11. We will now move on to studying  $\text{Poinc}$ . Since  $\text{Poinc}$  is not additive, we can only treat the case where the  $\text{coLie}$ -coalgebras involved are abelian, in which case, the statement is simply about symmetric powers.

**Proposition 5.3.12.** *Let*

$$\mathfrak{a} = \bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}[1] \oplus \mathfrak{u} \oplus \mathfrak{v}, \quad \mathfrak{b} = \bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}[1] \oplus \mathfrak{u}, \quad \mathfrak{c} = \bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}[1] \oplus \mathfrak{v}$$

be strongly unital  $\text{coLie}$ -coalgebras, where  $H^*(\mathfrak{u})$  and  $H^*(\mathfrak{v})$  are finite dimensional, and where the direct sum are taken in the category of  $\text{coLie}$ -coalgebras.<sup>15</sup> Then

$$\frac{\text{Poinc}(\overline{\text{coChev}^{\text{un}} \mathfrak{a}})}{\text{Poinc}(\overline{\text{coChev}^{\text{un}} \mathfrak{b}})} = \text{Poinc}(\overline{\text{coChev}^{\text{un}} \mathfrak{c}}) = \text{Poinc}(\text{oblv}_{\text{gr}}(\text{coChev}^{\text{un}} \mathfrak{v})).$$

*Proof.* We have,

$$\text{coChev}^{\text{un}} \mathfrak{a} \simeq \text{coChev}^{\text{un}} \mathfrak{b} \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \text{coChev}^{\text{un}} \mathfrak{c},$$

<sup>15</sup>I.e. no interaction between different summands.

and hence

$$\overline{\text{coChev}^{\text{un}} \mathfrak{a}} \simeq \overline{\text{coChev}^{\text{un}} \mathfrak{b}} \otimes_{\Lambda} \overline{\text{coChev}^{\text{un}} \mathfrak{c}}.$$

The first equality follows immediately from the multiplicative nature of Poinc. The second equality is due to Proposition 5.2.2.  $\square$

5.3.13. We will now turn to  $\text{Poinc}^{\text{vir}}$ . The upshot of Proposition 5.3.14 below is that when the decategorification procedure is additive, the decategorification of the stable piece can be extracted from the decategorification of the original object, but with the unit removed (hence, the appearance of  $\text{Quot}_{\text{un}}$  in the statement). This is not surprising since  $\text{Quot}_{\text{un}}$  of the  $\text{coLie}$ -coalgebra is equivalent to taking associated graded (see Lemma 5.2.4).

**Proposition 5.3.14.** *Let  $\mathcal{A} \in \text{ComAlg}^{\text{un}}(\text{Vect}^{\mathbb{Z}_{\geq 0}^m})$  where  $\mathcal{A} = \text{coChev}^{\text{un}} \mathfrak{a}$  for some strongly unital  $\text{coLie}$ -coalgebra  $\mathfrak{a} \in \text{coLie}^{\text{un}}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$ . Assume further that  $\mathfrak{a}$  is equipped with a weight filtration,  $H^*(\mathfrak{a})$  is finite dimensional, and moreover, the weights of  $H^*(\text{Quot}_{\text{un}}(\mathfrak{a}))$  is at least 1. Then, we have*

$$\text{Poinc}^{\text{vir}}(\overline{\text{coChev}^{\text{un}} \mathfrak{a}}, t) = \text{Poinc}^{\text{vir}}(\text{Sym}(\text{oblv}_{\text{gr}}(\text{Quot}_{\text{un}} \mathfrak{a}[-1])), t) = \exp \left( - \sum_{n>0} \frac{1}{n} \text{Poinc}^{\text{vir}}(\text{Quot}_{\text{un}}(\mathfrak{a}), t^n) \right).$$

*Proof.* We have,

$$\begin{aligned} \text{Poinc}^{\text{vir}}(\overline{\text{coChev}^{\text{un}} \mathfrak{a}}, t) &= \text{Poinc}^{\text{vir}}(\text{oblv}_{\text{gr}}(\text{ass-gr}(\text{coChev}^{\text{un}} \mathfrak{a})), t) \\ &= \text{Poinc}^{\text{vir}}(\text{oblv}_{\text{gr}}(\text{coChev}^{\text{un}}(\text{Quot}_{\text{un}} \mathfrak{a})), t) \\ &= \text{Poinc}^{\text{vir}}(\text{oblv}_{\text{gr}}(\text{Sym}(\text{Quot}_{\text{un}} \mathfrak{a}[-1])), t) \\ &= \text{Poinc}^{\text{vir}}(\text{Sym}(\text{oblv}_{\text{gr}}(\text{Quot}_{\text{un}} \mathfrak{a}[-1])), t) \\ &= \exp \left( - \sum_{n>0} \frac{1}{n} \text{Poinc}^{\text{vir}}(\text{Quot}_{\text{un}}(\mathfrak{a}), t^n) \right), \end{aligned}$$

where

$$\text{oblv}_{\text{gr}} : \text{Vect}^{\mathbb{Z}_{\geq 0}^m} \rightarrow \text{Vect}$$

is the functor of taking direct sum (see also §4.5.11).

The finite dimensionality of  $H^*(\mathfrak{a})$  ensures that homological stability happens for  $\mathcal{A}$ , by Corollary 5.1.15. In this case, information about the stable homology can be recovered by the associated graded, since  $\text{Poinc}^{\text{vir}}$  is additive, and we get the first equality. The second equality comes from Lemma 5.2.4. The third equality is due to §3.3. The fourth equality is due to the fact that  $\text{oblv}_{\text{gr}}$  is symmetric monoidal. And finally, as above, the last equality can be seen directly, since we are now simply dealing with symmetric algebras. Note that throughout, the conditions imposed on  $\mathfrak{a}$  ensures that for each power of  $t$ , the coefficient is a finite sum, and hence we do not need to worry about well-definedness.  $\square$

**Corollary 5.3.15.** *Suppose we have the following pushout diagram*

$$\begin{array}{ccc} \mathcal{A} & \longrightarrow & \mathcal{B} \\ \downarrow & & \downarrow \\ \Lambda[\mathbb{Z}_{\geq 0}^m] & \longrightarrow & \mathcal{B} \otimes_{\mathcal{A}} \Lambda[\mathbb{Z}_{\geq 0}^m] = \mathcal{Q} \end{array}$$

in  $\text{ComAlg}^{\text{un}}(\mathbb{Z}_{\geq 0}^m)$ . Assume further that,  $\mathcal{A} = \text{coChev}^{\text{un}} \mathfrak{a}$ ,  $\mathcal{B} = \text{coChev}^{\text{un}} \mathfrak{b}$ , and  $\mathcal{Q} = \text{coChev}^{\text{un}} \mathfrak{q}$  where  $\mathfrak{a}$ ,  $\mathfrak{b}$ , and  $\mathfrak{q}$  satisfy the conditions of the Proposition above, and all the maps respect the weight filtrations.

Then

$$(5.3.16) \quad \frac{\text{Poinc}^{\text{vir}}(\overline{\mathcal{B}})}{\text{Poinc}^{\text{vir}}(\overline{\mathcal{A}})} = \text{Poinc}^{\text{vir}}(\overline{\mathcal{B} \otimes_{\mathcal{A}} \Lambda}) = \text{Poinc}^{\text{vir}}(\text{Sym}(\text{oblv}_{\text{gr}}(0 \sqcup_{\mathfrak{a}} \mathfrak{b})[-1])).$$

*Proof.* We have

$$\begin{aligned} \mathrm{Poinc}^{\mathrm{vir}}(\overline{\mathcal{B}} \otimes_{\overline{\mathcal{A}}} \Lambda, t) &= \mathrm{Poinc}^{\mathrm{vir}}(\overline{\mathcal{B}} \otimes_{\mathcal{A}} \Lambda[\mathbb{Z}_{\geq 0}^m], t) \\ &= \mathrm{Poinc}^{\mathrm{vir}}(\overline{\mathrm{coChev}}^{\mathrm{un}} \mathfrak{q}, t) \\ &= \exp \left( - \sum_{n>0} \frac{1}{n} \mathrm{Poinc}^{\mathrm{vir}}(\mathrm{Quot}_{\mathrm{un}}(\mathfrak{q}), t^n) \right), \end{aligned}$$

where the two last equalities are by Proposition 5.3.14 above. But now, we have established the first equality of (5.3.16) since  $\mathrm{Poinc}^{\mathrm{vir}}$  is additive and since we have the following pushout diagram

$$\begin{array}{ccc} \mathrm{Quot}_{\mathrm{un}}(\mathfrak{a}) & \longrightarrow & \mathrm{Quot}_{\mathrm{un}}(\mathfrak{b}) \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathrm{Quot}_{\mathrm{un}}(\mathfrak{q}) \end{array}$$

For the second equality of (5.3.16), observe that we have the pushout diagram

$$\begin{array}{ccc} \mathfrak{a} & \longrightarrow & \mathfrak{b} \\ \downarrow & & \downarrow \\ \bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}[1] & \longrightarrow & \mathfrak{q} \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathrm{Quot}_{\mathrm{un}}(\mathfrak{q}) \end{array}$$

which implies that

$$(5.3.17) \quad \mathrm{Quot}_{\mathrm{un}}(\mathfrak{q}) \simeq 0 \sqcup_{\mathfrak{a}} \mathfrak{b}.$$

□

5.3.18. Finally, we will turn to trace,  $\chi_T$ . As mentioned above, more care has to be taken in this case since we have to deal with convergence issues.

**Proposition 5.3.19.** *Let  $V \in \mathrm{Vect}^{\mathbb{Z}_{\geq 0}^{m, \rightarrow}}$  living in cohomological degrees  $\geq 0$  and satisfies homological stability. Assume that  $V$  is equipped with an action of an operator  $T$  such that*

$$\mathrm{oblv}_{\mathrm{gr}}(\mathrm{ass}\text{-}\mathrm{gr}(V)) = \mathrm{oblv}_{\mathrm{gr}}(\Lambda \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} V)$$

*is  $T$ -summable. Then  $\overline{V}$  is  $T$ -summable, and moreover*

$$\chi_T^{\mathrm{gr}}(\overline{V}, 1) = \chi_T(\mathrm{oblv}_{\mathrm{gr}}(\mathrm{ass}\text{-}\mathrm{gr}(V))) = \chi_T(\mathrm{oblv}_{\mathrm{gr}}(\Lambda \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} V)).$$

*Proof.* For any  $\mathbf{d} \in \mathbb{Z}_{\geq 0}^m$ , using the fact that  $\mathrm{oblv}_{\mathrm{gr}}(\mathrm{ass}\text{-}\mathrm{gr}(V))$  is  $T$ -summable, it is easy to see that  $V_{\mathbf{d}}$  is also  $T$ -summable (see also [GL14, Remark 6.3.3]), and moreover,

$$|V_{\mathbf{d}}|_T \leq |\mathrm{oblv}_{\mathrm{gr}}(\mathrm{ass}\text{-}\mathrm{gr}(V))|_T.$$

Since  $V$  satisfies homological stability, we know that for any cohomological degree  $c$ ,  $\tau_{\leq c} \overline{V}$  is also  $T$ -summable, with

$$|\tau_{\leq c} \overline{V}|_T \leq |\mathrm{oblv}_{\mathrm{gr}}(\mathrm{ass}\text{-}\mathrm{gr}(V))|_T$$

since

$$\tau_{\leq c} \overline{V} \simeq \tau_{\leq c} V_{\mathbf{d}}$$

for some  $\mathbf{d}$ . But now, this implies that  $\overline{V}$  is  $T$ -summable since

$$|\overline{V}|_T = \lim_{c \rightarrow \infty} |\tau_{\leq c} \overline{V}|_T \leq |\mathrm{oblv}_{\mathrm{gr}}(\mathrm{ass}\text{-}\mathrm{gr}(V))|_T,$$



and we are done. □

## 6. FACTORIZATION HOMOLOGY

The discussion we have so far in this paper has no geometry. In this section, we will discuss factorization algebras and factorization homology, which is the geometric tool needed to bridge the two worlds of geometry and algebra. In §6.1–§6.4, we give a brief review of the theory and provide the first link to the geometric objects  $Z_n^m(X)$  that we are interested in. The materials appearing in these subsections are mostly adapted from [FG11, GL14, Gai15a, Gai15b]. In §6.5, we prove the second main result of this paper, which is a statement about homological stability of factorization cohomology. Finally, §6.6 provides a simple construction which allows one to produce new commutative factorization algebras from a given one. It will be used in §7 to compute the commutative factorization algebras responsible for the cohomology of  $Z_n^m(X)$ .

**6.1. Graded commutative factorization algebras.** We will take factorization homology with coefficients in graded commutative factorization algebras. In the introduction of this paper, we briefly reviewed the notion of the Ran space as well as commutative factorization algebras. We will now visit the graded variant of these notions. Most of what we present in this section come from [Gai15b, §4.1] or are otherwise straight-forward adaptations of [FG11, GL14, Gai15a].

**6.1.1. Colored Ran space/prestack.** Let  $X$  be a scheme.  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$  is the prestack defined as follows: for a test scheme  $S$ ,

$$\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)(S) = \{I \subset X(S), \phi : I \rightarrow \mathbb{Z}_{\geq 0, +}^m\}.$$

For  $\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m$ , we define  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^{\mathbf{d}}$  to be the component of  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$  consisting of those points  $(I, \phi)$  such that

$$\sum_{i \in I} \phi(i) = \mathbf{d}.$$

We have

$$\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m) = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m} \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^{\mathbf{d}}.$$

**6.1.2.** Unlike the usual (i.e. ungraded) Ran space, the colored variant is essentially finite dimensional. Indeed, we have the following canonical map

$$\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^{\mathbf{d}} \rightarrow \text{Sym}^{\mathbf{d}} X$$

that is an isomorphism after a sheafification in the topology generated by finite surjective maps (see §1.1.1 for the definition of  $\text{Sym}^{\mathbf{d}} X$ ). Thus, as far as sheaves and cohomology are concerned, the two are the same.

**6.1.3.** Fix an  $m$ -tuple of positive integers  $\mathbf{n}$ , we define  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}$  to be the open subfunctor of  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$  fitting into the following pullback diagram (see §1.5.6 for the definition of  $Z_{\mathbf{n}}^m(X)$ )

$$\begin{array}{ccc} \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}^{\mathbf{d}} & \longrightarrow & Z_{\mathbf{n}}^{\mathbf{d}}(X) \\ \downarrow & & \downarrow \\ \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^{\mathbf{d}} & \longrightarrow & \text{Sym}^{\mathbf{d}} X \end{array}$$

Similar to the above,

$$\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}^{\mathbf{d}} \rightarrow Z_{\mathbf{n}}^{\mathbf{d}}(X)$$

is an isomorphism after a sheafification.



6.1.4. For later use, we will adopt the following notation.

**Notation 6.1.5.** We will use (see §1.5.6 for the definition of  $Z_{\mathbf{d}}^m$ )

$$Z_{\mathbf{n}}^m(X) = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} Z_{\mathbf{n}}^{\mathbf{d}}(X)$$

to denote the disjoint union of all the schemes  $Z_{\mathbf{n}}^{\mathbf{d}}(X)$ , where when  $\mathbf{d} = \mathbf{0}$  we choose, by convention, that  $Z_{\mathbf{n}}^{\mathbf{0}}(X) = \text{pt}$ . In particular,

$$Z_{\infty}^m(X) = \text{Sym } X = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0}^m} \text{Sym}^{\mathbf{d}} X,$$

where the  $\infty$  in the subscript means  $\mathbf{n} = (\infty, \dots, \infty)$ .

We will use  $Z_{\mathbf{n}}^m(X)_{+}$  to denote

$$Z_{\mathbf{n}}^m(X)_{+} = Z_{\mathbf{n}}^m(X) - Z_{\mathbf{n}}^{\mathbf{0}}(X) = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m} Z_{\mathbf{n}}^{\mathbf{d}}(X),$$

and similarly for  $(\text{Sym } X)_{+}$ .

Similarly, for the Ran side, we have

**Notation 6.1.6.** We will use

$$\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}} = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m} \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}^{\mathbf{d}} \subset \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m).$$

to collect all the components of  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}^{\mathbf{d}}$ .

6.1.7. We define  $\text{Conf}_{\mathbf{d}}(X)$  to be the open subscheme of  $\text{Sym}^{\mathbf{d}} X$  such that no point is allowed to collide (i.e. all multiplicities have to be at most 1). And similarly, we define

$$\overset{\circ}{\text{Ran}}(X, \mathbb{Z}_{\geq 0, +}^m) = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m} \overset{\circ}{\text{Ran}}(X, \mathbb{Z}_{\geq 0, +}^m)^{\mathbf{d}}$$

to be the open subprestack of  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$  where  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^{\mathbf{d}}$  fits into the following pullback diagram

$$\begin{array}{ccc} \overset{\circ}{\text{Ran}}(X, \mathbb{Z}_{\geq 0, +}^m)^{\mathbf{d}} & \longrightarrow & \text{Conf}_{\mathbf{d}}(X) \\ \downarrow & & \downarrow \\ \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^{\mathbf{d}} & \longrightarrow & \text{Sym}^{\mathbf{d}} X \end{array}$$

6.1.8. *Graded commutative factorization algebras.* Similar to the case of the usual Ran space,  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$  has a natural commutative semi-group structure given by

$$\text{union} : \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k \rightarrow \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m).$$

One can thus define the  $\otimes^*$ -symmetric monoidal structure on  $\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m))$  by convolution in the usual way: for  $\mathcal{F}_1, \dots, \mathcal{F}_k \in \text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m))$ , we define

$$\mathcal{F}_1 \otimes^* \dots \otimes^* \mathcal{F}_k = \text{union}_!(\mathcal{F}_1 \boxtimes \dots \boxtimes \mathcal{F}_k).$$

6.1.9. As in the non-graded case, this allows us to define the category

$$\text{ComAlg}^*(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)) = \text{ComAlg}(\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m))^{\otimes^*})$$

of commutative algebra objects, as well as the full subcategory

$$\text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m) \subset \text{ComAlg}^*(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)),$$

consisting of factorizable objects, which will call commutative factorization algebras.

For the reader's convenient, let us quickly review what this means. An object  $\mathcal{F} \in \text{ComAlg}^*(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m))$  is equipped with multiplication maps

$$\text{union}_!(\mathcal{F} \boxtimes \cdots \boxtimes \mathcal{F}) \rightarrow \mathcal{F}$$

which, by adjunction, give rise to maps of the form

$$(6.1.10) \quad \mathcal{F} \boxtimes \cdots \boxtimes \mathcal{F} \rightarrow \text{union}^! \mathcal{F}.$$

We say that  $\mathcal{F}$  is factorizable if these maps induce equivalences on  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k_{\text{disj}}$ , i.e. we have

$$(\mathcal{F} \boxtimes \cdots \boxtimes \mathcal{F})|_{\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k_{\text{disj}}} \simeq (\text{union}^! \mathcal{F})|_{\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k_{\text{disj}}}.$$

6.1.11. *Graded commutative algebras.* The category  $\text{Shv}(X)$  of sheaves on  $X$  is equipped with the  $\otimes$ -symmetric monoidal structure. Thus, we can talk about categories of graded commutative algebras in it as in §4.2, i.e.  $\text{ComAlg}(\text{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m})$  and  $\text{ComAlg}(\text{Shv}(X)^{\mathbb{Z}_{\geq 0}^m})$  etc. Sometimes, to emphasize that we are using the  $\otimes$ -monoidal structure, we will write  $\text{ComAlg}^!$  instead of just  $\text{ComAlg}$ .

Alternatively, we can view  $\text{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m}$  as follows. Consider

$$X^{\mathbb{Z}_{\geq 0, +}^m} = \bigsqcup_{\mathbf{d} \in \mathbb{Z}_{\geq 0, +}^m} X,$$

i.e. disjoint copies of  $X$  indexed by  $\mathbb{Z}_{\geq 0, +}^m$ . Using the  $\otimes$ -monoidal structure on  $X$ , and keep tracks of the grading, we obtain a symmetric monoidal structure on  $\text{Shv}(X^{\mathbb{Z}_{\geq 0, +}^m})$  in an obvious way. Moreover, we have a natural equivalence of symmetric monoidal categories

$$\text{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m} \simeq \text{Shv}(X^{\mathbb{Z}_{\geq 0, +}^m}).$$

Using this identification, we will not distinguish between these two categories in what follows, unless confusion is likely to occur.

6.1.12. Let

$$\delta : X^{\mathbb{Z}_{\geq 0, +}^m} \rightarrow \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$$

be the “diagonal” embedding which sends  $x_{\mathbf{d}} \in X(S)$  to the pair  $(I, \phi) = (\{x\}, x \mapsto \mathbf{d})$ .

We have the following pair of adjoint functors

$$\delta_? : \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m}) \rightleftarrows \text{ComAlg}^*(\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m))) : \delta^!$$

where  $\delta^!$  is the functor of  $!$ -restricting to the diagonal, which commutes with limit and hence, admits a left adjoint denoted by  $\delta_?$ .

**Proposition 6.1.13** ([GL14, Thm. 5.6.4]). <sup>16</sup> *The pair of functors  $\delta_?$  and  $\delta^!$  induces an equivalence of categories*

$$\delta_? : \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m}) \simeq \text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m) : \delta^!$$

*Remark 6.1.14.* The curious readers can take a look at [Gai17, §2.3] for a concrete implementation of the functor  $\delta_?$ . We do not need this concrete description in this paper.

<sup>16</sup>The proof given in [GL14] is for the non-graded case. However, the same proof carries over to this setting verbatim.

6.1.15. Let  $\pi : X \rightarrow \text{pt}$  denote the structure map of  $X$ . For a similar reason as above, we have the following pair of adjoint functors

$$\pi_? : \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) \rightleftarrows \text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}) : \pi^!.$$

**Proposition 6.1.16** ([GL14, Expl. 5.6.9]). *We have the following commutative diagram*

$$(6.1.17) \quad \begin{array}{ccc} \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightarrow[\delta^!]{\delta_?} & \text{Fact}^*(X, \mathbb{Z}_{\geq 0,+}^m) \\ \pi_? \downarrow & \swarrow C_c^*(\text{Ran}(X, \mathbb{Z}_{\geq 0,+}^m), -) & \\ \text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}) & & \end{array}$$

More concretely, this Proposition states that the functor of taking cohomology of compact support along the Ran space implements the abstractly defined functor  $\pi_?$ . We will thus call  $\pi_?$  the functor of taking factorization homology.

**6.2. Koszul duality.** We will now review the interaction between  $\pi_?$ ,  $\pi^!$ , and Koszul duality.

6.2.1. Note that since  $\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}$  is pro-nilpotent (see Lemma 4.1.7), we have an equivalence of categories by Koszul duality

$$\text{coPrim}[1] : \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) \rightleftarrows \text{coLie}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) : \text{coChev}.$$

Since the functor  $\pi^!$  is monoidal, its left adjoint,  $\pi_!$  is left-lax monoidal, and hence, the adjoint pair  $\pi_! \dashv \pi^!$  upgrades to a pair of adjoint functors

$$\pi_! : \text{coLie}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) \rightleftarrows \text{coLie}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}) : \pi^!.$$

Since the monoidal functor  $\pi^!$  commutes with limits (being a right adjoint), the right adjoints (and hence, also the left adjoints) in the diagram below commute

$$\begin{array}{ccc} \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightleftharpoons[\text{coChev}]{\text{coPrim}[1]} & \text{coLie}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) \\ \pi^! \updownarrow \pi_? & & \pi^! \updownarrow \pi_! \\ \text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightleftharpoons[\text{coChev}]{\text{coPrim}[1]} & \text{coLie}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}) \end{array}$$

This implies that

$$\pi_? \circ \text{coChev} \simeq \text{coChev} \circ \text{coPrim}[1] \circ \pi_? \circ \text{coChev} \simeq \text{coChev} \circ \pi_! \circ \text{coPrim}[1] \circ \text{coChev} \simeq \text{coChev} \circ \pi_!.$$

Namely, the following diagram commutes

$$(6.2.2) \quad \begin{array}{ccc} \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) & \xleftarrow{\text{coChev}} & \text{coLie}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) \\ \pi_? \downarrow & & \pi_! \downarrow \\ \text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}) & \xleftarrow{\text{coChev}} & \text{coLie}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}) \end{array}$$

Similarly, we have

$$\pi^! \circ \text{coPrim}[1] \simeq \text{coPrim}[1] \circ \pi^!.$$

**6.3. Factorization cohomology.** We will now explain the process of taking factorization cohomology. Since the theory of sheaves on prestacks are biased toward the functor  $(-)^!$  and  $(-)_!$ , in general, one can only talk about homology rather than cohomology. However, suppose  $X$  admits a compactification

$$j : X \rightarrow \bar{X},$$

we can make sense of it in such a way that is compatible with all the functors we have defined so far. This is the goal of this subsection.

6.3.1. Let

$$j : X \rightarrow \bar{X}$$

be as above.

First, observe that both functors

$$j^* \simeq j^! : \mathrm{Shv}(\bar{X}) \rightleftarrows \mathrm{Shv}(X) : j_*$$

are symmetric monoidal. Hence, we have the following diagram

$$\begin{array}{ccc} \mathrm{ComAlg}^!(\mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightleftharpoons[\mathrm{coChev}]{\mathrm{coPrim}[1]} & \mathrm{coLie}^!(\mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) \\ j^* \updownarrow j_* & & j^* \updownarrow j_* \\ \mathrm{ComAlg}^!(\mathrm{Shv}(\bar{X})^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightleftharpoons[\mathrm{coChev}]{\mathrm{coPrim}[1]} & \mathrm{coLie}^!(\mathrm{Shv}(\bar{X})^{\mathbb{Z}_{\geq 0,+}^m}) \end{array}$$

where parallel arrows are adjoint pairs. Moreover, a priori, left/right adjoints commute with left/right adjoints, but since all the horizontal maps are equivalences everything commutes.

6.3.2. Arguing similarly, we have the following diagram, where everything commutes

$$\begin{array}{ccc} \mathrm{ComAlg}^!(\mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightleftharpoons[\delta^!]{\delta^?} & \mathrm{Fact}^*(X, \mathbb{Z}_{\geq 0,+}^m) \\ j^* \updownarrow j_* & & j^* \updownarrow j_* \\ \mathrm{ComAlg}^!(\mathrm{Shv}(\bar{X})^{\mathbb{Z}_{\geq 0,+}^m}) & \xrightleftharpoons[\delta^!]{\delta^?} & \mathrm{Fact}^*(\bar{X}, \mathbb{Z}_{\geq 0,+}^m) \end{array}$$

**Definition 6.3.3.** We define the functor of taking factorization cohomology

$$\pi_{?,*} : \mathrm{ComAlg}^!(\mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) \rightarrow \mathrm{ComAlg}^!(\mathrm{Vect}^{\mathbb{Z}_{\geq 0,+}^m})$$

to be

$$\pi_{X,?,*} = \pi_{\bar{X},?} \circ j_*$$

where  $\pi_X$  and  $\pi_{\bar{X}}$  are structure maps of  $X$  and  $\bar{X}$  respectively. Namely,

$$\mathrm{ComAlg}^!(\mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m}) \xrightarrow{j_*} \mathrm{ComAlg}^!(\mathrm{Shv}(\bar{X})^{\mathbb{Z}_{\geq 0,+}^m}) \xrightarrow{\pi_{?}} \mathrm{ComAlg}(\mathrm{Vect}^{\mathbb{Z}_{\geq 0,+}^m}).$$

$\pi_{?,*}$

*Remark 6.3.4.* Observe that

$$\begin{aligned} \pi_{?,*} &\simeq \pi_{?} \circ j_* \simeq C_c^*(\mathrm{Ran}(\bar{X}, \mathbb{Z}_{\geq 0,+}^m), \delta_{?}(j_*(-))) \\ &\simeq C_c^*(\mathrm{Ran}(\bar{X}, \mathbb{Z}_{\geq 0,+}^m), j_*(\delta_{?}(-))) \simeq C^*(\mathrm{Ran}(X, \mathbb{Z}_{\geq 0,+}^m), \delta_{?}(-)), \end{aligned}$$

where the last term can be made sense of using the fact that as far as sheaves and cohomology are involved,  $\mathrm{Ran}(X, \mathbb{Z}_{\geq 0,+}^m)^{\mathbf{d}}$  is the same as  $\mathrm{Sym}^{\mathbf{d}} X$ . This implies that  $\pi_{?,*}$  does not depend on the choice of the compactification  $\bar{X}$  of  $X$ .

6.3.5. Note that the discussion above also implies that our functor  $\pi_{?*}$  behaves nicely with respect to  $\text{coChev}$ . Indeed, let  $\mathfrak{a} \in \text{coLie}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m})$ . Then

$$(6.3.6) \quad \pi_{?*}(\text{coChev } \mathfrak{a}) \simeq \pi_?(j_*(\text{coChev } \mathfrak{a})) \simeq \pi_?(\text{coChev}(j_*\mathfrak{a})) \simeq \text{coChev}(\pi_!(j_*\mathfrak{a})) \simeq \text{coChev}(\pi_*\mathfrak{a}),$$

where we have used  $\pi$  to denote both the structure map of  $X$  and of  $\bar{X}$ .

The  $\text{coLie}$ -coalgebra structure of  $\pi_*\mathfrak{a}$  (in (6.3.6)) is induced by the fact that

$$\pi_* : \text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m} \rightarrow \text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}$$

is left-lax symmetric monoidal. Indeed, this is because

$$\pi_* \simeq \pi_! \circ j_*,$$

where  $j_*$  is monoidal, and  $\pi_!$  is left-lax monoidal.

6.3.7. When our  $\text{coLie}$ -coalgebra has the form  $\pi^!\mathfrak{a}$  for some  $\mathfrak{a} \in \text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}$ , we have

$$(6.3.8) \quad \pi_*\pi^!\mathfrak{a} \simeq C^*(X, \omega_X) \otimes \mathfrak{a} \simeq \pi_*\omega_X \otimes \mathfrak{a}.$$

Since  $C^*(X, -)$  is left-lax monoidal,  $C^*(X, \omega_X) \in \text{Vect}$  has a natural co-commutative coalgebra structure since  $\omega_X$  does, namely

$$\omega_X \overset{!}{\otimes} \omega_X \simeq \omega_X.$$

The  $\text{coLie}$ -coalgebra structure on  $\pi_*\omega_X \otimes \mathfrak{a}$  is then induced by that of  $\mathfrak{a}$  and the co-commutative co-algebra structure on  $C^*(X, \omega_X)$ . See [GR17, §IV.2.1.2] and [Ho17, Expl. 4.2.8] for a discussion of the dual situation, where one tensors a Lie-algebra with a commutative algebra to get a new Lie-algebra. See also [Ho17, §5.3] for a computational example where the resulting Lie-algebra is abelian.

*Remark 6.3.9.* When the co-algebra  $C^*(X, \omega_X)$  is trivial (for eg. when  $X = \mathbb{A}^d$ ),  $\pi_*\omega_X \otimes \mathfrak{a}$  is abelian. In many cases, this can be exploited to compute the factorization cohomology in the case where  $X = \mathbb{A}^d$ .

6.3.10. Let  $\mathcal{A} \in \text{ComAlg}^{\text{un, au}}(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m})$  (see §2.3) with augmentation ideal  $\mathcal{A}_+$ . We will use

$$\pi_{?*}^{\text{un}} \mathcal{A} = \Lambda \oplus \pi_{?*}(\mathcal{A}_+)$$

to denote the unital factorization cohomology. It is immediate that when the Koszul dual of  $\mathcal{A}_+$  is  $\mathfrak{a}$ , we have

$$\pi_{?*}^{\text{un}} \mathcal{A} \simeq \text{coChev}^{\text{un}}(\pi_*\mathfrak{a}).$$

When  $\mathcal{A} \in \text{ComAlg}(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m})$ , we will abuse notation and write

$$\pi_{?*}^{\text{un}} \mathcal{A} = \Lambda \oplus \pi_{?*}\mathcal{A}.$$

If  $\mathfrak{a}$  is the Koszul dual of  $\mathcal{A}$ , then we still have

$$\pi_{?*}^{\text{un}} \mathcal{A} \simeq \text{coChev}^{\text{un}}(\pi_*\mathfrak{a}).$$

**6.4. A multiplicative trace formula.** We will now come to an analog of the Grothendieck-Lefschetz trace formula, but for the functor  $\pi_{?*}$  instead of  $\pi_*$ . If  $\pi_*$  is an additive functor, in some sense, we will see that  $\pi_{?*}$  is multiplicative. The result presented here is a straightforward adaptation of [GL14, §6.5]. Our case is simpler since we do not have to worry about convergence issues.

We start with the following finiteness condition.

**Definition 6.4.1.** For a category  $\mathcal{C}$ , we will use  $\mathcal{C}^{c, \mathbb{Z}_{\geq 0}^m}$  to denote the fullsubcategory of  $\mathcal{C}^{\mathbb{Z}_{\geq 0}^m}$  consisting objects that are compact in each degree.

For example,  $\text{Shv}(X)^{c, \mathbb{Z}_{\geq 0}^m}$  consists of graded-sheaves that are constructible in each graded-degree;  $\text{Vect}^{c, \mathbb{Z}_{\geq 0}^m}$  consists of graded-chain complexes such that are bounded (on both sides) at each graded-degree.

6.4.2. It is easy that for  $\mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0}^m}$  and  $\mathrm{Vect}^{\mathbb{Z}_{\geq 0}^m}$ ,  $\mathrm{coChev}$  and  $\mathrm{coPrim}[1]$  preserves compactness. Indeed, this is because when we restrict to each graded degrees, the limits/colimits computing them are finite (essentially the same argument used in the case of pro-nilpotent categories). But now, for schemes, since the functor  $\pi_*$  preserves compactness, so is the functor  $\pi_{?,*}$  since it's computed as

$$\pi_{?,*} \simeq \mathrm{coChev} \circ \pi_* \circ \mathrm{coPrim}[1].$$

**Proposition 6.4.3.** *When the scheme  $X$  and  $\mathcal{A} \in \mathrm{ComAlg}^{\mathrm{un}, \mathrm{au}}(\mathrm{Shv}(X)^{c, \mathbb{Z}_{\geq 0}^m})$  are obtained from pulling back objects defined over a finite field  $\mathbb{F}_q$ . Then  $\pi_{?,*}\mathcal{A}$  naturally carries a Frobenius action and we have the following identity*

$$\chi_{\mathrm{Frob}_q^{-1}}^{\mathrm{gr}}(\pi_{?,*}^{\mathrm{un}} \mathcal{A}, t) = \prod_{x \in |X|} \chi_{\mathrm{Frob}_x^{-1}}^{\mathrm{gr}}(i_x^! \mathcal{A}, t^{\deg(x)}),$$

where  $\mathrm{Frob}^{-1}$  denotes the inverse of the geometric Frobenius, and  $|X|$  denotes the set of all closed points of  $X$ .

Before we start the proof of Proposition 6.4.3, let us recall the following (Verdier-dual) form of the Grothendieck-Lefschetz formula.

**Theorem 6.4.4** (Grothendieck). *Let  $X$  be a scheme and  $\mathcal{F} \in \mathrm{Shv}(X)$  a constructible sheaf. Assume that they both are pullbacks of objects defined over a finite field  $\mathbb{F}_q$ . Then, we have the following equality*

$$\chi_{\mathrm{Frob}_q^{-1}}(C^*(X, \mathcal{F})) = \sum_{x \in X(\mathbb{F}_q)} \chi_{\mathrm{Frob}_q^{-1}}(i_x^! \mathcal{F}),$$

where  $\chi_{\mathrm{Frob}_q^{-1}}(-)$  denotes the trace of  $\mathrm{Frob}_q^{-1}$ .<sup>17</sup>

*Proof of Proposition 6.4.3.* Let  $\mathfrak{a}$  be the Koszul dual of  $\mathcal{A}_+$ . Then for each  $x \in |X|$ ,  $i_x^! \mathfrak{a}$  is the Koszul dual of  $i_x^! \mathcal{A}_+$ , since  $i_x^!$  is symmetric monoidal and commute with limits. In particular,

$$(6.4.5) \quad i_x^! \mathcal{A} \simeq \mathrm{coChev}^{\mathrm{un}}(i_x^! \mathfrak{a}).$$

---

<sup>17</sup>This is sometimes denoted as  $\mathrm{tr}(\mathrm{Frob}_q^{-1}, -)$ .

We have

$$\begin{aligned}
& \lambda_{\mathrm{Frob}_q^{-1}}^{\mathrm{gr}}(\pi_{? *}^{\mathrm{un}} \mathcal{A}, t) \\
(\S 6.3.10) \quad &= \lambda_{\mathrm{Frob}_q^{-1}}^{\mathrm{gr}}(\mathrm{coChev}^{\mathrm{un}} \pi_* \mathfrak{a}, t) \\
(\text{Proposition 5.3.7}) \quad &= \exp \left( - \sum_{n>0} \frac{1}{n} \lambda_{\mathrm{Frob}_q^{-n}}^{\mathrm{gr}}(\mathfrak{a}, t^n) \right) \\
(\text{Theorem 6.4.4}) \quad &= \exp \left( - \sum_{n>0} \sum_{x \in X(\mathbb{F}_{q^n})} \frac{1}{n} \lambda_{\mathrm{Frob}_q^{-n}}^{\mathrm{gr}}(i_x^! \mathfrak{a}, t^n) \right) \\
&= \exp \left( - \sum_{n>0} \sum_{\substack{x \in |X| \\ \deg x | n}} \frac{\deg x}{n} \lambda_{\mathrm{Frob}_x^{-n/\deg x}}^{\mathrm{gr}}(i_x^! \mathfrak{a}, t^n) \right) \\
&= \exp \left( - \sum_{x \in |X|} \sum_{m>0} \frac{1}{m} \lambda_{\mathrm{Frob}_x^{-m}}^{\mathrm{gr}}(i_x^! \mathfrak{a}, t^{m \deg x}) \right) \\
&= \prod_{x \in |X|} \exp \left( - \sum_{m>0} \frac{1}{m} \lambda_{\mathrm{Frob}_x^{-m}}^{\mathrm{gr}}(i_x^! \mathfrak{a}, t^{m \deg x}) \right) \\
(\text{Proposition 5.3.7 and (6.4.5)}) \quad &= \prod_{x \in X} \lambda_{\mathrm{Frob}_x^{-1}}^{\mathrm{gr}}(i_x^! \mathcal{A}, t^{\deg x}).
\end{aligned}$$

□

**6.5. Homological stability.** We will now come to the second main result of the paper: homological stability of factorization cohomology with commutative factorization algebra coefficients. Thanks to the preparation that we have done so far, the result is now quite straightforward to prove.

6.5.1. In this subsection, we will work with irreducible schemes of dimension  $d$ . Let  $X$  be such a scheme. Then

$$\dim H^{-2d}(X, \omega_X) = \dim H_c^{2d}(X, \Lambda) = 1,$$

where the first equality is due to Verdier duality. Moreover, we have a natural map of algebras

$$C_c^*(X, \Lambda) \rightarrow H_c^{2d}(X, \Lambda)[-2d] \simeq \Lambda[-2d](-d)$$

inducing an equivalence at cohomological degree  $2d$ , where the RHS is equipped with the trivial algebra structure. Dually, we have the following map of co-algebras

$$\Lambda[2d](d) \rightarrow C^*(X, \omega_X),$$

inducing an equivalence at cohomological degree  $-2d$ , where the co-algebra structure on the LHS is trivial.

6.5.2. In parallel to the situation at §5.1, we have the following

**Definition 6.5.3.** An  $\mathfrak{a} \in \mathrm{coLie}(\mathrm{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  is said to be  $d$ -shifted unital if it is equipped with a map of  $\mathrm{coLie}$ -coalgebras<sup>18</sup>

$$\bigoplus_{k=1}^m \Lambda_{1_k}[-2d+1](-d) \rightarrow \mathfrak{a},$$

where the LHS is equipped with the trivial  $\mathrm{coLie}$ -coalgebra structure.

<sup>18</sup>Note that the extra Tate twist is to make sure that we will eventually fall into the setting of Remark 5.1.17.

Such an  $\mathfrak{a}$  is said to be  $d$ -shifted strongly unital if it lives in cohomological degrees  $\geq 2d - 1$  and the map above induces an equivalence at cohomological degree  $2d - 1$ .

**Lemma 6.5.4.** *Let  $X$  be an irreducible scheme of dimension  $d$ , and  $\mathfrak{a} \in \text{coLie}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  is a  $d$ -shifted strongly unital  $\text{coLie}$ -coalgebra. Then*

$$C^*(X, \omega_X) \otimes \mathfrak{a} \in \text{coLie}^{\text{un}}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$$

*is strongly unital.*

*Proof.* Indeed, the unit map is given by

$$\bigoplus_{k=1}^m \Lambda_{1_k}[1] \simeq \Lambda[2d](d) \otimes \bigoplus_{k=1}^m \Lambda_{1_k}[-2d+1](-d) \rightarrow C^*(X, \omega_X) \otimes \mathfrak{a}.$$

□

6.5.5. We are now in the purview of Theorem 5.1.12, which could be used to investigate homological stability of the factorization cohomology of  $X$  with coefficient in  $\pi^!(\text{coChev } \mathfrak{a}) \simeq \text{coChev}(\pi^! \mathfrak{a})$  since, by (6.3.6) and (6.3.8),

$$\pi_{?*}(\pi^! \text{coChev } \mathfrak{a}) \simeq \text{coChev}(C^*(X, \omega_X) \otimes \mathfrak{a}).$$

In particular, the following result follows immediately from Corollary 5.1.15.

**Theorem 6.5.6.** *Let  $X$  be an irreducible scheme of dimension  $d$ . Let  $\mathcal{A} \in \text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$  such that  $\mathfrak{a} = \text{coPrim}[1](\mathcal{A})$  is a finite dimensional  $d$ -shifted strongly unital  $\text{coLie}$ -coalgebra. Then,*

$$\pi_{?*}^{\text{un}} \pi^! \mathcal{A} \simeq \text{coChev}^{\text{un}}(C^*(X, \omega_X) \otimes \mathfrak{a})$$

*satisfies homological stability.*

A couple of remarks are in order.

*Remark 6.5.7.* Morally speaking, Theorem 6.5.6 states that factorization cohomology of an irreducible scheme  $X$  of dimension  $d$  with coefficients in a graded commutative algebra  $\mathcal{A}$  satisfies homological stability if the cotangent fiber of  $\mathcal{A}$  is finite dimensional.<sup>19</sup> The extra unital assumption is there just so that we can construct stabilization maps and hence to make sense of questions about homological stability.

In this sense, this result is closely related to the one of [KM18]. Indeed, the finiteness condition on  $\mathfrak{a}$  (or more generally, the bound on the cohomological amplitudes stated in Theorem 5.1.12) could be viewed as the analog of the bounded generation property of loc. cit. As we mentioned in Remark 5.1.16, this Theorem can be extended beyond the case of finite dimensionality. We choose not to do that since the numerics involved would look somewhat complicated,<sup>20</sup> and moreover, in practice, we can just apply Theorem 5.1.12 directly to get stability if we care about the stable range.

*Remark 6.5.8.* Note that by Theorem 5.1.12, the homological stability range of  $\pi_{?*} \mathcal{A}$  depends solely on the cohomological amplitudes of  $H^*(X, \omega_X)$  and  $H(\mathfrak{a})$ . In §7, we will present many examples where the ranges for homological stability can be easily extracted from Theorem 5.1.12.

**6.6. Variants.** For the application of this paper, we also need various variants of the  $\otimes^*$ -monoidal structure, along with the corresponding notions of commutative algebras and commutative factorization algebras. More specifically, we will need an analog of the  $\otimes^*$ -monoidal structure on  $\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_n)$ . Just like in the case of  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$ , this is done by equipping  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_n$  with the structure of a commutative semi-group, and then, the monoidal structure is defined as convolution. However, this time, it is a commutative semi-group object in the category  $\text{Corr}(\text{PreStk})$  of correspondences in prestacks.

This theory is developed in [Ras15]. We used it in [Ho17] to study the cohomology of configuration spaces. The construction we give here is a variant of the one given there.

<sup>19</sup>Cotangent fiber in the sense [GL14, §6.1]. This is linked to our  $\text{coPrim}[1]$  via [GL14, Thm. 6.1.10].

<sup>20</sup>Even though it would be just some shift of the conditions listed in Theorem 5.1.12.



6.6.1. Since  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$  is a semi-group object in  $\text{PreStk}$ , it is automatically a semi-group object in  $\text{Corr}(\text{PreStk})$ , i.e. the multiplication maps are given by

$$\begin{array}{ccc} & \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k & \\ \swarrow & & \searrow \text{union} \\ \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k & & \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m) \end{array}$$

From this, we can equip  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}$  with the structure of a semi-group object by pulling back the multiplication map (i.e. union) along the open morphism

$$\iota_{\mathbf{n}} : \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}} \hookrightarrow \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m).$$

Namely, the commutative semi-group structure on  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}$  is given by

$$(6.6.2) \quad \begin{array}{ccc} & \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k \times_{\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)} \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}} & \\ \swarrow j & & \searrow \text{union} \\ (\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})^k & & \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}} \end{array}$$

Note that union is pseudo-proper, since it is a pullback of a pseudo-proper morphism. Moreover, we see easily that  $j$  is an open embedding.

6.6.3. We are now ready to define the  $\otimes^*$ -monoidal structure on  $\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})$ : for  $\mathcal{F}_1, \dots, \mathcal{F}_k \in \text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})$ , we define

$$\mathcal{F}_1 \otimes^* \dots \otimes^* \mathcal{F}_k = \text{union}_! j^!(\mathcal{F}_1 \boxtimes \dots \boxtimes \mathcal{F}_k) \simeq \text{union}_! j^*(\mathcal{F}_1 \boxtimes \dots \boxtimes \mathcal{F}_k),$$

where the equivalence is due to the fact that  $j$  is an open embedding.

As in the case of  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$ , we will use

$$\text{ComAlg}^*(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}) = \text{ComAlg}(\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})^{\otimes^*})$$

to denote the category of commutative algebra objects in  $\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})$  with respect to the  $\otimes^*$ -monoidal structure.

We define the full subcategory

$$\text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}} \subset \text{ComAlg}^*(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})$$

consisting of factorizable objects, namely, those commutative algebras  $\mathcal{F}$  such that the maps (obtained from the multiplication maps by adjunction)

$$(6.6.4) \quad j^!(\mathcal{F} \boxtimes \dots \boxtimes \mathcal{F}) \rightarrow \text{union}_! \mathcal{F}$$

induces equivalences

$$(j^!(\mathcal{F} \boxtimes \dots \boxtimes \mathcal{F}))|_{(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})^k_{\text{disj}}} \rightarrow (\text{union}_! \mathcal{F})|_{(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})^k_{\text{disj}}}.$$

6.6.5. We will now study the interaction between the  $\otimes^*$ -monoidal structures on  $\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})$  and on  $\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m))$ . We start with the following

**Lemma 6.6.6.** *The functor*

$$\iota_{\mathbf{n}}^* \simeq \iota_{\mathbf{n}}^* : \text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)) \rightarrow \text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}})$$

is symmetric monoidal with respect to the  $\otimes^*$ -monoidal structures on both sides. As a result, its right adjoint  $\iota_{\mathbf{n},*}$  is right-lax monoidal.

*Proof.* This follows immediately from the base-change theorem for pseudo-proper morphisms and the fact that the semi-group structure on  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_n$  is defined via pulling-back along the pseudo-proper map union for  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$ .  $\square$

**Corollary 6.6.7.** *The adjoint pair  $\iota_n^* \dashv \iota_{n,*}$  upgrades to a pair of adjoint functors between the corresponding commutative algebra objects:*

$$\iota_n^* \simeq \iota_n^! : \text{ComAlg}^*(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)) \rightleftarrows \text{ComAlg}^*(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_n) : \iota_{n,*}.$$

6.6.8. The adjoint pair above also behaves nicely with respect to factorizability.

**Lemma 6.6.9.** *The functors  $\iota_n^*$  and  $\iota_{n,*}$  preserve factorizability. Hence, we have a pair of adjoint functors*

$$\iota_n^! \simeq \iota_n^* : \text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m) \rightleftarrows \text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m)_n : \iota_{n,*}.$$

*Proof.* We start with the case for  $\iota_n^!$ . The statement follows from the following pullback diagram

$$\begin{array}{ccc} (\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_n)^k_{\text{disj}} & \longrightarrow & \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k \times_{\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)} \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_n \\ \downarrow & & \downarrow j \\ \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k_{\text{disj}} & \longrightarrow & \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k \end{array}$$

Indeed, for  $\mathcal{F} \in \text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m)$ , the maps

$$j^!(\mathcal{F} \boxtimes \cdots \boxtimes \mathcal{F}) \rightarrow \text{union}^! \mathcal{F},$$

of (6.6.4) is obtained from the corresponding map for  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$ , (6.1.10), by pulling along  $j$ . Proving factorizability amounts to showing that pulling this back further to  $(\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_n)^k_{\text{disj}}$  along the top horizontal arrow gives an equivalence. But by functoriality of  $(-)^!$ , we can pullback along the other circuit of the diagram above. As an intermediate step, however, when we pull back to  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)^k_{\text{disj}}$ , we get an equivalence, provided by the given factorizability condition on  $\text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$ , and we are done.

The case for  $\iota_{n,*}$  is argued similarly, using base change for  $(-)_*$  and  $(-)^!$  in place of functoriality of  $(-)^!$ .  $\square$

6.6.10. Now, we want to understand what  $\iota_{n,*} \iota_n^!$  does to commutative factorization algebras. More specifically, we want to understand the following composition of functors (the dotted arrow below)

$$\begin{array}{ccc} \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m}) & \xrightarrow[\simeq]{\delta^?} & \text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m) \\ \vdots \downarrow & & \downarrow \iota_n^! \\ & & \text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m)_n \\ & & \downarrow \iota_{n,*} \\ \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m}) & \xleftarrow[\simeq]{\delta^!} & \text{Fact}^*(X, \mathbb{Z}_{\geq 0, +}^m) \end{array}$$

Consider the following pullback diagram

$$\begin{array}{ccccc} (X^{\mathbb{Z}_{\geq 0, +}^m})_n & \xrightarrow{\delta} & \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_n & \longrightarrow & Z_n^m(X) \\ \iota_n \downarrow & & \downarrow \iota_n & & \downarrow \\ X^{\mathbb{Z}_{\geq 0, +}^m} & \xrightarrow{\delta} & \text{Ran}(X, \mathbb{Z}_{\geq 0, +}^m) & \longrightarrow & Z_\infty^m(X) \end{array}$$

where  $(X^{\mathbb{Z}_{\geq 0, +}^m})_{\mathbf{n}} \subset X^{\mathbb{Z}_{\geq 0, +}^m}$  consists of copies of  $X$  indexed by  $\mathbf{d}$  where for at least one  $k, 1 \leq k \leq m$ , we have  $\mathbf{d}_k < \mathbf{n}_k$ . Now,

$$\delta^! \circ \iota_{\mathbf{n},*} \circ \iota_{\mathbf{n}}^! \circ \delta_? \simeq \iota_{\mathbf{n},*} \circ \delta^! \circ \iota_{\mathbf{n}}^! \circ \delta_? \simeq \iota_{\mathbf{n},*} \circ \iota_{\mathbf{n}}^! \circ \delta^! \circ \delta_? \simeq \iota_{\mathbf{n},*} \circ \iota_{\mathbf{n}}^!,$$

where the first equivalence is by base change, the second by commutativity of the diagram, and the third by Proposition 6.1.13. But now,  $\iota_{\mathbf{n},*} \circ \iota_{\mathbf{n}}^!$  (on the left side of the diagram) is easy to understand: for  $\mathcal{F} \in \mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m}$  we have

$$(\iota_{\mathbf{n},*}(\iota_{\mathbf{n}}^! \mathcal{F}))_{\mathbf{d}} \simeq \begin{cases} 0, & \text{if } \mathbf{d}_k \geq \mathbf{n}_k, 1 \leq k \leq m, \\ \mathcal{F}_{\mathbf{d}}, & \text{otherwise.} \end{cases}$$

We have thus proved the following

**Proposition 6.6.11.** *Let  $\mathcal{A} \in \mathrm{ComAlg}^1(\mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0, +}^m})$ , then the commutative algebra  $\mathcal{A}' = \delta^!(\iota_{\mathbf{n},*}(\iota_{\mathbf{n}}^!(\delta_?(\mathcal{A}))))$  is obtained from  $\mathcal{A}$  by setting all components  $\mathcal{A}_{\mathbf{d}}$  to 0, for all  $\mathbf{d}$  such that  $\mathbf{d}_k \geq \mathbf{n}_k$ , for all  $1 \leq k \leq m$ .*

## 7. COHOMOLOGY OF $Z_{\mathbf{n}}^m(X)$ AS FACTORIZATION COHOMOLOGY

We will now use the theory developed so far to study cohomology of the spaces  $Z_{\mathbf{n}}^m(X)$  (see Notation 6.1.5). As we have seen, the cohomology of  $Z_{\mathbf{n}}^m(X)_+$  is equivalent to that of  $\mathrm{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)_{\mathbf{n}}$ , and similarly for  $Z_{\infty}^m(X)_+$  and  $\mathrm{Ran}(X, \mathbb{Z}_{\geq 0, +}^m)$ . We can thus perform our manipulations on the  $\mathrm{Ran}$  side.

Since this section denotes different (but related) objects with the same letter, differed only by the fonts used, let us quickly orient the readers with the notation used. In general, we adopt the following convention: the “fancier” the object, the “fancier” the font used to denote it. We start with augmented unital algebras  $A_{m, \mathbf{n}}$  in  $\mathrm{Vect}^{\mathbb{Z}_{\geq 0}^m}$ , along with its augmentation ideal  $(A_{m, \mathbf{n}})_+$ , and its Koszul dual  $a_{m, \mathbf{n}}$ . These objects are very simple, as reflected by the font used. !-pulling back to  $X$  gives us corresponding objects in  $\mathrm{Shv}(X)^{\mathbb{Z}_{\geq 0}^m}$ , which we will denote by  $A_{m, \mathbf{n}}(X)$ ,  $(A_{m, \mathbf{n}}(X))_+$ , and  $a_{m, \mathbf{n}}(X)$  respectively. Taking factorization cohomology of  $(A_{m, \mathbf{n}}(X))_+$ , we obtain  $(\mathcal{A}_{m, \mathbf{n}}(X))_+$ , which is the augmentation ideal of  $\mathcal{A}_{m, \mathbf{n}}(X)$ , and which has Koszul dual  $a_{m, \mathbf{n}}(X)$ . These objects are more complicated, also reflected by the font used. Indeed,  $\mathcal{A}_{m, \mathbf{n}}(X)$  is precisely the cohomology of  $Z_{\mathbf{n}}^m(X)$ . Finally, we use  $\overline{\mathcal{A}}_{m, \mathbf{n}}(X)$  to denote the stabilization of  $\mathcal{A}_{m, \mathbf{n}}(X)$ .

We will now briefly review the content of this section. We start, in §7.1, with the definition of the algebras  $A_{m, \mathbf{n}}$  and show that the factorization cohomology with coefficients in  $A_{m, \mathbf{n}}(X)$  computes the cohomology of  $Z_{\mathbf{n}}^m(X)$ . The algebras  $A_{m, \mathbf{n}}$  are of a very simple nature: they are the quotient of a symmetric algebra  $A_{m, \infty}$  by one monomial. In §7.2, we perform the simple computation of  $\Lambda \otimes_{A_{m, \infty}} A_{m, \mathbf{n}}$  where coincidences readily manifest themselves. Via factorization cohomology, these coincidences get translated to the level of cohomology of the spaces  $Z_{\mathbf{n}}^m(X)$ . In §7.3, we compute the Koszul duals of the  $A_{m, \mathbf{n}}$ ’s. It is shown in §7.4 that homological stability for  $Z_{\mathbf{n}}^m(X)$  is now a direct consequence of Theorem 5.1.12. In §7.5, building on the coincidences among the densities of  $\mathcal{A}_{m, \mathbf{n}}(X)$ , we prove coincidences at the limits, i.e. of  $\overline{\mathcal{A}}_{m, \mathbf{n}}(X)$ . In §7.6, we decategorify these coincidences to get interesting identities, some of which are new, others are known previously but only as combinatorial coincidences. Finally, in §7.7, through a simple  $L_{\infty}$ -algebra computation, we deal with the case of Poincaré polynomials, which is less well-behaved. Using the same technique, we obtain a simple expression for the stable homological density  $\Lambda \otimes_{\overline{\mathcal{A}}_{m, \infty}(X)} \overline{\mathcal{A}}_{m, \mathbf{n}}(X)$ .

Throughout this section, we fix an irreducible scheme  $X$  of dimension  $d$ . In what follows, the algebras that we define will have their cohomological gradings depending on this  $d$ . We will, however, suppress it from the notation unless confusion is likely to occur.

**7.1. Commutative factorization algebras.** We will now produce commutative factorization algebras whose factorization cohomology groups compute the cohomology of  $Z_{\mathbf{n}}^m(X)$ .

7.1.1. Let

$$A_{m,\infty} \in \text{ComAlg}^{\text{un,au}}(\text{Vect}^{\mathbb{Z}_{\geq 0}^m})$$

be the free (unital) commutative algebra generated by

$$(7.1.2) \quad \bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}[-2d](-d).$$

We write symbolically

$$A_{m,\infty} = \Lambda[x_1, \dots, x_m]$$

where  $x_k$  sits in graded degree  $\mathbf{1}_k$ , cohomological degree  $2d$ , and  $-d$  Tate twist.

Its augmentation ideal, is

$$(A_{m,\infty})_+ = \Lambda[x_1, \dots, x_m]_+ \in \text{ComAlg}(\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}).$$

7.1.3. For an  $m$ -tuple of positive integers  $\mathbf{n}$ , let

$$A_{m,\mathbf{n}} = A_{m,\infty} / (\prod_{k=1}^m x_k^{\mathbf{n}_k}) = \Lambda[x_1, \dots, x_m] / (\prod_{k=1}^m x_k^{\mathbf{n}_k}).$$

In other words,  $A_{m,\mathbf{n}}$  is obtained from  $A_{m,\infty}$  by setting  $(A_{m,\infty})_{\mathbf{d}}$  to 0 for all  $\mathbf{d} \in \mathbb{Z}_{\geq 0,+}^m$  such that  $\mathbf{d}_k \geq \mathbf{n}_k$  for all  $1 \leq k \leq m$ . Similarly, we let  $(A_{m,\mathbf{n}})_+$  be the augmentation ideal of  $A_{m,\mathbf{n}}$ .

7.1.4. Let  $\pi : X \rightarrow \text{pt}$  be the structure map of  $X$ . We let

$$A_{m,\infty}(X) = \pi^! A_{m,\infty} = \omega_X \otimes A_{m,\infty} \quad \text{and} \quad A_{m,\mathbf{n}}(X) = \pi^! A_{m,\mathbf{n}} = \omega_X \otimes A_{m,\mathbf{n}},$$

where  $\omega_X$  is the dualizable sheaf on  $X$ .

Note that the generators of  $A_{m,\infty}(X)$  and  $A_{m,\mathbf{n}}(X)$  live in cohomological degrees 0 and have no Tate twist, since the generators of  $A_{m,\infty}$  and  $A_{m,\mathbf{n}}$  live in cohomological degrees  $2d$  and have  $-d$  Tate twist.

However, since these are algebras with respect to the  $\otimes$ -monoidal structure, their products do not live in cohomological degrees 0 and have non-trivial Tate twists.

7.1.5. We will make use of the following

**Notation 7.1.6.** Let  $S = \sqcup S_i$  be a possibly infinite disjoint union of irreducible schemes. We will use  $\vec{\omega}_S$  to denote the sheaf  $\omega_S[-2 \dim S](-\dim S)$ , by which we mean the sheaf

$$\bigoplus \omega_{S_i}[-2 \dim S_i](-\dim S_i).$$

Note that when  $S$  is smooth,  $\vec{\omega}_S \simeq \Lambda_S$ .

More generally, when we shift a sheaf by some factor of the dimension of  $S$ , we mean on each component, we shift using the dimension of that factor. The same convention applies to Tate twists as well.

7.1.7. Since  $\pi^!$  is symmetric monoidal,

$$A_{m,\infty}(X)_+ \in \text{ComAlg}^!(\text{Shv}(X)^{\mathbb{Z}_{\geq 0,+}^m})$$

is the free (non-unital) commutative algebra generated by<sup>21</sup>

$$\bigoplus_{k=1}^m (\vec{\omega}_X)_{\mathbf{1}_k}.$$

Thus, by inspecting the free commutative algebra on  $\text{Shv}(\text{Ran}(X, \mathbb{Z}_{\geq 0,+}^m))$  using the  $\otimes^*$ -monoidal structure, we get

$$\delta_? A_{m,\infty}(X)_+ \simeq \vec{\omega}_{\text{Ran}(X, \mathbb{Z}_{\geq 0,+}^m)} \in \text{Fact}^*(X, \mathbb{Z}_{\geq 0,+}^m).$$

By Proposition 6.6.11, we see immediately that

$$\delta_? A_{m,\mathbf{n}}(X)_+ \simeq \iota_{\mathbf{n},*} \iota_{\mathbf{n}}^! \vec{\omega}_{\text{Ran}(X, \mathbb{Z}_{\geq 0,+}^m)} \simeq \iota_{\mathbf{n},*} \vec{\omega}_{\text{Ran}(X, \mathbb{Z}_{\geq 0,+}^m)_{\mathbf{n}}} \in \text{Fact}^*(X, \mathbb{Z}_{\geq 0,+}^m).$$

---

<sup>21</sup>When  $X$  is smooth, the formula becomes simply  $\bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}$ .

The discussion above implies the following result, which links the cohomologies of  $Z_{\mathbf{n}}^m(X)$  to factorization cohomology (see Notation 6.1.5 for the notations, and §6.3 for the discussion of factorization cohomology).

**Proposition 7.1.8.** *We have*

$$\pi_{?*}(A_{m,\mathbf{n}}(X)_+) \simeq C^*(Z_{\mathbf{n}}^m(X)_+, \vec{\omega}_{Z_{\mathbf{n}}^m(X)}).$$

*In particular, when  $X$  is smooth, we have*

$$\pi_{?*}(A_{m,\mathbf{n}}(X)_+) \simeq C^*(Z_{\mathbf{n}}^m(X)).$$

**Notation 7.1.9.** We will use  $\mathcal{A}_{m,\mathbf{n}}(X)$  to denote the algebra  $C^*(Z_{\mathbf{n}}^m(X), \vec{\omega}_{Z_{\mathbf{n}}^m(X)})$ , and

$$\mathcal{A}_{m,\mathbf{n}}(X)_+ = \pi_{?*}(A_{m,\mathbf{n}}(X)_+) \simeq C^*(Z_{\mathbf{n}}^m(X)_+, \vec{\omega}_{Z_{\mathbf{n}}^m(X)_+}),$$

including the case where  $\mathbf{n} = \infty$ .

*Remark 7.1.10.* This result generalizes that of [Knu14]. The algebra structure on  $\mathcal{A}_{m,\mathbf{n}}(X)$  can also be constructed geometrically in a similar way as in [Knu14, §5.2]. We do not need this geometric description in the sequel.

**7.2. Densities.** We will now perform an easy, but fundamental, computation, Proposition 7.2.2, from which coincidences observed at the level of cohomology is an easy consequence, Theorem 7.2.4.

Consider the following pushout in  $\text{ComAlg}^{\text{un, au}}(\text{Vect}^{\mathbb{Z}_{\geq 0}^m})$

$$(7.2.1) \quad \begin{array}{ccc} A_{m,\infty} & \longrightarrow & A_{m,\mathbf{n}} \\ \downarrow & & \downarrow \\ \Lambda & \longrightarrow & \Lambda \otimes_{A_{m,\infty}} A_{m,\mathbf{n}} \end{array}$$

Since  $A_{m,\mathbf{n}}$  is defined by one equation, a simple computation yields the following

**Proposition 7.2.2.** *We have the following equivalence*

$$\Lambda \otimes_{A_{m,\infty}} A_{m,\mathbf{n}} \simeq \Lambda_0 \oplus \Lambda_{\mathbf{n}}[1 - 2d|\mathbf{n}|](-d|\mathbf{n}|).$$

*Proof.* Consider the following monoidal self-equivalence  $\vec{\text{sh}}$  of  $\text{Vect}^{\mathbb{Z}_{\geq 0}^m}$ , where

$$\vec{\text{sh}}(V)_{\mathbf{d}} = V_{\mathbf{d}}[-2d|\mathbf{d}|](-d|\mathbf{d}|),$$

and its inverse<sup>22</sup>

$$\overleftarrow{\text{sh}}(V)_{\mathbf{d}} = V_{\mathbf{d}}[2d|\mathbf{d}|](d|\mathbf{d}|).$$

Observe that  $\overleftarrow{\text{sh}}(A_{m,\infty})$  and  $\overleftarrow{\text{sh}}(A_{m,\mathbf{n}})$  are in the heart of the  $t$ -structure of  $\text{Vect}^{\mathbb{Z}_{\geq 0}^m}$ , and hence, we can compute the tensor using classical means. Namely, we have the following two-step resolution of  $\overleftarrow{\text{sh}}(A_{m,\mathbf{n}})$

$$0 \longrightarrow \Lambda[x_1, x_2, \dots, x_m] \longrightarrow \Lambda[x_1, x_2, \dots, x_m] \longrightarrow \Lambda[x_1, \dots, x_m] / (\cap_{k=q}^m x_k^{\mathbf{n}_k}) \simeq \overleftarrow{\text{sh}}(A_{m,\mathbf{n}}),$$

where the first non-zero term is shifted to graded degrees  $\mathbf{n}$ . This gives us

$$\Lambda \otimes_{\overleftarrow{\text{sh}}(A_{m,\infty})} \overleftarrow{\text{sh}}(A_{m,\mathbf{n}}) \simeq \Lambda_0 \oplus \Lambda_{\mathbf{n}}[1].$$

Sheering back using  $\vec{\text{sh}}$ , we get the desired result. □

<sup>22</sup>Here,  $\vec{\text{sh}}$  and  $\overleftarrow{\text{sh}}$  stand for sheering right and left respectively. Note also that the definitions of these functors depend on  $d = \dim X$  we fixed throughout.

7.2.3. The computation above shows that

$$\mathrm{add}_!(\Lambda \otimes_{A_{m,\infty}} A_{m,\mathbf{n}}) \in \mathrm{ComAlg}(\mathrm{Vect}^{\mathbb{Z}_{\geq 0}})$$

depends only on  $|\mathbf{n}|$ . We can amplify this fact, by taking factorization cohomology, to obtain a statement about the cohomology of  $Z_{\mathbf{n}}^m(X)$ .

**Theorem 7.2.4.** *The relative tensor*

$$\mathrm{add}_!(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)) \in \mathrm{ComAlg}^{\mathrm{un},\mathrm{au}}(\mathrm{Vect}^{\mathbb{Z}_{\geq 0,+}})$$

*depends only on  $|\mathbf{n}|$ .*

*Proof.* It suffices to show the statement for

$$\mathrm{add}_!(0 \sqcup_{\mathcal{A}_{m,\infty}(X)_+} \mathcal{A}_{m,\mathbf{n}}(X)_+).$$

But this is equivalent to

$$\begin{aligned} \mathrm{add}_!(0 \sqcup_{\pi_{?,*} A_{m,\infty}(X)_+} \pi_{?,*} A_{m,\mathbf{n}}(X)_+) &\simeq \mathrm{add}_!(\pi_{?,*}(0 \sqcup_{A_{m,\infty}(X)} A_{m,\mathbf{n}}(X))) \\ &\simeq \mathrm{add}_!(\pi_{?,*}(\Lambda \otimes_{A_{m,\infty}(X)} A_{m,\mathbf{n}}(X)))_+ \\ &\simeq \pi_{?,*}(\mathrm{add}_!(\Lambda \otimes_{A_{m,\infty}(X)} A_{m,\mathbf{n}}(X)))_+, \end{aligned}$$

where the last equivalence is due to the fact that  $\mathrm{add}_!$  is continuous and symmetric monoidal. Now, we are done, by Proposition 7.2.2.  $\square$

*Remark 7.2.5.* In §7.5, we will give an explicit map equivalence between the  $\mathrm{add}_!(\Lambda \otimes_{A_{m,\infty}} A_{m,\mathbf{n}})$ 's, and hence, also the  $\mathrm{add}_!(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X))$ 's for various  $m$  and  $\mathbf{n}$  with  $|\mathbf{n}|$  fixed. See also Remarks 7.5.7 and 7.5.8.

7.3. **coLie-coalgebras.** Via Koszul duality, there exist coLie-coalgebras  $\mathfrak{a}_{m,\mathbf{n}}$  and  $\mathfrak{a}_{m,\infty}$  in  $\mathrm{Vect}^{\mathbb{Z}_{\geq 0,+}^m}$  such that

$$\mathrm{coChev} \mathfrak{a}_{m,\mathbf{n}} \simeq (A_{m,\mathbf{n}})_+ \quad \text{and} \quad \mathrm{coChev} \mathfrak{a}_{m,\infty} \simeq (A_{m,\infty})_+,$$

or equivalently,

$$\mathrm{coChev}^{\mathrm{un}} \mathfrak{a}_{m,\mathbf{n}} \simeq A_{m,\mathbf{n}} \quad \text{and} \quad \mathrm{coChev}^{\mathrm{un}} \mathfrak{a}_{m,\infty} \simeq A_{m,\infty}.$$

7.3.1. Similar to the case of algebras, we use  $\mathfrak{a}_{m,\mathbf{n}}(X)$  and  $\mathfrak{a}_{m,\infty}(X)$  to denote  $\pi^! \mathfrak{a}_{m,\mathbf{n}}$  and  $\pi^! \mathfrak{a}_{m,\infty}$  respectively. From (6.2.2) and (6.3.8), we see that

$$\mathcal{A}_{m,\mathbf{n}}(X)_+ \simeq \mathrm{coChev}(\pi_*(\mathfrak{a}_{m,\mathbf{n}}(X))) \simeq \mathrm{coChev}(\pi_* \omega_X \otimes \mathfrak{a}_{m,\mathbf{n}})$$

and similarly, that

$$\mathcal{A}_{m,\infty}(X)_+ \simeq \mathrm{coChev}(\pi_* \omega_X \otimes \mathfrak{a}_{m,\infty}).$$

7.3.2. We will use  $\mathfrak{a}_{m,\mathbf{n}}(X)$  and  $\mathfrak{a}_{m,\infty}(X)$  to denote  $\pi_* \omega_X \otimes \mathfrak{a}_{m,\mathbf{n}}$  and  $\pi_* \omega_X \otimes \mathfrak{a}_{m,\infty}$  respectively. In particular, we have

$$\mathcal{A}_{m,\mathbf{n}}(X)_+ \simeq \mathrm{coChev} \mathfrak{a}_{m,\mathbf{n}}(X),$$

including the case where  $\mathbf{n} = \infty$ .

7.3.3. The goal now is to compute  $\mathfrak{a}_{m,\mathbf{n}}$  and  $\mathfrak{a}_{m,\infty}$ . Since  $A_{m,\infty}$  is a free commutative algebra, the corresponding coLie-coalgebra  $\mathfrak{a}_{m,\infty}$  is thus abelian (i.e. trivial co-multiplication map). Moreover, the underlying object in  $\mathrm{Vect}^{\mathbb{Z}_{\geq 0,+}^m}$  given by

$$\mathfrak{a}_{m,\infty} \simeq \bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}[-2d+1](-d).$$

Note that there is an extra cohomological shift to the left by 1 compared to (7.1.2). This is because for a trivial coLie-coalgebra  $L$ ,  $\mathrm{coChev} L \simeq (\mathrm{Sym} L[-1])_+$ .

**Lemma 7.3.4.** *The algebra  $(\Lambda \otimes_{A_{m,\infty}} A_{m,\mathbf{n}})_+$  is both a free and a trivial algebra. Hence,*

$$\begin{aligned} \text{coPrim}[1](\Lambda \otimes_{A_{m,\infty}} A_{m,\mathbf{n}})_+ &\simeq \text{coFree}_{\text{coLie}}(\Lambda_{\mathbf{n}}[2 - 2d|\mathbf{n}|](-d|\mathbf{n}|)) \\ &\simeq \text{cotriv}_{\text{coLie}}(\Lambda_{\mathbf{n}}[2 - 2d|\mathbf{n}|](-d|\mathbf{n}|)). \end{aligned}$$

*Proof.* The string of equivalences comes from the first assertion, since Koszul duality exchanges free objects and trivial objects. But now, the first assertion comes from the fact that a free commutative algebra generated by a one dimensional vector space living in an odd cohomological degree is trivial.  $\square$

**Corollary 7.3.5.** *We have the following equivalence*

$$(7.3.6) \quad \text{oblv}_{\text{coLie}} \mathfrak{a}_{m,\mathbf{n}} \simeq \bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}[-2d+1](-d) \oplus \Lambda_{\mathbf{n}}[2 - 2d|\mathbf{n}|](-d|\mathbf{n}|).$$

Moreover,  $\mathfrak{a}_{m,\mathbf{n}}$  is equipped with the structure of a  $d$ -shifted strongly unital (see Definition 6.5.3).

*Proof.* The pushout diagram (7.2.1) gives us the following pushout diagram in  $\text{Vect}^{\mathbb{Z}_{\geq 0,+}^m}$

$$\begin{array}{ccc} \bigoplus_{k=1}^m \Lambda_{\mathbf{1}_k}[-2d+1](-d) & \longrightarrow & \text{oblv}_{\text{coLie}} \mathfrak{a}_{m,\mathbf{n}} \\ \downarrow & & \downarrow \\ 0 & \longrightarrow & \Lambda_{\mathbf{n}}[2 - 2d|\mathbf{n}|](-d|\mathbf{n}|) \end{array}$$

By graded-degree consideration, we get the computation of  $\text{oblv}_{\text{coLie}} \mathfrak{a}_{m,\mathbf{n}}$ .

Also by degree consideration, we see that the only possible non-zero co-bracket of  $\mathfrak{a}_{m,\mathbf{n}}$  is given by

$$(\mathfrak{a}_{m,\mathbf{n}})_{\mathbf{n}} \rightarrow \bigotimes_{k=1}^m (\mathfrak{a}_{m,\mathbf{n}})_{\mathbf{1}_k}^{\otimes \mathbf{n}_k}.$$

This implies the second assertion, where the first summand of (7.3.6) gives the  $d$ -shifted strongly unital structure.<sup>23</sup>  $\square$

The computation above, coupled with Lemma 6.5.4, implies the following

**Corollary 7.3.7.** *We have*

$$\begin{aligned} &\text{oblv}_{\text{coLie}}(\mathfrak{a}_{m,\mathbf{n}}(X)) \\ &\simeq \left( \bigoplus_{k=1}^m C^*(X, \omega_X) \otimes \Lambda_{\mathbf{1}_k}[-2d+1](-d) \right) \oplus (C^*(X, \omega_X) \otimes \Lambda_{\mathbf{n}}[2 - 2d|\mathbf{n}|](-d|\mathbf{n}|)) \end{aligned}$$

and

$$\mathfrak{a}_{m,\infty}(X) \simeq \bigoplus_{k=1}^m (C^*(X, \omega_X) \otimes \Lambda_{\mathbf{1}_k}[-2d+1](-d)),$$

with trivial  $\text{coLie}$ -structure.

Moreover, both  $\text{coLie}$ -coalgebras are strongly unital.

**Corollary 7.3.8.** *We have the following equivalences*

$$\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X) \simeq \text{Sym}(C^*(X, \omega_X) \otimes \Lambda_{\mathbf{n}}[1 - 2d|\mathbf{n}|](-d|\mathbf{n}|)).$$

*Proof.* A priori,  $\text{coChev}^{\text{un}}$  rather than  $\text{Sym}$  should appear on the RHS of the equivalence above. However, the  $\text{coLie}$ -coalgebra under consideration is trivial, by Lemma 7.3.4, and we are done.  $\square$

**Remark 7.3.9.** Note that since  $|\mathbf{n}| \geq 2$  (by our convention),  $\text{oblv}_{\text{gr}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X))$  is  $\text{Frob}_q^{-1}$ -summable, since the expression inside  $\text{Sym}$  is finite dimensional has positive weight.

<sup>23</sup>See §7.7 for a similar discussion, where we dualize and deal with Lie-algebras rather than  $\text{coLie}$ -coalgebras, which is, psychologically, a bit easier to think about.

*Remark 7.3.10.* In the case of  $X = \mathbb{A}^d$ , the computation of Corollary 7.3.5 allows us to compute  $Z_{\mathbf{n}}^m(\mathbb{A}^d)$  easily, since  $C^*(X, \omega_X) \simeq \Lambda[2d](d)$  has trivial co-algebra structure in this case (see also Remark 6.3.9). Thus,

$$\mathcal{A}_{m,\mathbf{n}}(\mathbb{A}^d) \simeq \text{Sym} \left( \bigoplus_{k=1}^m \Lambda_{1_k} \oplus \Lambda_{\mathbf{n}}[1 - 2d|\mathbf{n}| + 2d](-d|\mathbf{n}| + d) \right).$$

**7.4. Homological stability.** The computation of  $a_{m,\mathbf{n}}$  above allows us to deduce homological stability for  $Z_{\mathbf{n}}^m(X)$ .

**Theorem 7.4.1.** *Let  $X$  be an irreducible scheme of dimension  $d \geq 1$ . For each  $1 \leq k \leq m$  and  $c \geq 0$ , there exists a natural map*

$$H^c(Z_{\mathbf{n}}^{\mathbf{d}}(X), \vec{\omega}_{Z_{\mathbf{n}}^{\mathbf{d}}(X)}) \rightarrow H^c(Z_{\mathbf{n}^{\mathbf{d}+1_k}}^{\mathbf{d}+1_k}(X), \vec{\omega}_{Z_{\mathbf{n}^{\mathbf{d}+1_k}}^{\mathbf{d}+1_k}(X)})$$

that is

- (i) an equivalence when  $\mathbf{d}_k \geq 2c$ , and injective when  $\mathbf{d}_k = 2c - 1$ , when  $d = 1, m = 1, n = 2$  (the case of configuration spaces of a curve),
- (ii) an equivalence when  $\mathbf{d}_k \geq c$ , and injective when  $\mathbf{d}_k = c - 1$  otherwise.

*Proof.* The result is now a direct consequence of Theorem 5.1.12, using the computation of Corollary 7.3.7.

In the first case (i.e. the case of configuration spaces of a curve), we can take  $s = 2, s_i = 0, \forall i$  in (5.1.13). For the other cases, with  $n$  finite, we can take  $s = 1, s_i = 0, \forall i$ .

The case where  $n = \infty$  is treated similarly, and in fact, simpler, since the coLie-coalgebra under consideration is trivial.  $\square$

**7.5. Stable homological densities.** Theorem 7.2.4 and Corollary 7.3.8 show that the quotients

$$\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)$$

depends only on  $|\mathbf{n}|$ . In this section, we will show how this implies the same statement about the stable homology. Namely, we want to show the following

**Theorem 7.5.1.** *Then, we have a natural equivalence*

$$\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X) \simeq \Lambda \otimes_{\overline{\mathcal{A}}_{1,\infty}(X)} \overline{\mathcal{A}}_{1,|\mathbf{n}|}(X).$$

*In particular, the relative tensor depends only on  $|\mathbf{n}|$ .*

**7.5.2.** Before proving the Proposition, we will need some preparation. Choosing a smooth point on  $X$ , and hence, a smooth point on  $Z_{\infty}^{\mathbf{d}}(X)$  for each  $\mathbf{d}$ , we get a map of algebras

$$\mathcal{A}_{m,\infty}(X) \rightarrow \Lambda[\mathbb{Z}_{\geq 0}^m],$$

which is compatible with the unit map

$$\Lambda[\mathbb{Z}_{\geq 0}^m] \rightarrow \mathcal{A}_{m,\infty}(X)$$

induced by the structure map  $Z_{\infty}^m(X) \rightarrow \text{pt}$ .

The rough idea now is to construct a natural equivalence

$$\Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X) \rightarrow \Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\mathcal{A}_{1,\infty}(X)} \mathcal{A}_{1,|\mathbf{n}|}(X).$$

The theorem is then obtained by taking

$$\text{colim} = \overline{(-)} : \text{Vect}^{\mathbb{Z}_{\geq 0}^m \rightarrow} \rightarrow \text{Vect}.$$

**7.5.3.** We will now make this idea rigorous by using §4.4. As in there, to keep the notation less cluttered, we will drop  $\text{add}_!$  from the notation. For example, whenever we have a map where the target is  $\mathbb{Z}_{\geq 0}$ -graded, and the source is  $\mathbb{Z}_{\geq 0}^m$ -graded,  $\text{add}_!$  is implicit in the source.



7.5.4. First, we have the following natural map of algebras

$$\Lambda[\mathbb{Z}_{\geq 0}^m] \rightarrow \Lambda[\mathbb{Z}_{\geq 0}]$$

by sending all generators of the LHS to the generator of the RHS. Similarly, we have the following maps of algebras

$$A_{m,\infty} \rightarrow A_{1,\infty}$$

and

$$A_{m,\mathbf{n}} \rightarrow A_{1,|\mathbf{n}|},$$

which, respectively, induce maps of algebras

$$\mathcal{A}_{m,\infty}(X) \rightarrow \mathcal{A}_{1,\infty}(X)$$

and

$$\mathcal{A}_{m,\mathbf{n}}(X) \rightarrow \mathcal{A}_{1,|\mathbf{n}|}(X).$$

These maps, in turn, induce a map between algebras

$$\Lambda[\mathbb{Z}_{\geq 0}^m] \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X) \rightarrow \Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\mathcal{A}_{1,\infty}(X)} \mathcal{A}_{1,|\mathbf{n}|}(X),$$

which factors through

$$(7.5.5) \quad \Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X) \rightarrow \Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\mathcal{A}_{1,\infty}(X)} \mathcal{A}_{1,|\mathbf{n}|}(X).$$

Since  $\Lambda \otimes_{\Lambda[\mathbb{Z}_{\geq 0}]}$  – of the map (7.5.5) is an equivalence, by Theorem 7.2.4, so is (7.5.5) itself, since  $\Lambda \otimes_{\Lambda[\mathbb{Z}_{\geq 0}]}$  – is the functor of taking associated graded, by §4.2.8, which is conservative, by Lemma 4.1.15. We thus obtain the following

**Proposition 7.5.6.** *The natural map of algebras*

$$\Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X) \rightarrow \Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\mathcal{A}_{1,\infty}(X)} \mathcal{A}_{1,|\mathbf{n}|}(X)$$

*is an equivalence.*

*Proof of Theorem 7.5.1.* We will now complete the proof of Theorem 7.5.1. Indeed, taking colimit along  $\mathbb{Z}_{\geq 0}^{\rightarrow}$ , keeping in mind that we have the following natural equivalences, we have

$$\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \overline{\mathcal{A}_{m,\mathbf{n}}(X)} \simeq \overline{\Lambda[\mathbb{Z}_{\geq 0}^m] \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)} \simeq \overline{\Lambda[\mathbb{Z}_{\geq 0}] \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)},$$

where the middle item is obtained by taking colimit along  $\mathbb{Z}_{\geq 0}^{m,\rightarrow}$  and the last one along  $\mathbb{Z}_{\geq 0}^{\rightarrow}$ . The first equivalence is due to the fact that  $\overline{(-)}$  along  $\mathbb{Z}_{\geq 0}^{m,\rightarrow}$  is, by Corollary 4.5.7, a left adjoint, and hence, commutes with pushouts of algebras. The second equivalence is due to Lemma 4.5.10.  $\square$

A couple of remarks are in order.

*Remark 7.5.7.* Observe that we have the following diagram

$$\begin{array}{ccc} A_{m,\infty} & \longrightarrow & A_{m,\mathbf{n}} \\ \downarrow & & \downarrow \\ A_{1,\infty} & \longrightarrow & A_{1,|\mathbf{n}|} \\ \downarrow & & \downarrow \\ \Lambda & \longrightarrow & \Lambda \otimes_{\mathcal{A}_{m,\infty}} A_{m,\mathbf{n}} \simeq \Lambda \otimes_{\mathcal{A}_{1,\infty}} A_{1,|\mathbf{n}|} \end{array}$$

where each square is a pushout square in  $\text{ComAlg}(\mathbb{Z}_{\geq 0})$ .<sup>24</sup> This provides us with a canonical map witnessing the coincidences observed in Proposition 7.2.2.

<sup>24</sup>One way to see that the squares are indeed pushouts is to use the same trick as in Proposition 7.2.2.

By tensoring (over  $\Lambda[\mathbb{Z}_{\geq 0}]$ ) the map in Proposition 7.5.6 with  $\Lambda$ , we get a canonical map witnessing the coincidences observed in Theorem 7.2.4.

*Remark 7.5.8.* Theorem 7.5.1 allow us to produce a natural equivalence

$$\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X) \simeq \Lambda \otimes_{\overline{\mathcal{A}}_{m',\infty}(X)} \overline{\mathcal{A}}_{m',\mathbf{n}'}$$

whenever  $|\mathbf{n}| = |\mathbf{n}'|$  by first passing through

$$\Lambda \otimes_{\overline{\mathcal{A}}_{1,\infty}(X)} \overline{\mathcal{A}}_{1,|\mathbf{n}|}(X).$$

We can link the cases of  $(m, \mathbf{n})$  and  $(m', \mathbf{n}')$  directly by considering a more general form of the functor  $\text{add}_!$ . Indeed, instead of only consider  $\text{add} : \mathbb{Z}_{\geq 0}^m \rightarrow \mathbb{Z}_{\geq 0}$ , one can consider, more generally, homomorphisms  $\text{add} : \mathbb{Z}_{\geq 0}^m \rightarrow \mathbb{Z}_{\geq 0}^{m'}$  in an analogous way. We can then define the functor  $\text{add}_!$  by left Kan extension as usual, and the whole argument goes through unchanged. We do not need this in the sequel.

*Remark 7.5.9.* The stable homological density

$$\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)$$

is in fact computable and has a simple expression. This gives an alternative proof of Theorem 7.5.1. Since it relies on  $L_\infty$ -algebras, we postpone the actual computation to Proposition 7.7.8.

**7.6. Decategorifications.** Using §5.3, we will now see how Theorem 7.2.4 and Theorem 7.5.1 immediately give us the coincidences observed in quotients of (graded) Euler characteristics, and when our scheme  $X$  comes from pulling back  $X_0$  over a finite field  $\mathbb{F}_q$ , also of virtual Poincaré series and  $L$ -series. Since Poincaré series are not additive, more care has to be taken. We will return to this in §7.7

As before, throughout this subsection, we will assume that  $X$  is an irreducible scheme of dimension  $d$ .

**Proposition 7.6.1.** *The quotient of graded Euler characteristics*

$$\frac{\chi^{\text{gr}}(\mathcal{A}_{m,\mathbf{n}}(X))}{\chi^{\text{gr}}(\mathcal{A}_{m,\infty}(X))} = \chi^{\text{gr}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)),$$

and hence, depends only  $|\mathbf{n}|$ .

When  $X$  is a pullback of a scheme  $X_0$  over  $\mathbb{F}_q$ , then all the objects inside has the action of the geometric Frobenius  $\text{Frob}$ . We have the same statement for

$$(7.6.2) \quad \frac{\chi_{\text{Frob}_q^{-1}}^{\text{gr}}(\mathcal{A}_{m,\mathbf{n}}(X), t)}{\chi_{\text{Frob}_q^{-1}}^{\text{gr}}(\mathcal{A}_{m,\infty}(X), t)} = \chi_{\text{Frob}_q^{-1}}^{\text{gr}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X), t) = \prod_{x \in |X|} (1 - q^{-d|\mathbf{n}| \deg x} t^{|\mathbf{n}| \deg x}).$$

*Proof.* The equality comes directly from Corollary 5.3.10. The fact that it depends only on  $|\mathbf{n}|$  is a consequence of Theorem 7.2.4. The case of  $\chi_F^{\text{gr}}$  is treated the same way (see also Remark 5.1.17).

The second equality involving  $\chi_{\text{Frob}_q^{-1}}^{\text{gr}}$  follows from Proposition 6.4.3 and Proposition 7.2.2, using the fact that

$$\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X) \simeq \pi_{? *}^{\text{un}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)).$$

□

Combined with Proposition 5.3.19, noting that

$$\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X) \simeq \Lambda \otimes_{\Lambda[\mathbb{Z}_{\geq 0}^m]} \Lambda[\mathbb{Z}_{\geq 0}^m] \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X),$$

which is also  $\text{Frob}_q^{-1}$ -summable (by the explicit computation at Corollary 7.3.8), we obtain the following

**Corollary 7.6.3.** *When  $X$  is as in the second part of Proposition 7.6.1. Then*

$$(7.6.4) \quad \chi_{\text{Frob}_q^{-1}}(\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)) = \chi_{\text{Frob}_q^{-1}}^{\text{gr}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X), 1) = \zeta_{X_0}(d|\mathbf{n}|)^{-1}.$$

*Remark 7.6.5.* When  $\mathbf{n} = (n, n, \dots, n)$ , evaluating  $t = 1$  and using the Grothendieck-Lefschetz trace formula, Theorem 6.4.4, we see that the left most term of (7.6.2) captures the arithmetic densities

$$\lim_{d \rightarrow \infty} \frac{|Z_{\mathbf{n}}^d(X_0)(\mathbb{F}_q)|}{|Z_{\infty}^d(X_0)(\mathbb{F}_q)|}$$

and the middle term becomes

$$\chi_{\text{Frob}_q}^{\text{gr}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X), 1) = \chi_{\text{Frob}_q}^{-1}(\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)) = \zeta(d|\mathbf{n}|) = \zeta(dmn)^{-1}.$$

Thus, equality (7.6.4) categorifies arithmetic density of Theorem 1.1.4.

7.6.6. Arguing similarly as Proposition 7.6.1, we get

**Proposition 7.6.7.** *When  $X$  is a pullback of a scheme  $X_0$  over  $\mathbb{F}_q$ , then we have*

$$\frac{\text{Poinc}^{\text{vir}}(\overline{\mathcal{A}}_{m,\mathbf{n}}(X))}{\text{Poinc}^{\text{vir}}(\overline{\mathcal{A}}_{m,\infty}(X))} = \text{Poinc}^{\text{vir}}(\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)),$$

and hence, only depends on  $|\mathbf{n}|$ .

*Proof.* We argue similarly as the Proposition above, but using Theorem 7.5.1 and Corollary 5.3.15 in place of Theorem 7.2.4 and Corollary 5.3.10.  $\square$

*Remark 7.6.8.* Note that all the terms appearing in Propositions 7.6.1 and 7.6.7 are readily available in explicit forms using Propositions 5.3.7 and 5.3.14.

7.7.  **$L_{\infty}$ -algebras, the case of Poincaré series, and stable homological densities.** As we have seen above, the various equivalences of algebras given in Theorem 7.2.4 and Theorem 7.5.1 allow us to systematically recover information about densities of Euler-characteristics and virtual Poincaré polynomials by decategorification. This comes from the fact that these invariants are additive, i.e. they behave nicely with respect to fiber sequences. Poincaré polynomials are, unfortunately, not additive, and only exhibit good behaviors when  $\text{coLie}$ -coalgebras involved are abelian. In the first part of this subsection, we will find the conditions on the dimension of  $X$  as well as the numbers  $m, \mathbf{n}$  so that  $\mathfrak{a}_{m,\mathbf{n}}(X)$  is abelian. In the second part, using the same technique, we obtain an expression for the stable homological density  $\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)$ .

Since it is more convenient to manipulate Lie-algebras than  $\text{coLie}$ -coalgebras, in this subsection, we will exclusively work with the linear dual  $\mathfrak{a}_{m,\mathbf{n}}(X)^{\vee}$ , which is now a Lie-algebra. We can take the double-dual to get back to the original  $\text{coLie}$ -coalgebra, since everything is finite dimensional.

We start with the following dual description of Corollary 7.3.7. We ignore Tate twists, since it is not relevant for the purposes of taking Poincaré polynomials.

**Lemma 7.7.1.** *We have*

$$\text{oblv}_{\text{Lie}}(\mathfrak{a}_{m,\mathbf{n}}(X)^{\vee}) \simeq \left( \bigoplus_{k=1}^m C_c^*(X) \otimes \Lambda_{\mathbf{1}_k}[2d-1] \right) \oplus (C_c^*(X) \otimes \Lambda_{\mathbf{n}}[-2+2d|\mathbf{n}|]).$$

Here, we use  $C_c^*(X)$  to denote  $C_c^*(X, \Lambda) \simeq C^*(X, \omega_X)^{\vee}$ , where the equivalence is due to Verdier duality. Note that  $C_c^*(X)$  lives in cohomological degrees  $[0, 2d]$  in general. When  $X$  is smooth affine,  $C_c^*(X)$  lives in cohomological degrees  $[d, 2d]$ .

7.7.2. By homotopy transfer ([LV12, §10.3]), one obtains the structure of a graded  $L_{\infty}$ -algebra on

$$\bigoplus_{k=1}^m H_c^*(X)[2d-1]_{\mathbf{1}_k} \oplus H_c^*(X)[-2+2d|\mathbf{n}|]_{\mathbf{n}}.$$

By degree consideration, the only possibly non-vanishing bracket is the  $|\mathbf{n}|$ -th bracket, which has cohomological degree  $2 - |\mathbf{n}|$

$$l_{|\mathbf{n}|} : \bigotimes_{k=1}^m (H_c^*(X)[2d-1]_{1_k})^{\otimes \mathbf{n}_k} \rightarrow H_c^*(X)[2d|\mathbf{n}| - |\mathbf{n}|]_{\mathbf{n}}$$

The goal now is to see when this bracket vanishes.

7.7.3. Suppose that  $H_c^*(X)$  lives in cohomological degrees  $[c, 2d]$  for some  $c \geq 0$ . We see that the lowest cohomological degree on the LHS is  $|\mathbf{n}|(-2d + c + 1)$  whereas the highest cohomological degree on the RHS is  $2d + |\mathbf{n}|(1 - 2d)$ . The bracket is forced to vanish when the former is greater than the latter, i.e. when

$$(7.7.4) \quad |\mathbf{n}|(-2d + c + 1) > 2d + |\mathbf{n}|(1 - 2d),$$

which is equivalent to

$$|\mathbf{n}|c > 2d.$$

This can only happen when  $c > 0$ , in which case, we have

$$|\mathbf{n}| > \frac{2d}{c}.$$

Coupled with Proposition 5.3.12 and Theorem 7.5.1, we have the following

**Proposition 7.7.5.** *Let  $X$  be an irreducible scheme of dimension  $d$  such that  $H_c^*(X)$  concentrates in cohomological degrees  $[c, 2d]$ , where  $c > 0$ . Then, for all  $m, \mathbf{n}$  such that  $|\mathbf{n}| > 2d/c$ , we have*

$$(7.7.6) \quad \frac{\text{Poinc}(\overline{\mathcal{A}}_{m,\mathbf{n}}(X))}{\text{Poinc}(\overline{\mathcal{A}}_{m,\infty}(X))} = \text{Poinc}(\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X)).$$

In particular, the quotient depends only on  $|\mathbf{n}|$ .

7.7.7. In general, we do not have equality (7.7.6). The RHS of (7.7.6), which should be thought of as the “real” density, is computable. The following result computes the categorified homological density at the limit.

**Proposition 7.7.8.** *We have the following equivalences*

$$\Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,\mathbf{n}}(X) \simeq \text{oblv}_{\text{gr}}(\text{Sym}(C^*(X, \omega_X) \otimes \Lambda_{\mathbf{n}}[1 - 2d|\mathbf{n}|](-d|\mathbf{n}|))) \simeq \text{oblv}_{\text{gr}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X)).$$

*Proof.* Consider the pushout in  $\text{coLie}(\text{Vect}^{\mathbb{Z}_{\geq 0, +}^m})$

$$\begin{array}{ccc} \mathfrak{a}_{m,\infty}(X) & \longrightarrow & \mathfrak{a}_{m,\mathbf{n}}(X) \\ \downarrow & & \downarrow \\ \bigoplus_{k=1}^m \Lambda_{1_k}[1] & \longrightarrow & \mathfrak{q} \end{array}$$

which corresponds to the quotient

$$\Lambda[\mathbb{Z}_{\geq 0}^m] \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,\mathbf{n}}(X) = \text{coChev}^{\text{un}} \mathfrak{q}.$$

The coincidence here is that  $\mathfrak{q}$  is abelian. Indeed,

$$\text{oblv}_{\text{Lie}}(\mathfrak{q}^{\vee}) \simeq \bigoplus_{k=1}^m \Lambda_{1_k}[-1] \oplus (C_c^*(X) \otimes \Lambda_{\mathbf{n}}[-2 + 2d|\mathbf{n}|]),$$

and the inequality corresponding to (7.7.4) is

$$|\mathbf{n}| > 2d + |\mathbf{n}|(1 - 2d) \Leftrightarrow 0 > 2d(1 - |\mathbf{n}|)$$

which always happens since  $|\mathbf{n}| \geq 2$ .

But now, we have

$$\begin{aligned}
 \Lambda \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,n}(X) &\simeq \overline{\Lambda[\mathbb{Z}_{\geq 0}^m]} \otimes_{\overline{\mathcal{A}}_{m,\infty}(X)} \overline{\mathcal{A}}_{m,n}(X) \\
 &\simeq \overline{\Lambda[\mathbb{Z}_{\geq 0}^m]} \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,n}(X) \\
 \text{(Proposition 5.2.2)} \quad &\simeq \text{oblv}_{\text{gr}}(\text{coChev}^{\text{un}}(\text{Quot}_{\text{un}} \mathfrak{q})) \\
 \text{(by (5.3.17))} \quad &\simeq \text{oblv}_{\text{gr}}(\text{coChev}^{\text{un}}(0 \sqcup_{\mathfrak{a}_{m,\infty}(X)} \mathfrak{a}_{m,n}(X))) \\
 &\simeq \text{oblv}_{\text{gr}}(\Lambda \otimes_{\mathcal{A}_{m,\infty}(X)} \mathcal{A}_{m,n}(X)) \\
 \text{(Corollary 7.3.8)} \quad &\simeq \text{oblv}_{\text{gr}}(\text{Sym}(C^*(X, \omega_X) \otimes \Lambda_{\mathbf{n}}[1 - 2d|\mathbf{n}|](-d|\mathbf{n}|))).
 \end{aligned}$$

The second equivalence is the content of Corollary 7.3.8.  $\square$

#### ACKNOWLEDGMENTS

This paper owes an obvious intellectual debt to the illuminating treatments of factorization homology by J. Francis, D. Gaitsgory, and J. Lurie in [GR17, Gai15a, FG11]. I would like to thank B. Farb and J. Wolfson for bringing the question of explaining coincidences in homological densities to your attention. Moreover, I thank J. Wolfson for many helpful conversations on the subject.

Our work is supported by Advanced Grant “Arithmetic and Physics of Higgs moduli spaces” No. 320593 of the European Research Council. I thank T. Hausel and his group at IST Austria for useful discussions and for providing a wonderful research environment.

#### REFERENCES

- [FG11] J. Francis and D. Gaitsgory, *Chiral Koszul duality*, *Selecta Mathematica* **18** (September 2011), no. 1, 27–87 (en). [↑9, 10, 11, 16, 44, 65](#)
- [FWW16] B. Farb, J. Wolfson, and M. M. Wood, *Coincidences of homological densities, predicted by arithmetic* (November 2016), arXiv:1611.04563 [math]. [↑1, 3, 4, 8](#)
- [Gai15a] D. Gaitsgory, *The Atiyah-Bott formula for the cohomology of the moduli space of bundles on a curve* (May 2015), arXiv:1505.02331 [math]. [↑13, 16, 44, 65](#)
- [Gai15b] Dennis Gaitsgory, *Eisenstein series and quantum groups* (May 2015), arXiv:1505.02329 [math]. [↑44](#)
- [Gai17] ———, *The semi-infinite intersection cohomology sheaf-II: The Ran space version* (August 2017), arXiv:1708.07205 [math]. [↑46](#)
- [GK94] V. Ginzburg and M. Kapranov, *Koszul duality for operads*, *Duke Mathematical Journal* **76** (1994), no. 1, 203–272. [↑9](#)
- [GL14] D. Gaitsgory and J. Lurie, *Weil’s conjecture for function fields* (September 2014). [↑6, 13, 16, 40, 41, 43, 44, 46, 47, 49, 52](#)
- [GR17] D. Gaitsgory and N. Rozenblyum, *A study in derived algebraic geometry*, 1st ed., *Mathematical Surveys and Monographs*, American Mathematical Society, 2017 (English). [↑9, 17, 18, 20, 22, 23, 25, 41, 49, 65](#)
- [Ho16] Q. P. Ho, *The Atiyah-Bott formula and connectivity in chiral Koszul duality* (October 2016), arXiv:1610.00212 [math]. [↑17](#)
- [Ho17] ———, *Free factorization algebras and homology of configuration spaces in algebraic geometry*, *Selecta Mathematica* (July 2017), 1–53 (en). [↑5, 15, 16, 49, 52](#)
- [KM16] A. Kupers and J. Miller, *Homological stability for topological chiral homology of completions*, *Advances in Mathematics* **292** (April 2016), 755–827. [↑4](#)
- [KM18] ———,  *$E_n$ -cell attachments and a local-to-global principle for homological stability*, *Mathematische Annalen* **370** (February 2018), no. 1-2, 209–269 (en). [↑4, 5, 7, 52](#)
- [Knu14] B. Knudsen, *Betti numbers and stability for configuration spaces via factorization homology* (May 2014), arXiv:1405.6696 [math]. [↑5, 34, 57](#)
- [Lur17a] J. Lurie, *Higher algebra*, 2017. [↑9, 24](#)
- [Lur17b] Jacob Lurie, *Higher topos theory*, *Annals of mathematics studies*, Princeton University Press, Princeton, N.J., 2017. OCLC: ocn244702012. [↑9](#)
- [LV12] Jean-Louis Loday and Bruno Vallette, *Algebraic operads*, Vol. 346, Springer Science & Business Media, 2012. [↑9, 63](#)
- [LZ12] Y. Liu and W. Zheng, *Enhanced six operations and base change theorem for Artin stacks* (November 2012), arXiv:1211.5948 [math]. [↑13](#)
- [LZ14] ———, *Enhanced adic formalism and perverse t-structures for higher artin stacks* (April 2014), arXiv:1404.1128 [math]. [↑13](#)
- [May72] J. P. May, *The geometry of iterated loop spaces*, Springer Berlin, 1972. [↑4](#)
- [MS76] D. McDuff and G. Segal, *Homology fibrations and the “group-completion” theorem*, *Inventiones Mathematicae* **31** (October 1976), no. 3, 279–284 (en). [↑4, 7, 31](#)
- [Qui69] D. Quillen, *Rational homotopy theory*, *Annals of Mathematics* **90** (September 1969), no. 2, 205–295. [↑9](#)
- [Ras15] S. Raskin, *Chiral categories* (January 2015). [↑52](#)

- [Seg73] G. Segal, *Configuration-spaces and iterated loop-spaces*, *Inventiones mathematicae* **21** (1973), no. 3, 213–221. ↑[4](#)
- [VW15] R. Vakil and M. Wood, *Discriminants in the Grothendieck ring*, *Duke Mathematical Journal* **164** (2015), no. 6, 1139–1185. ↑[1](#), [4](#), [8](#)

INSTITUTE OF SCIENCE AND TECHNOLOGY AUSTRIA, KLOSTERNEUBURG, AUSTRIA  
Email address: qho@ist.ac.at