

# 1 Recap

We have the following locuses

$$C_d^r(C) = \left\{ D \in C^{(d)} \mid h^0(D) \geq r + 1 \right\}$$

$$W_d^r(C) = \left\{ L \in \text{Pic}^d(C) \mid h^0(L) \geq r + 1 \right\}$$

**Theorem 1.1** (Brill-Noether). *Let  $C$  be a curve of genus  $g$ , and let  $d \geq 1, r \geq 0$  be integers. Consider the Brill-Noether number:*

$$\rho = g - (r + 1)(g - d + r)$$

*If  $\rho \geq 0$  then  $W_d^r(C)$  is non empty. Now if  $C$  is a general curve, and  $\rho < 0$ , then  $W_d^r(C)$  is empty.*

Henry had also told us that there is always a  $D = g_6^2$  on a smooth curve of genus 6. If  $D$  has base points, then we have the following cases:

- If  $D$  has 2 base points, then (mapping using  $D$  is the same as mapping using  $D$  minus base points) it gives a map  $\phi_D : C \rightarrow \mathbb{P}^2$  which is either a birational map to a quadric, in which case the genus doesn't match, a 2-1 map to a conic, in which case  $C$  is hyperelliptic, or a 4-1 map to a line. The last case doesn't happen because  $\phi_D(C)$  is nondegenerate.
- If  $D$  has 1 base point, then  $\phi_D$  embeds  $C$  as a smooth quintic curve (5-1 map can't happen because nondegeneracy).
- Can use a similar argument to rule out the case of  $\geq 3$  base points.

If  $D$  has no base point, then  $6 = \deg \phi_D \cdot \deg \phi_D(C)$ , and we have the following cases:

- $\phi_D$  maps  $C$  in a 3-1 manner onto a conic. Here  $C$  is trigonal.
- $\phi_D$  maps  $C$  in a 2-1 manner onto a smooth plane cubic. Here  $C$  is bi-elliptic.
- $\phi_D$  maps  $C$  in a 2-1 manner onto a singular plane cubic. Here  $C$  is hyperelliptic.
- $\phi_D$  maps  $C$  birationally to a plane sextic curve  $C_0$ . In this case  $C_0$  cannot have a point of multiplicity  $\geq 4$  (genus drops too much). If  $C_0$  has a triple point, then  $C$  is trigonal.

**Note 1.1.** The genus formula for a plane curve  $C \subset \mathbb{P}^2$  of degree  $d$  with singularities of multiplicities  $m_i$  is

$$g = \binom{d-1}{2} - \sum \binom{m_i}{2}$$

## 2 General curve of genus 6

We want to know what a general curve of genus 6 looks like. Brill-Noether gives us this for free, but for genus 6, we can do some hands on inspections instead. Recall we have the moduli  $\mathcal{M}_g$ , and the Hurwitz space:

$$\mathcal{H}_{d,g} = \left\{ (C, f) \mid C \in \mathcal{M}_g, f : C \rightarrow \mathbb{P}^1 \text{ a simply branched cover of degree } d \right\}$$

which has dimension  $2d + 2g - 2$  by Riemann-Hurwitz (once we know the branched points, monodromy, which is finite, gives the cover). We have the projection map  $\pi : \mathcal{H}_{d,g} \rightarrow \mathcal{M}_g$ . Say  $C \in \text{im } \pi$ , i.e.,  $C$  has a  $g_d^1$ , then we expect

$$\dim \pi^{-1}(C) \geq \dim \text{PGL}(2) + \dim W_d^r(C) = 3 + \dim W_d^r(C)$$

where  $W_d^r(C)$  is the locus of line bundles  $\mathcal{L}$  of degree  $d$  with  $h^0(\mathcal{L}) \geq r + 1$ . In our case,  $g = 6$ , and for  $d = 2, 3$ , we have

$$\dim \mathcal{H}_{d,g} = 2d + 10 < 18 = \dim \mathcal{M}_g + 3$$

so a general curve of genus 6 cannot have a  $g_2^1$  or  $g_3^1$ . For  $d = 4$ , we have that a general curve of genus 6 can only have finitely many  $g_4^1$ .

If  $C$  is bi-elliptic or a smooth plane quintic, then  $C$  has at least 1-dimension worth of  $g_4^1$  (since we get a  $g_2^1$  from any point on elliptic curve), so  $C$  cannot be general either. It follows that a general curve of genus 6 is birational to a plane sextic with 4 simple nodes.

Our goal is to investigate the locus  $W_4^1(C)$  of  $g_4^1$  on a general genus 6 curve. Such a divisor looks like  $D = q_1 + q_2 + q_3 + q_4$ ; we can always assume these are distinct points. For this to be a  $g_4^1$ , we need  $h^0(D) \geq 2$ . Recall Riemann-Roch

$$h^0(D) - h^0(K_C - D) = 4 - 6 + 1$$

so we want  $h^0(K_C - D) \geq 3$ . But  $h^0(K_C) = 6$ , so  $D$  moves in a pencil iff it fails to impose independent conditions on  $K_C$ .

Back to our plane model, let's first consider  $C_0 \subset \mathbb{P}^2$  with 4 nodes  $\{p_1, p_2, p_3, p_4\}$ , no 3 of which are collinear. We can blow up these 4 points to get the normalization  $\widetilde{C}_0$ , and by universal property  $C$  is isomorphic to  $\widetilde{C}_0$ .

$$\begin{array}{ccc} C & \longrightarrow & S = \text{Bl}_{p_1, \dots, p_4} \mathbb{P}^2 \\ \downarrow & & \downarrow \pi \\ C_0 & \longrightarrow & \mathbb{P}^2 \end{array}$$

Here  $S$  is a quintic del Pezzo surface. Let  $\text{Pic}(S) = \langle H, E_1, E_2, E_3, E_4 \rangle$  where  $H$  is the pullback of a line in  $\mathbb{P}^2$ , and  $E_i$  are exceptional divisors. We have

$$E_i^2 = -1, \quad H^2 = 1, \quad H \cdot E_i = 0, \quad E_i \cdot E_j = 0$$

and

$$K_S = -3H + E_1 + E_2 + E_3 + E_4$$

$$C = \pi^* C_0 - \sum_1^4 \text{mult}(C, p_i) \cdot E_i = 6H - 2 \sum_1^4 E_i$$

$$K_C = (K_S + C)|_C = 3H - \sum_1^4 E_i$$

**Note 2.1.** Notice that  $K_C = -K_S|_C$ , so this shows that if we embeds

$$S \xrightarrow{-K_S} \mathbb{P}^5$$

then the image of  $C$  coincides with  $\phi_K(C)$  of the canonical embedding.

In other words, the canonical divisor  $K_C$  is cut out by cubics through 4 points  $\Gamma$ . Then  $D = \sum_1^4 q_i$  failing to impose independent conditions on  $K_C$  is equivalent to  $\Gamma = \{p_1, \dots, p_4, q_1, \dots, q_4\}$  failing to impose independent conditions on cubics.

**Proposition 2.1.** *Let  $\Gamma$  be a set of 8 points in  $\mathbb{P}^2$ , then  $\Gamma$  fails to impose independent conditions on cubics iff*

- $\Gamma$  contains 5 collinear points, or
- $\Gamma$  is contained in a conic.

Applying the proposition, we get that either  $\Gamma$  is contained in a conic, or 5 of them are collinear. By our choice, no 3 of  $\{p_i\}$  are collinear, so either  $\{p_1, p_2, q_1, q_2, q_3\}$  are collinear or  $\{p_1, q_i\}$ . In the first case, let  $l$  be the line, then

$$l \cap C_0 = \{p_1, p_2, q_1, q_2, q_3\}$$

so the intersection number is 7, which contradicts that  $C_0$  is a sextic. It follows that a  $g_4^1$  is cut out by either conics through  $\{p_i\}$  or lines through each  $p_i$ . That gives us all 5  $g_4^1$  on a general curve of genus 6.

### 3 Scheme structure on $W_4^1(C)$

Now let  $p_1, p_2, p_3$  be collinear. We can still only have the previous 2 cases because of intersection number. A  $g_4^1$  is, once again, cut out by conics through  $\{p_i\}$  or lines through each  $p_i$ . There is an issue here, which is that each conic through  $\{p_i\}$  contains a line through  $p_4$ , and vice versa. As a result we only have four  $g_4^1$ .

This has to do with the scheme structure on  $W_4^1(C)$ . In the previous case of 4 general points,  $W_4^1(C)$  has 5 points each of which is reduced. In this collinear case, we have 4 points but the one corresponding to conics is nonreduced. The usual way of checking whether a point is smooth is by looking at the tangent space.

We have  $\text{Pic}^d(C) \simeq \text{Pic}^0(C) \simeq J(C)$ , and we know the tangent space to  $J(C)$ , thus for any  $L \in \text{Pic}^d(C)$  we have

$$T_L \text{Pic}^d(C) \simeq H^1(C, \mathcal{O}_C)$$

and on the other hand we have the identification

$$H^1(C, \mathcal{O}_C) \simeq \left\{ \mathcal{L} \in \text{Pic} \left( \text{Spec} \frac{\mathbb{C}[\epsilon]}{(\epsilon^2)} \times C \right) \middle| \mathcal{L}|_C \simeq L \right\}$$

$$\{\phi_{\alpha\beta}\} \leftrightarrow \tau_{\alpha\beta}(1 + \epsilon\phi_{\alpha\beta})$$

which is the space of first order deformations of  $L$ . Then if  $L \in W_d^r(C) \setminus W_d^{r+1}(C)$ , i.e.,  $r(L) = r$ , we have

$$T_L W_d^r(C) \simeq \left\{ \mathcal{L} \in \text{Pic} \left( \text{Spec} \frac{\mathbb{C}[\epsilon]}{(\epsilon^2)} \times C \right) \middle| \mathcal{L}|_C \simeq L, \text{ every } \sigma \in H^0(L) \text{ extends to } H^0(\mathcal{L}) \right\}$$

Let  $v \in H^1(C, \mathcal{O}_C)$  corresponding to some  $\mathcal{L}$ , one can check that a section  $\sigma \in H^0(L)$  extends to a section of  $\mathcal{L}$  iff

$$H^1(C, \mathcal{O}_C) \otimes H^0(C, L) \rightarrow H^1(C, L)$$

$$v \smile \sigma \mapsto 0$$

so all sections extend iff  $v \smile H^0(C, L) = 0$ . By Serre Duality we have

$$H^0(C, K_C)^\vee \otimes H^0(C, L) \rightarrow H^0(C, K_C \otimes L^\vee)^\vee$$

and dualizing again to get multiplication map

$$H^0(C, K_C \otimes L^\vee) \otimes H^0(C, L) \xrightarrow{\mu} H^0(C, K_C)$$

and we can identify  $T_L W_d^r(C) \simeq \text{Ann}(\text{im } \mu)$ .

Back to our case of interest,  $W_4^1(C)$  is 0 dimensional. Let  $D$  be the  $g_4^1$  cut out by conics through  $\{p_i\}$ , then  $D = 2H - \sum E_i$  then look at the multiplication map

$$H^0(C, H) \otimes H^0 \left( C, 2H - \sum E_i \right) \rightarrow H^0 \left( C, 3H - \sum E_i \right)$$

If  $p_1, p_2, p_3$  are collinear, on a line  $l$ , then any conic through those  $p_i$  has to contain  $l$ . On the other hand, cubics through 4 points  $p_i$  don't have to contain  $l$  (take product of 3 lines  $\overline{p_4 p_i}$ ), so this map cannot be surjective. It follows that  $\text{Ann}(\text{im } \mu)$  has positive dimension, and  $D$  is not a smooth point.

We claimed that  $D$  is reduced if  $\{p_i\}$  are in general position. This seemingly implies that the map

$$H^0(C, H) \otimes H^0 \left( C, 2H - \sum E_i \right) \rightarrow H^0 \left( C, 3H - \sum E_i \right)$$

is surjective, i.e., any cubic through 4 general points can be written as a linear combination of reducible cubics, which is quite surprising. This is actually just a straightforward consequence of  $AF + BG$  theorem; here the key point is that 4 general points is a complete intersection of 2 conics.

**Note 3.1.** The version we are using is that for  $f, g \in \mathbb{C}[x, y, z]$  two homogenous polynomials and  $V(f) \cap V(g) = \Gamma$  transversely, i.e., in a finite number of reduced points, then any curve  $V(h)$  containing  $\Gamma$  has  $h \in (f, g)$ . This generalizes to transverse intersection in higher dimensions as well.