

RANK TWO COHOMOLOGY JUMP LOCI ON PROJECTIVE VARIETIES

QUOC ANH TRAN

ABSTRACT. Let X be a smooth connected complex projective variety. We classify the irreducible components of the rank two cohomology jump loci on X .

CONTENTS

1. Introduction	2
1.1. Hodge structures on moduli spaces	2
1.2. Classification of jump components	2
2. Preliminaries	4
2.1. Jump ideals	4
2.2. Variations of complex mixed Hodge and twistor structures	5
2.3. Variations with artinian coefficients	7
2.4. Absolute subset and rank one theory	8
2.5. Determinant sheet	9
3. Hodge theory of R_B and M_B	11
3.1. Betti universal deformation algebra and Deligne-Hitchin space	11
3.2. Jump loci are Hodge subvarieties	12
3.3. Hodge structures on good moduli image	16
3.4. Hodge structures on normalization	18
4. Virtually abelian jump components	21
4.1. Generic projective type	21
4.2. Reducible jump component	22
4.3. Dihedral jump component	24
4.4. Projectively finite and rigid jump components	25
5. Nonabelian jump component	27
5.1. Local systems on finite quotient curves	27
5.2. A factorization theorem following Corlette-Simpson	29
5.3. $C \rightarrow T$ is either a product or generically finite	33
5.4. Finiteness of $C \rightarrow T$	37
5.5. Étaleness of $C^\nu \rightarrow T$	40
References	42

1. INTRODUCTION

Let X be a smooth projective variety, and

$$\begin{aligned} R_B(X, x, r) &:= \operatorname{Hom}(\pi_1(X, x), \operatorname{GL}_r) \\ \mathcal{M}_B(X, r) &:= [R_B(X, x, r)/\operatorname{GL}_r] \\ M_B(X, r) &:= R_B(X, x, r)//\operatorname{GL}_r \end{aligned}$$

be, respectively, the moduli of framed local systems, the quotient stack, and the GIT quotient/good moduli space for the stack. We have the cohomology jump loci

$$\Sigma_k^i(X, x, r) := \left\{ \mathbb{V} \in R_B(X, x, r) \mid \dim H^i(X, \mathbb{V}) \geq k \right\}$$

which is GL_r -invariant hence descends to a substack $\Sigma_k^i(X, r) \subset \mathcal{M}_B(X, r)$. Denote $\overline{\Sigma}_k^i(X, r)$ its reduced image in $M_B(X, r)$.

1.1. Hodge structures on moduli spaces.

Theorem 1.1 (Theorem 3.9). *For smooth projective X , the jump locus $\Sigma_k^i(X, x, r) \subseteq R_B(X, x, r)$ is a Hodge subvariety. Furthermore, it is closed under semisimplification, and $\Sigma_k^i(X, r)_{\text{red}}$ is a Hodge subvariety which is absolute $\mathbb{Q}$ -closed.*

Every irreducible component of $\Sigma_k^i(X, r)_{\text{red}}$ is a Hodge subvariety and is absolute $\overline{\mathbb{Q}}$ -closed.

In particular, at a VHS point $\mathbb{V} \in \Sigma_k^i(X, x, r)$ there is a pro-MHS-algebra structure on $\hat{\mathcal{O}}_{\Sigma_k^i(X, x, r), \mathbb{V}}$. This persists for $\overline{\Sigma}_k^i(X, r) \subseteq M_B(X, r)$ by Proposition 3.11, and also on normalizations by Proposition 3.16.

1.2. Classification of jump components. Let $A = M_B(X, 1)$.

Proposition 1.2 (Proposition 4.3). *Let $Z \subseteq \overline{\Sigma}_k^i(X, r)$ be an irreducible component. Consider a general point $\rho \in Z$, with $\bar{\rho} : \pi_1(X, x, r) \rightarrow \operatorname{PGL}_2(\mathbb{C})$ the projective quotient. Then ρ is exactly one of*

- (A): ρ is reducible, i.e., $\rho \simeq \chi_1 \oplus \chi_2$;
- (B): ρ is irreducible, and, up to conjugation, $G_{\bar{\rho}} \subseteq D_{\infty}$;
- (C): $G_{\bar{\rho}} \simeq \bar{\rho}(\pi_1(X, x)) \simeq A_4, S_4$ or A_5 ;
- (D): $\bar{\rho}$ has Zariski dense image and is rigid;
- (E): $\bar{\rho}$ has Zariski dense image and is non-rigid.

For obvious reason, we call the first 4 cases the virtually abelian jump components, and the last one the nonabelian component. We have a structure theorem for the virtually abelian components:

Theorem 1.3.

- (A): (Proposition 4.5) either $Z \simeq \operatorname{Sym}^2(N)$ or $Z^{\nu} \simeq N_1 \times N_2$ with $N, N_1, N_2 \subseteq A$ translated subtori of A ;
- (B): (Proposition 4.7) either $Z^{\nu} \simeq N$ or $Z \simeq N/\langle \tau^* \rangle$ with N a translated subtorus of A and τ^* an involution on N ;
- (C): (Proposition 4.9) Z^{ν} is an algebraic torus;

(D): ([Proposition 4.10](#)) Z^ν is an algebraic torus.

For the nonabelian jump component, there is a torsion lift $\epsilon \in A$ of some $\bar{\epsilon} \in A/A^2$ such that $Z \subseteq M_B(X, 2, \det \in \epsilon A^2)$. Consider the diagram:

$$\begin{array}{ccc} M_B(X, 2, \det = \epsilon) \times A & \xrightarrow[\Phi_\epsilon]{(\mathbb{V}, \mathbb{L}) \mapsto \mathbb{V} \otimes \mathbb{L}} & M_B(X, 2, \det \in \epsilon A^2) \\ \text{pr}_2 \downarrow & & \downarrow \det \\ A & \xrightarrow{\mathbb{L} \mapsto \epsilon \otimes \mathbb{L}^{\otimes 2}} & A \end{array}$$

and let C be an irreducible component of $\Phi_\epsilon^{-1}(Z)$. By [Proposition 2.19](#), $Z^\nu \simeq C^\nu/H_C$ is an étale quotient of C^ν . Denote

$$\begin{array}{ccc} C \subseteq \Phi_\epsilon^{-1}(Z) \subseteq M_B(X, 2, \det = \epsilon) \times A & & \\ \text{pr}_1 \downarrow & \searrow \text{pr}_2 & \\ M_B(X, 2, \det = \epsilon) & & A \end{array}$$

$$D = \overline{\text{pr}_1(C)}_{\text{red}} \subseteq M_B(X, 2, \det = \epsilon), \quad T = \overline{\text{pr}_2(C)}_{\text{red}} \subseteq A$$

For more details, see [Section 2.5](#). As a consequence of [\[CS08\]](#) we have a factorization of D through a DM-curve:

Proposition 1.4 ([Proposition 5.8](#)). *There is a proper hyperbolic DM curve $\mathcal{Y} = [Y/G]$ and a morphism $s : X \rightarrow \mathcal{Y}$ and an irreducible component Q of a fixed-determinant, fixed-inertia locus $M_B(\mathcal{Y}, 2, \det = \delta, \tau)$ such that*

- (1) $s_* : \pi_1(X, x) \rightarrow \pi_1(\mathcal{Y}, \bar{y})$ is a surjection.
- (2) with the closed embedding $s^* : M_B(\mathcal{Y}, 2) \hookrightarrow M_B(X, 2)$, $D \subseteq s^*(Q)$.

Hence we can view C as a subvariety of $Q \times T \subseteq M_B(\mathcal{Y}, 2) \times A$.

Theorem 1.5 ([Proposition 5.18](#), [Proposition 5.23](#), [Proposition 5.26](#)). *Either $C = Q \times T$, $C = \{\mathbb{V}\} \times T$, or $\text{pr}_2 : C \rightarrow T$ is finite and surjective. In the latter case, the normalization $\text{pr}_2^\nu : C^\nu \rightarrow T$ is étale, and C^ν is an algebraic torus.*

Corollary 1.6. *If the general point of Z is of type (E), then either Z is an algebraic torus, Z^ν is an algebraic torus, or Z^ν is an étale quotient of $Q^\nu \times T$.*

2. PRELIMINARIES

We work with schemes over $\mathbb{C}$. For a scheme S , let S_{red} be the reduced scheme, and S^ν be the normalization of S .

2.1. Jump ideals. Let A be a noetherian ring, and $K^\bullet$ a bounded above complex of A -modules such that $H^i(K^\bullet)$ is a finitely generated A -module for all i . By [Har77, Lemma III.12.3], there exists a quasi-isomorphism $\phi : L^\bullet \rightarrow K^\bullet$ where $(L^\bullet, d^\bullet)$ is a bounded above complex of finitely generated free A -modules.

Definition 2.1. The cohomological jump ideals of $K^\bullet$ are defined to be

$$J_k^i(K^\bullet) := I_{\text{rank}_A(L^i) - k + 1}(d^{i-1} \oplus d^i) \subset A$$

where I denotes the determinantal ideal generated by $(\text{rank}_A(L^i) - k + 1)^2$ -minors. This definition does not depend on the choice of $L^\bullet$ [BW15, Definition-Proposition 2.2]. We also have that, for a base change $A \rightarrow B$,

$$J_k^i(K^\bullet)B = J_k^i(K^\bullet \otimes_A^L B) \quad (2.1)$$

$$V(J_k^i(K^\bullet)) = \{s \in \text{Spec } A \mid \dim_{\kappa(s)} H^i(K^\bullet \otimes_A^L \kappa(s)) \geq k\} \quad (2.2)$$

Let $S = \text{Spec } A \rightarrow R_B(X, x, r) = \text{Hom}(\pi_1(X, x), \text{GL}_r)$ be a classifying map. This is equivalent to a representation

$$\rho_S : \pi_1(X, x) \rightarrow \text{GL}_r(A)$$

which we might think of as a $A[\pi_1(X, x)]$ -module V_S . Let the corresponding local system of free A -modules on X be denoted $\mathbb{V}_S$. Here we have an explicit free model for $\text{R}\Gamma(X, \mathbb{V}_S)$. Fix a finite CW complex structure for X , we can form the cellular complex $C_\bullet(\tilde{X}, \mathbb{Z})$ on the universal cover $\tilde{X} \rightarrow X$; this is a complex of free $\mathbb{Z}[\pi_1(X, x)]$ -modules. Then

$$C^\bullet(X, \mathbb{V}_S) := \text{Hom}_{\mathbb{Z}[\pi_1(X, x)]}(C_\bullet(\tilde{X}, \mathbb{Z}), V_S)$$

is a complex of free A -modules. Cellular cohomology with local system coefficients identifies this with $\text{R}\Gamma(X, \mathbb{V}_S)$. Furthermore, for $f : \text{Spec } B \rightarrow S$, we have that

$$C^\bullet(X, \mathbb{V}_S \otimes_{A_X} B_X) = C^\bullet(X, \mathbb{V}_S) \otimes_A B \quad (2.3)$$

with A_X, B_X being the constant local systems on X . Recall that $R_B(X, x, r)$ is affine, and denote the universal local system, which is a local system of free $\Gamma(R_B(X, x, r), \mathcal{O}_{R_B(X, x, r)})$ -modules on X , to be $\mathbb{V}^{\text{univ}}$ (which is just $\mathbb{V}_S$ for $S = R_B(X, x, r)$). We can then give a scheme structure to

$$\Sigma_k^i(X, r) = \{\mathbb{V} \in R_B(X, x, r) \mid \dim H^i(X, \mathbb{V}) \geq k\}$$

as $\Sigma_k^i(X, r) = V(J_k^i(\text{R}\Gamma(X, \mathbb{V}^{\text{univ}})))$.

2.2. Variations of complex mixed Hodge and twistor structures.

Definition 2.2. A complex pure Hodge structure (C-HS) of weight n is a finite-dimensional $\mathbb{C}$ -vector space V with n -opposed decreasing filtrations $F^\bullet, F'^\bullet$, i.e.,

$$\mathrm{gr}_F^p \mathrm{gr}_{F'}^q V = 0 \quad p + q \neq n, \quad V = \bigoplus_{p+q=n} V^{p,q}, \quad V^{p,q} = F^p \cap F'^q$$

Morphisms are filtered morphisms. A polarization of V is a hermitian form h such that $V = \bigoplus_{p+q=n} V^{p,q}$ is h -orthogonal, and $\bigoplus_{p+q=n} (-1)^p h|_{V^{p,q}}$ is positive definite.

Definition 2.3. A complex mixed Hodge structure (C-MHS) is $(V, W_\bullet, F^\bullet, F'^\bullet)$, with $W_\bullet$ an increasing (weight) filtration, such that for any n , the filtrations induced by $F^\bullet, F'^\bullet$ on $\mathrm{gr}_n^W V$ define a pure Hodge structure of weight n .

Morphisms are those compatible with all three filtrations. A graded polarization is a polarization h_n on $\mathrm{gr}_n^W V$ for each n .

Definition 2.4. A complex pure twistor structure (C-TS) of weight n is a vector bundle $\mathcal{V} \simeq \mathcal{O}_{\mathbb{P}^1}(n)^{\oplus r}$. A complex mixed twistor structure (C-MTS) is a pair $(\mathcal{V}, W_\bullet)$ where $\mathcal{V}$ is a vector bundle on $\mathbb{P}^1$, $W_\bullet \mathcal{V}$ is a locally split increasing (weight) filtration by subbundles satisfying

$$\mathrm{gr}_n^W \mathcal{V} \simeq \mathcal{O}_{\mathbb{P}^1}(n)^{\oplus r_n}$$

Morphisms of complex mixed twistor structures are $\mathcal{O}_{\mathbb{P}^1}$ -linear morphisms.

The complex mixed Hodge and twistor structures form categories $\mathrm{MHS}_{\mathbb{C}}$ and $\mathrm{MTS}_{\mathbb{C}}$. We have fiber functors ω , with image $\mathbb{C}$ -vector spaces, where $\omega(V, W_\bullet, F^\bullet, F'^\bullet) = V$, and $\omega(\mathcal{V}, W_\bullet) = \mathcal{V}|_{\lambda=1}$. Both are then neutral Tannakian categories.

Definition 2.5. A graded polarized complex variation of mixed Hodge structures (C-VMHS) on X is $(\mathbb{V}, W_\bullet, F^\bullet, F'^\bullet)$ where

- $\mathbb{V}$ is a complex local system on X and $W_\bullet$ is a flat increasing (weight) filtration;
- $F^\bullet, F'^\bullet$ are locally split decreasing filtrations

$$F^p \subset \mathcal{O}_X \otimes_{\mathbb{C}_X} \mathbb{V}, \quad F'^q \subset \overline{\mathcal{O}}_X \otimes_{\mathbb{C}_X} \mathbb{V}$$

by holomorphic and anti-holomorphic subbundles respectively;

- for the natural flat connections $\nabla, \overline{\nabla}$ coming from the exterior derivative, these must satisfy Griffiths transversality

$$\nabla F^p \subset F^{p-1} \otimes \Omega_X^1, \quad \overline{\nabla} F'^q \subset F'^{q-1} \otimes \Omega_X^1$$

- there exists a flat hermitian form h_n on each $\mathrm{gr}_n^W \mathbb{V}$ such that for each $x \in X$, the restriction of the tuple to x gives a graded polarized complex mixed hodge structure.

For the twistor analogue, we first review the theory of harmonic bundle developed by Corlette and Simpson in [Cor88; Sim92].

Definition 2.6. Let $\mathcal{E}$ be a complex $\mathcal{C}^\infty$ -vector bundle on a projective complex manifold X equipped with a flat $\mathcal{C}^\infty$ -connection D . For h a hermitian metric on $\mathcal{E}$, we have a decomposition

$$D = D_h + \Psi$$

where D_h is unitary with respect to h , and Ψ is self-adjoint for h . Take a further type decomposition into $(1, 0)$ and $(0, 1)$ components

$$D_h = \partial_h + \bar{\partial}_h, \quad \Psi = \theta + \theta^*$$

We say that h is pluriharmonic if $(\theta + \bar{\partial}_h)^2 = 0$. A harmonic bundle $(\mathcal{E}, D, h)$, equivalently $(\mathcal{E}, \bar{\partial}_h, \theta, h)$, is then a smooth complex vector bundle $\mathcal{E}$ with a flat smooth connection D and a pluriharmonic metric h .

Let $\mathcal{A}_X := \mathcal{C}_X^\infty \boxtimes \mathcal{O}_{\mathbb{P}^1}$ be the sheaf of C^∞ -functions on $X_{\mathbb{P}^1} := X \times \mathbb{P}^1$ which are holomorphic in the $\mathbb{P}^1$ -direction, and choose generating sections $x_0, x_\infty \in H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}(1))$ vanishing at 0 and ∞ respectively. Then we get a derivative

$$x_0 \partial_{X_{\mathbb{P}^1}/\mathbb{P}^1} + x_\infty \bar{\partial}_{X_{\mathbb{P}^1}/\mathbb{P}^1} : \mathcal{A}_X \rightarrow \mathcal{A}_X \otimes \Omega_{X_{\mathbb{P}^1}/\mathbb{P}^1}^1(1)$$

with $\Omega_{X_{\mathbb{P}^1}/\mathbb{P}^1}^1(1) := \Omega_X^1 \boxtimes \mathcal{O}_{\mathbb{P}^1}(1)$, where Ω_X^1 is the sheaf of C^∞ -one-forms on X . Let $(\mathcal{E}, \bar{\partial}_h, \theta, h)$ be a harmonic bundle on X , and define $\mathcal{E}_{\mathbb{P}^1} := \mathcal{E} \boxtimes \mathcal{O}_{\mathbb{P}^1}$. Then we have a $x_0 \partial_{X_{\mathbb{P}^1}/\mathbb{P}^1} + x_\infty \bar{\partial}_{X_{\mathbb{P}^1}/\mathbb{P}^1}$ -connection

$$\mathcal{D} := x_0(\partial_h + \theta^*) + x_\infty(\theta + \bar{\partial}_h) : \mathcal{E}_{\mathbb{P}^1} \rightarrow \mathcal{E}_{\mathbb{P}^1} \otimes \Omega_{X_{\mathbb{P}^1}/\mathbb{P}^1}^1(1), \quad \mathcal{D}^2 = 0$$

where we abuse notation so that $\partial_h + \theta^*, \theta + \bar{\partial}_h$ really mean their natural extensions to $\mathcal{E}_{\mathbb{P}^1}$.

Definition 2.7. The resulting $(\mathcal{E}_{\mathbb{P}^1}, \mathcal{D})$ is the variation of pure twistor structures ($\mathbb{C}$ -VTS) of weight 0 associated to the harmonic bundle $(\mathcal{E}, \bar{\partial}_h, \theta, h)$. We get the weight k $\mathbb{C}$ -VTS associated to $(\mathcal{E}, \bar{\partial}_h, \theta, h)$ by twisting $\mathcal{E}_{\mathbb{P}^1, k} := \mathcal{E} \boxtimes \mathcal{O}_{\mathbb{P}^1}(k)$.

Remark 2.8. Every semisimple complex local system on X underlies a harmonic bundle, hence a variation of pure twistor structures.

Definition 2.9. A graded-polarized variation of mixed twistor structures ($\mathbb{C}$ -VMTS) on X is a triple $(\mathcal{V}, W_\bullet, \mathcal{D})$ such that

- $\mathcal{V}$ is an $\mathcal{A}_X$ -module on $X_{\mathbb{P}^1}$;
- $\mathcal{D} : \mathcal{V} \rightarrow \mathcal{V} \otimes \Omega_{X_{\mathbb{P}^1}/\mathbb{P}^1}^1(1)$ is a $x_0 \partial_{X_{\mathbb{P}^1}/\mathbb{P}^1} + x_\infty \bar{\partial}_{X_{\mathbb{P}^1}/\mathbb{P}^1}$ -connection with $\mathcal{D}^2 = 0$;
- $W_\bullet \mathcal{V}$ is a $\mathcal{D}$ -flat increasing weight filtration of $\mathcal{E}_{\mathbb{P}^1}$;
- each $(\text{gr}_k^W \mathcal{V}, \text{gr}_k^W \mathcal{D})$ has a pluriharmonic hermitian metric h_k making it a $\mathbb{C}$ -VTS of weight k associated to a harmonic bundle.

Morphisms are $\mathcal{A}_X$ -linear morphisms which preserve $W_\bullet$ and commutes with $\mathcal{D}$.

Remark 2.10. On an affine chart $\mathbb{A}_\lambda^1 = \mathbb{P}^1 - \{\infty\}$, let $D_\lambda = x_\infty^{-1} \mathcal{D}|_{X \times \{\lambda\}}$. The data of a $\mathbb{C}$ -VMTS is equivalently a family of integrable λ -connections

$$\nabla_\lambda(fs) = \lambda df \otimes s + f \nabla_\lambda(s), \quad \nabla_\lambda^2 = 0$$

where $\nabla_\lambda = D_\lambda^{0,1}$ on the holomorphic vector bundle $(\mathcal{V}|_\lambda, D_\lambda^{0,1})$. The fibers $\lambda = 1$ and $\lambda = 0$ give a flat bundle, hence an underlying local system by Riemann-Hilbert correspondence, and a Higgs bundle respectively.

2.3. Variations with artinian coefficients. Let $\mathbf{MS}_{\mathbb{C}}$ be either $\mathbf{MHS}_{\mathbb{C}}$ or $\mathbf{MTS}_{\mathbb{C}}$, and $\mathbf{T}(0)$ be the tensor unit in $\mathbf{MS}_{\mathbb{C}}$.

Definition 2.11. A $\mathbf{T}(0)$ -MS-algebra is a commutative unital algebra object $\mathbf{A} \in \mathbf{MS}_{\mathbb{C}}$, i.e., $\mathbf{A} \in \mathbf{MS}_{\mathbb{C}}$ with morphisms in $\mathbf{MS}_{\mathbb{C}}$:

$$\mathbf{T}(0) \xrightarrow{u} \mathbf{A}, \quad \mathbf{A} \otimes \mathbf{A} \xrightarrow{\mu} \mathbf{A}$$

satisfying the usual unit, associativity, and commutativity diagrams. Similarly, we can define $\mathbf{A}$ -MS-algebra (and module) from a $\mathbf{T}(0)$ -MS-algebra $\mathbf{A}$.

We say that $\mathbf{A}$ is artinian local if $\omega(\mathbf{A})$ is an artinian local $\mathbb{C}$ -algebra, and the residue map $\mathbf{A} \rightarrow \mathbf{T}(0)$ is a morphism in $\mathbf{MS}_{\mathbb{C}}$. Since $\mathbf{MS}_{\mathbb{C}}$ is abelian, $\mathfrak{m}_{\mathbf{A}} := \ker(\mathbf{A} \rightarrow \mathbf{T}(0))$ is a subobject of $\mathbf{A}$ in $\mathbf{MS}_{\mathbb{C}}$, and we have a splitting $\mathbf{A} = \mathbf{T}(0) \oplus \mathfrak{m}_{\mathbf{A}}$.

Definition 2.12.

- (1) A pro-MS is a pro-object in $\mathbf{MS}_{\mathbb{C}}$.
- (2) A pro- $\mathbf{T}(0)$ -MS-algebra is a $\mathbf{T}(0)$ -algebra object in the category of pro-MS. Equivalently, it is an inverse system of $\mathbf{T}(0)$ -MS-algebras.
- (3) A pro- $\mathbf{T}(0)$ -MS-algebra $\mathbf{\Lambda}$ is complete, noetherian, artinian, or local when $\omega(\mathbf{\Lambda})$ is as a $\mathbb{C}$ -algebra.
- (4) An ideal subobject $\mathbf{I}$ of a (pro-) $\mathbf{T}(0)$ -MS-algebra is a (pro-) $\mathbf{T}(0)$ -MS-ideal if it is a sub pro- $\mathbb{C}$ -MS.
- (5) If $\mathbf{\Lambda}$ is a pro- $\mathbf{T}(0)$ -MS-algebra, we define a pro- $\mathbf{\Lambda}$ -MS-module to be a pro- $\mathbf{\Lambda}$ -module object in $\mathbf{MS}_{\mathbb{C}}$.

Let $a_X : X \rightarrow \text{pt}$, and denote $\mathcal{A} = a_X^* \mathbf{A}$ for $\mathbf{A} \in \mathbf{MS}_{\mathbb{C}}$, i.e., $\mathcal{A}$ is the constant variation with fiber $\omega(\mathbf{A})$.

Definition 2.13. Let $\mathbf{A} \in \mathbf{MS}_{\mathbb{C}}$ be an artinian local $\mathbf{T}(0)$ -MS-algebra. An $\mathbf{A}$ -variation of mixed structures ($\mathbf{A}$ -VMS) is a complex variation of mixed Hodge/twistor structures $\mathcal{V}$ together with an action morphism

$$\mathcal{A} \otimes \mathcal{V} \rightarrow \mathcal{V}$$

which is a morphism of variations of mixed structures and which satisfies unit and associativity diagrams. We also require the underlying local system of $\mathcal{V}$ to be a local system of free $\omega(\mathbf{A})$ -modules.

If $\mathbf{\Lambda}$ is a pro- $\mathbf{T}(0)$ -MS-algebra with

$$\mathbf{\Lambda} = \varprojlim_n \mathbf{\Lambda}_n, \quad \mathbf{\Lambda}_n = \mathbf{\Lambda} / \mathfrak{m}_{\mathbf{\Lambda}}^{n+1}$$

where each $\mathbf{\Lambda}_n$ is an artinian local $\mathbf{T}(0)$ -MS-algebra with $\mathbf{\Lambda}_{n+1} \rightarrow \mathbf{\Lambda}_n$ being a morphism in $\mathbf{MS}_{\mathbb{C}}$, then a pro- $\mathbf{\Lambda}$ -VMS is of the form

$$\mathcal{V} = \varprojlim_n \mathcal{V}_n, \quad \mathcal{V}_n = \mathcal{V} / \mathfrak{m}_{\mathbf{\Lambda}}^{n+1} \mathcal{V}$$

where $\mathcal{V}$ has an underlying local system of free $\omega(\mathbf{\Lambda})$ -modules. Each $\mathcal{V}_n$ is a $\mathbb{C}$ -VMS for which the maps $\mathcal{V}_{n+1} \rightarrow \mathcal{V}_n$ are morphisms of $\mathbb{C}$ -VMS, each $\mathbf{\Lambda} / \mathfrak{m}_{\mathbf{\Lambda}}^{n+1} \subset \text{End}(\mathcal{V}_n)$ is a sub $\mathbb{C}$ -MS.

Proposition 2.14. *Let X be a smooth projective complex variety. Let $\mathbf{A} \in \mathbf{MS}_{\mathbb{C}}$ be an artinian local $\mathbf{T}(0)$ -algebra, and $\mathcal{V}$ an $\mathbf{A}$ -VMS with underlying local system $\mathbb{V}$. Then there is a functorial*

$$K_{\mathbf{A}}(\mathcal{V}) = \mathrm{Ra}_{X,*}^{\mathrm{MS}} \mathcal{V} \in D^b(\mathbf{A}\text{-MS-mod})$$

such that $\omega(K_{\mathbf{A}}(\mathcal{V})) \simeq \mathrm{R}\Gamma(X, \mathbb{V})$ in $D^b(\omega(\mathbf{A})\text{-mod})$. Furthermore, for every morphism $\mathbf{A} \rightarrow \mathbf{B}$ in $\mathbf{MS}_{\mathbb{C}}$ with the corresponding $\mathcal{A} \rightarrow \mathcal{B}$ of constant variations on X , we have base change

$$K_{\mathbf{A}}(\mathcal{V}) \otimes_{\mathbf{A}}^{\mathrm{L}} \mathbf{B} \simeq K_{\mathbf{B}}(\mathcal{V} \otimes_{\mathcal{A}} \mathcal{B})$$

Proof. Forget about the coefficient $\mathbf{A}$, we can regard $\mathcal{V}$ as a smooth object in the derived category of algebraic complex mixed modules. Then the fact that we have $K(\mathcal{V}) = \mathrm{Ra}_{X,*}^{\mathrm{MS}} \mathcal{V} \in D^b(\mathbf{MS}_{\mathbb{C}})$ with $\omega(K(\mathcal{V})) \simeq \mathrm{R}\Gamma(X, \mathbb{V})$ follows from the six functor formalism. In the Hodge setting, this follows from work of Saito [Sai90, Theorem 0.1]; in the twistor setting, this follows from works of Sabbah and Mochizuki [Sab05; Moc15]. We now show compatibility with the $\mathbf{A}$ -structure. Apply the projection formula, which in this specific case is a consequence of six functor formalism and dualizability of $\mathbf{A}$, to the action morphism $(a_X)^* \mathbf{A} \otimes \mathcal{V} \rightarrow \mathcal{V}$ we get

$$\mathbf{A} \otimes \mathrm{Ra}_{X,*}^{\mathrm{MS}} \mathcal{V} \rightarrow \mathrm{Ra}_{X,*}^{\mathrm{MS}} \mathcal{V}$$

and similarly for the unit and associativity diagrams. Regarding the base change $\mathbf{A} \rightarrow \mathbf{B}$, notice that we have a base change morphism by adjunction

$$\mathbf{B} \otimes_{\mathbf{A}}^{\mathrm{L}} \mathrm{Ra}_{X,*}^{\mathrm{MS}} \mathcal{V} \rightarrow \mathrm{Ra}_{X,*}^{\mathrm{MS}} (\mathcal{V} \otimes_{\mathcal{A}} \mathcal{B})$$

which, after applying ω , becomes the base change morphism $\mathrm{R}\Gamma(X, \mathbb{V}) \otimes_{\omega(\mathbf{A})}^{\mathrm{L}} \omega(\mathbf{B}) \rightarrow \mathrm{R}\Gamma(X, \mathbb{V} \otimes_{\mathcal{A}_X} \mathcal{B}_X)$. This follows from the compatibility of ω with the six functors. The base change $\mathrm{R}\Gamma(X, \mathbb{V}) \otimes_{\omega(\mathbf{A})}^{\mathrm{L}} \omega(\mathbf{B}) \rightarrow \mathrm{R}\Gamma(X, \mathbb{V} \otimes_{\mathcal{A}_X} \mathcal{B}_X)$ is an isomorphism due to identifications with free cellular complexes, see (2.3). Finally, ω is faithful and exact, hence its derived version is conservative, and $\mathbf{B} \otimes_{\mathbf{A}}^{\mathrm{L}} \mathrm{Ra}_{X,*}^{\mathrm{MS}} \mathcal{V} \rightarrow \mathrm{Ra}_{X,*}^{\mathrm{MS}} (\mathcal{V} \otimes_{\mathcal{A}} \mathcal{B})$ must be an isomorphism to begin with. $\square$

2.4. Absolute subset and rank one theory. We have a quasi-projective moduli space $R_{\mathrm{DR}}(X, x, r)$ of framed flat connections on X [Sim94, Theorem 6.13]. The Riemann-Hilbert correspondence then induces a complex analytic isomorphism [Sim94, Theorem 7.1]:

$$\mathrm{RH} : R_{\mathrm{DR}}(X, x, r) \xrightarrow{\simeq} R_B(X, x, r) \quad (2.4)$$

Given $(E, \nabla) \in R_{\mathrm{DR}}(X, x, r)$ and $\sigma \in \mathrm{Aut}(\mathbb{C}/\mathbb{Q})$, we have a base change $X^{\sigma} \rightarrow X$ and pulling back gives $(E^{\sigma}, \nabla^{\sigma}) \in R_{\mathrm{DR}}(X^{\sigma}, x, r)$. In fact, this gives a bijection $\sigma : R_{\mathrm{DR}}(X, x, r) \xrightarrow{\simeq} R_{\mathrm{DR}}(X^{\sigma}, x, r)$.

Definition 2.15. A GL_r -invariant subset $\Sigma \subseteq R_B(X, x, r)$ is absolute $\overline{\mathbb{Q}}$ -constructible if for any $\sigma \in \mathrm{Aut}(\mathbb{C}/\mathbb{Q})$,

$$\Sigma^{\sigma} := \mathrm{RH} \circ \sigma \circ \mathrm{RH}^{-1}(\Sigma) \subseteq R_B(X^{\sigma}, x, r)$$

is $\overline{\mathbb{Q}}$ -constructible, i.e., a finite union of locally closed subvarieties defined over $\overline{\mathbb{Q}}$.

We have similar notions for the stacks $\mathcal{M}_{B/\text{DR}}(X, r)$ as well as the good moduli spaces $M_{B/\text{DR}}(X, r)$. Now, consider the rank one character variety

$$M_B(X, 1) = \text{Hom}(\pi_1(X, x), \mathbb{C}^*) \simeq (\mathbb{C}^*)^{\text{rank } H_1(X, \mathbb{Z})} \times \text{Hom}(H_1(X, \mathbb{Z})_{\text{tors}}, \mathbb{C}^*) \quad (2.5)$$

which is a finite disjoint union of algebraic tori.

Theorem 2.16 ([BW20; Sim93]). *Every irreducible absolute $\overline{\mathbb{Q}}$ -closed subset $\Sigma \subseteq M_B(X, 1)$ is a torsion-translated algebraic subtorus*

$$\Sigma = \tau \cdot N$$

where $N \subseteq M_B(X, 1)$ is a subtorus, and $\text{Lie}(N) \subseteq H^1(X, \mathbb{C})$ is a complex Hodge substructure. Absolute $\overline{\mathbb{Q}}$ -constructible subsets are obtained from a finite sequence of unions, intersections and complements.

Proof. The first and last statement follows from [Sim93, Theorem 6.1] and [BW20, Theorem 9.1.2]. Regarding the Lie algebra, by [Sim93, Theorem 6.1], N is a triple torus, hence by [Sim93, Theorem 2.1] there exists abelian subvariety $P \subseteq \text{Alb}(X)$ such that N is the image of $M_B(\text{Alb}(X)/P, 1) \hookrightarrow M_B(X, 1)$. The map on tangent space is

$$H^1(\text{Alb}(X)/P, \mathbb{C}) \hookrightarrow H^1(X, \mathbb{C})$$

is induced by $\pi^* : H^1(\text{Alb}(X)/P, \mathbb{C}) \rightarrow H^1(\text{Alb}(X), \mathbb{C}) \simeq H^1(X, \mathbb{C})$ where $\pi : \text{Alb}(X) \rightarrow \text{Alb}(X)/P$ is the projection. Pullback is a Hodge morphism hence we are done. $\square$

Corollary 2.17 ([BW20, Theorem 11.1.1]). *Let $\mathcal{F} \in D_c^b(X, \mathbb{C})$ be of geometric origin, then the twisted jump loci*

$$\Sigma_k^i(X, \mathcal{F}) = \{\mathbb{V} \in R_B(X, 1) \mid \dim \mathbb{H}^i(X, \mathcal{F} \otimes \mathbb{V}) \geq k\}$$

is absolute $\overline{\mathbb{Q}}$ -closed and thus a finite union of torsion-translated algebraic subtori. If $\mathcal{F}$ is not of geometric origin, then $\Sigma_k^i(X, \mathcal{F})$ is only a finite union of translated algebraic subtori.

2.5. Determinant sheet. In this section, let $A = M_B(X, 1)$ and the decomposition (2.5) be $A = A^\circ \times \hat{T}$, where $T = H_1(X, \mathbb{Z})_{\text{tors}}$. Since T is finite, its dual $\hat{T}$ is also finite. We have the squaring map

$$[2] : A \rightarrow A, \quad \mathbb{L} \mapsto \mathbb{L}^{\otimes 2}$$

and let $A^2 = \text{im}[2]$, $A[2] = \ker[2]$. Then $A/A^2 \simeq \hat{T}/\hat{T}^2$ is a finite group. For each $\bar{\epsilon} \in A/A^2$ we can pick a torsion lift $\epsilon \in A$. Let ϵA^2 be the coset $\epsilon A^2 = \{\epsilon \otimes \mathbb{L}^{\otimes 2} \mid \mathbb{L} \in A\}$, then

$$A = \bigsqcup_{\bar{\epsilon} \in A/A^2} \epsilon A^2$$

and thus

$$M_B(X, 2) = \bigsqcup_{\bar{\epsilon} \in A/A^2} M_B(X, 2, \det \in \epsilon A^2)$$

For each $\bar{\epsilon} \in A/A^2$ with torsion lift $\epsilon \in A$, we have the following proposition

Proposition 2.18. *There is a Cartesian diagram*

$$\begin{array}{ccc}
 M_B(X, 2, \det = \epsilon) \times A & \xrightarrow[\Phi_\epsilon]{(\mathbb{V}, \mathbb{L}) \mapsto \mathbb{V} \otimes \mathbb{L}} & M_B(X, 2, \det \in \epsilon A^2) \\
 \text{pr}_2 \downarrow & & \downarrow \det \\
 A & \xrightarrow{\mathbb{L} \mapsto \epsilon \otimes \mathbb{L}^{\otimes 2}} & A
 \end{array} \tag{2.5}$$

and Φ_ϵ is a finite étale $A[2]$ -torsor over $M_B(X, 2, \det \in \epsilon A^2)$, with the action

$$\rho \cdot (\mathbb{V}, \mathbb{L}) = (\mathbb{V} \otimes \rho, \mathbb{L} \otimes \rho^{-1})$$

The same is true for the quotient stacks and framed moduli schemes.

Proof. Notice that

$$M_B(X, 2, \det \in \epsilon A^2) \times_A A = \{(\mathbb{W}, \mathbb{L}) \mid \det \mathbb{W} = \epsilon \otimes \mathbb{L}^{\otimes 2}\}$$

which is isomorphic to $M_B(X, 2, \det = \epsilon) \times A$ via the maps

$$\begin{aligned}
 M_B(X, 2, \det \in \epsilon A^2) \times_A A &\rightarrow M_B(X, 2, \det = \epsilon) \times A \\
 (\mathbb{W}, \mathbb{L}) &\mapsto (\mathbb{W} \otimes \mathbb{L}^{-1}, \mathbb{L}) \\
 (\mathbb{V} \otimes \mathbb{L}, \mathbb{L}) &\leftarrow (\mathbb{V}, \mathbb{L})
 \end{aligned}$$

Next, the map downstairs is a $A[2]$ -torsor over ϵA^2 , with the action $\rho \cdot \mathbb{L} = \mathbb{L} \otimes \rho^{-1}$. Hence Φ_ϵ is also a $A[2]$ -torsor. The induced action on $M_B(X, 2, \det \in \epsilon A^2) \times_A A$ is $\rho \cdot (\mathbb{W}, \mathbb{L}) = (\mathbb{W}, \mathbb{L} \otimes \rho^{-1})$ which translate to the stated action on $M_B(X, 2, \det = \epsilon) \times A$. $\square$

Now, let $Z \subset M_B(X, 2)$ be an irreducible subvariety, then it must be contained in some $M_B(X, 2, \det \in \epsilon A^2)$. We can then look at $\phi_\epsilon^{-1}(Z)$ and pick an irreducible component C . Set $H_C = \{\rho \in A[2] \mid \rho \cdot C = C\}$.

Proposition 2.19. *The induces map between normalizations $C^\nu \rightarrow Z^\nu$ is a finite étale H_C -torsor, and $Z^\nu \simeq C^\nu/H_C$.*

Proof. We have a base change diagram

$$\begin{array}{ccc}
 W & \xrightarrow{f} & \Phi_\epsilon^{-1}(Z) \\
 g \downarrow & & \downarrow \phi_\chi \\
 Z^\nu & \xrightarrow{\nu} & Z
 \end{array}$$

where g is a finite étale $A[2]$ -torsor. Let η_Z be the generic point of Z , then the generic point η_C of C lies over η_Z . Notice that we have an isomorphism of function fields $\kappa(\eta_{Z^\nu}) \simeq \kappa(\eta_Z)$, hence the an isomorphism of generic fibers $W_{\eta_{Z^\nu}} \simeq \Phi_\epsilon^{-1}(Z)_{\eta_Z}$. Thus there is a $W_{\eta_{Z^\nu}} \ni \eta \mapsto \eta_C$ under this identification. Let $\overline{C}$ be the closure of η in W . Since $W \rightarrow Z^\nu$ is finite étale, $\overline{C}$ is an irreducible component of W . It follows that $\overline{C}$ is open and closed in W , and $\overline{C} \rightarrow Z^\nu$ is a finite étale H_C -torsor.

On the other hand, $W \rightarrow \Phi_\epsilon^{-1}(Z)$ is finite, since it is the base change of ν which is finite. Thus $\overline{C} \rightarrow C$ is finite. It is also birational by construction, hence $\overline{C} = C^\nu$. $\square$

3. HODGE THEORY OF R_B AND M_B

In this section, we prove that the cohomological jump loci enjoy several nice Hodge theoretic properties. We first briefly recall some results and definitions, following the exposition of [BBT24].

3.1. Betti universal deformation algebra and Deligne-Hitchin space.

Theorem 3.1 ([BBT24, Theorem 4.2]). *Let $(\mathbb{V}, \phi)$ be a framed rank r local system underlying a $\mathbb{C}$ -VMS $\mathcal{V}$, then the completed local ring $\Lambda_{(\mathbb{V}, \phi)} := \widehat{\mathcal{O}}_{R_B(X, x, r), (\mathbb{V}, \phi)}$ has a pro- $\mathbf{T}(0)$ -MS-algebra structure $\Lambda_{(\mathbb{V}, \phi)}$ with*

$$\omega(\Lambda_{(\mathbb{V}, \phi)}) = \Lambda_{(\mathbb{V}, \phi)}$$

and the universal formal local system $\widehat{\mathbb{V}}$ underlies a free pro- $\Lambda_{(\mathbb{V}, \phi)}$ -VMS $\widehat{\mathcal{V}}$. The universal framing $\widehat{\phi} : \widehat{\mathbb{V}}_x \rightarrow \Lambda_{(\mathbb{V}, \phi)} \widehat{\otimes}_{\mathbb{C}} \mathbb{C}^r$ is the image under ω of a morphism of pro- $\Lambda_{(\mathbb{V}, \phi)}$ -MS-modules $\widehat{\phi} : \widehat{\mathbb{V}}_x \rightarrow \Lambda_{(\mathbb{V}, \phi)} \widehat{\otimes}_{\mathbf{T}(0)} \mathbf{T}(0)^{\oplus r}$.

These structures are uniquely determined by the following universal property: For any artinian local $\mathbf{T}(0)$ -MS-algebra $\mathbf{A}$ and any $\mathbf{A}$ -VMS $\mathcal{U}$ with a framing $\psi : \mathcal{U}_x \rightarrow \mathbf{A} \otimes_{\mathbf{T}(0)} \mathbf{T}(0)^{\oplus r}$ which restricts to $(\mathcal{V}, \phi) \bmod \mathfrak{m}_{\mathbf{A}}$, there is a unique morphism $\Lambda_{(\mathbb{V}, \phi)} \rightarrow \mathbf{A}$ of local pro- $\mathbf{T}(0)$ -MS-algebras such that

$$(\mathbf{A}, (\mathcal{U}, \psi)) \simeq \mathbf{A} \widehat{\otimes}_{\Lambda_{(\mathbb{V}, \phi)}} \left(\Lambda_{(\mathbb{V}, \phi)}, (\widehat{\mathcal{V}}, \widehat{\phi}) \right)$$

or, in other words, $\Lambda_{(\mathbb{V}, \phi)}$ represents the functor

$$\mathbf{A} \mapsto \{\text{framed free } \mathbf{A}\text{-VMS deformations of } (\mathcal{V}, \phi)\} / \simeq$$

For a $\mathbb{C}$ -scheme T , denote $X_T = X \times T$ with projection $\pi_T : X_T \rightarrow T$.

Proposition 3.2 ([Sim96, Proposition 4.1]). *The moduli functor*

$$R_{\text{Hod}}(r)(T) = \left\{ \left((\lambda, E_T, \nabla_{\lambda, T}, \phi) \mid \begin{array}{l} E_T \text{ locally free of rank } r \text{ on } X_T, \\ \lambda \in \Gamma(T, \mathcal{O}_T), \\ \pi_T^{-1} \mathcal{O}_T \text{--linear } \nabla_{\lambda, T} : E_T \rightarrow E_T \otimes \Omega_{X_T/T}^1, \\ \nabla_{\lambda, T}^2 = 0, \\ \nabla_{\lambda, T}(fs) = \lambda d_{X_T/T} f \otimes s + f \nabla_{\lambda, T} s \\ \phi : E_T|_{x \times T} \simeq \mathcal{O}_T^r \end{array} \right) \right\} / \simeq$$

is representable by a relative scheme $R_{\text{Hod}}(X, x, r) \xrightarrow{\lambda} \mathbb{A}_\lambda^1$ parametrizing λ -connections with a framing. Furthermore, there is a $\mathbb{G}_m$ -action on $R_{\text{Hod}}(X, x, r)$, covering the scaling action on $\mathbb{A}_\lambda^1$, by scaling the λ -connection.

Definition 3.3. Consider the complex conjugate variety X^c , we can similarly define $R_{\text{Hod}}(X^c, x, r)$ and there is a holomorphic isomorphism

$$R_{\text{Hod}}(X, x, r)|_{\mathbb{G}_m} \simeq R_{\text{Hod}}(X^c, x, r)|_{\mathbb{G}_m} \quad (3.1)$$

and there is a gluing to get the *Deligne-Hitchin* scheme $R_{\text{DH}}(X, x, r) \rightarrow \mathbb{P}^1$. We have the derivations associated to the $\mathbb{G}_m$ -actions on $R_{\text{Hod}}(X, x, r)$ and $R_{\text{Hod}}(X^c, x, r)$ which glue to a derivation Θ on $R_{\text{DH}}(X, x, r)$.

Proposition 3.4 ([BBT24, Corollary 8.8]). *Let $\mathbf{A}$ be an artinian local $\mathbf{T}(0)$ -MS-algebra. An $\mathbf{A}$ -VMTS $\mathcal{V}$ is equivalent to a $\mathbb{P}^1$ -morphism $\text{Spec}_{\mathbb{P}^1} \mathbf{A} \rightarrow R_{\text{DH}}(X, x, r)$.*

Definition 3.5. If $\mathcal{V}$ is a $\mathbb{C}$ -VTS on X , then the resulting $\mathbb{P}^1 \rightarrow R_{\text{DH}}(X, x, r)$ is called a preferred section.

More concretely, if $(\mathbb{V}, \phi) \in R_B(X, x, r)$ is a semisimple framed local system, then it underlies a canonical weight 0 $\mathbb{C}$ -VTS, giving a preferred section

$$s_{(\mathbb{V}, \phi)} : \mathbb{P}^1 \rightarrow R_{\text{DH}}(X, x, r) \quad (3.2)$$

where $s_{(\mathbb{V}, \phi)}(1)$ is the flat bundle corresponding to $(\mathbb{V}, \phi)$ via the Riemann-Hilbert correspondence.

Proposition 3.6 ([BBT24, Lemma 8.11]). *We have a classifying map for the universal pro- $\mathbf{A}_{(\mathbb{V}, \phi)}$ -VMS $\widehat{\mathcal{V}}$,*

$$\widehat{s}_{(\mathbb{V}, \phi)} : \text{Spf}_{\mathbb{P}^1} \mathbf{A}_{(\mathbb{V}, \phi)} \xrightarrow{\sim} \widehat{R}_{\text{DH}}(X, x, r)_{s_{(\mathbb{V}, \phi)}} \quad (3.3)$$

which is an isomorphism onto the completion of $R_{\text{DH}}(X, x, r)$ along the preferred section.

3.2. Jump loci are Hodge subvarieties. Let $\mathcal{M}_B(X, r)$ be the quotient stack of local systems of rank r , and $\mathcal{Z} \subseteq \mathcal{M}_B(X, r)$ locally closed substack. Let $\mathbb{V} \in \mathcal{Z}(\mathbb{C})$ be a semisimple local system, and pick a framing $\phi : \mathbb{V}_x \xrightarrow{\sim} \mathbb{C}^r$. Denote $R\mathcal{Z} \subseteq R_B(X, x, r)$ the base change to the framed space.

Definition 3.7 ([BBT24, Section 8.5]). $\mathcal{Z}$ is *Hodge* at $\mathbb{V}$ if

- (H1): equipping $\mathbb{V}$ with the canonical weight 0 $\mathbb{C}$ -VTS, the quotient $\Lambda_{(\mathbb{V}, \phi)} \twoheadrightarrow \widehat{\mathcal{O}}_{R\mathcal{Z}, (\mathbb{V}, \phi)}$ underlies a quotient map of pro- $\mathbf{T}(0)$ -MTS-algebras. If $\mathbb{V}$ is a $\mathbb{C}^*$ -fixed point, then we can equip $\mathbb{V}$ with a canonical weight 0, and the quotient $\Lambda_{(\mathbb{V}, \phi)} \twoheadrightarrow \widehat{\mathcal{O}}_{R\mathcal{Z}, (\mathbb{V}, \phi)}$ underlies a quotient map of pro- $\mathbf{T}(0)$ -MHS-algebra. Denote $\mathbf{\Gamma}_{\mathcal{Z}, (\mathbb{V}, \phi)}$ the corresponding pro- $\mathbf{T}(0)$ -MS-algebra on $\widehat{\mathcal{O}}_{R\mathcal{Z}, (\mathbb{V}, \phi)}$.
- (H2): The formal twistor subspace $\text{Spf}_{\mathbb{P}^1} \mathbf{\Gamma}_{\mathcal{Z}, (\mathbb{V}, \phi)} \subseteq \text{Spf}_{\mathbb{P}^1} \mathbf{A}_{(\mathbb{V}, \phi)}$ is tangent to the derivation Θ in Definition 3.3.
- (H3): We have an induced map by composition

$$\widehat{s}_{R\mathcal{Z}, (\mathbb{V}, \phi)} : \text{Spf}_{\mathbb{P}^1} \mathbf{\Gamma}_{\mathcal{Z}, (\mathbb{V}, \phi)} \rightarrow \widehat{R}_{\text{DH}}(X, x, r)_{s_{(\mathbb{V}, \phi)}}$$

Denote Z_{Hod} (resp. $Z_{\overline{\text{Hod}}}$) the Zariski closure of the image of $\widehat{s}_{R\mathcal{Z}, (\mathbb{V}, \phi)}|_{\mathbb{A}_\lambda^1}$ (resp. $\widehat{s}_{R\mathcal{Z}, (\mathbb{V}, \phi)}|_{\mathbb{P}^1 \setminus \{0\}}$) in $R_{\text{Hod}}(X, x, r)$ (resp. $R_{\text{Hod}}(X^c, x, r)$). Then $\widehat{s}_{R\mathcal{Z}, (\mathbb{V}, \phi)}|_{\mathbb{G}_m}$ induces an isomorphism onto the completion of $Z_{\text{Hod}}|_{\mathbb{G}_m}$ along the image of $s_{(\mathbb{V}, \phi)}$, and the same is true for $\widehat{s}_{R\mathcal{Z}, (\mathbb{V}, \phi)}|_{\mathbb{G}_m}$ and $Z_{\overline{\text{Hod}}}|_{\mathbb{G}_m}$.

$\mathcal{Z}$ is a *Hodge substack* if it is closed under semisimplification and Hodge at every semisimple $\mathbb{V} \in \mathcal{Z}(\mathbb{C})$. Notice that these conditions rely only on $R\mathcal{Z} \subseteq R_B(X, x, r)$, so we have a similar notion of *Hodge subvariety* for a GL_r -invariant subvariety

$Z \subseteq R_B(X, x, r)$. We also have obvious generalization to Hodge subvarieties of products of character varieties.

Lemma 3.8. *Let $\mathbf{A} \in \mathbf{MS}_{\mathbb{C}}$ be an artinian local $\mathbf{T}(0)$ -MS-algebra, with $A = \omega(\mathbf{A})$, and $\mathbf{K} \in D^b(\mathbf{A}\text{-MS-mod})$ with $K = \omega(\mathbf{K})$ quasi-isomorphic to a bounded above complex of free A -modules. Then $J_k^i(K) \subseteq A$ is the image under ω of an ideal subobject $J_k^i(\mathbf{K}) \subseteq \mathbf{A}$. This construction commutes with $\mathbf{T}(0)$ -MS-algebra coefficient base change.*

Proof. Let $G = \text{Aut}^{\otimes}(\omega)$ be the Tannakian dual of $\mathbf{MS}_{\mathbb{C}}$. Then we have an equivalence of category $\mathbf{MS}_{\mathbb{C}} \simeq \text{Rep}_{\mathbb{C}}^{\text{fd}}(G)$. Under this equivalence, $\mathbf{A}$ is identified with $A = \omega(\mathbf{A})$ with an algebraic G -action, i.e., a comodule structure $\delta_A : A \rightarrow A \otimes \mathcal{O}(G)$; this is an action by algebra automorphism. Similarly, $\mathbf{K}$ gives a G -equivariant complex $K = \omega(\mathbf{K})$ of A -modules.

Let B be a $\mathbb{C}$ -algebra, and $g \in G(B)$, all the tensor product below will be over $\mathbb{C}$ unless specified otherwise. We have a B -algebra automorphism $g : A \otimes B \rightarrow A \otimes B$. Since B is flat over $\mathbb{C}$, the action of g on $K \otimes^L B = K \otimes B$ satisfies

$$g \cdot (bm) = g(b)(g \cdot m) \quad b \in A \otimes B, m \in K^i \otimes B$$

coming from applying ω to the action morphism $\mathbf{A} \otimes \mathbf{K} \rightarrow \mathbf{K}$. It follows that

$$g^*(K \otimes B) := (A \otimes B) \otimes_g (K \otimes B) \simeq K \otimes B$$

since $g : A \otimes B \rightarrow A \otimes B$ is an isomorphism. Determinantal ideals commute with arbitrary base change, and by (2.1) we get

$$g(J_k^i(K \otimes B)) = J_k^i(g^*(K \otimes B)) \simeq J_k^i(K \otimes B)$$

thus $J_k^i(K \otimes B)$ is stable under the action of $g \in G(B)$. It follows that $J_k^i(K)$ is a G -subrepresentation. The equivalence then gives a unique subobject $J_k^i(\mathbf{K}) \subseteq \mathbf{A}$ in $\mathbf{MS}_{\mathbb{C}}$ with $\omega(J_k^i(\mathbf{K})) = J_k^i(K)$. To check that $J_k^i(\mathbf{K})$ is an ideal subobject, it suffices to check that the composition

$$\mathbf{A} \otimes J_k^i(\mathbf{K}) \rightarrow \mathbf{A} \otimes \mathbf{A} \xrightarrow{\mu} \mathbf{A}$$

factors through $J_k^i(\mathbf{K}) \subseteq \mathbf{A}$. Equivalently,

$$\mathbf{A} \otimes J_k^i(\mathbf{K}) \rightarrow \mathbf{A} \rightarrow \mathbf{A}/J_k^i(\mathbf{K})$$

is zero. This follows from ω being faithful and the image of that map under ω is zero. Compatibility with coefficient base change follows from (2.1). $\square$

The jump loci $\Sigma_k^i(X, r) \subseteq R_B(X, x, r)$ is invariant under the conjugation action by GL_r , so we can ask whether it is a Hodge subvariety.

Theorem 3.9. *For smooth projective X , the jump locus $\Sigma_k^i(X, r) \subseteq R_B(X, x, r)$ is a Hodge subvariety. Furthermore, it is closed under semisimplification, and $\Sigma_k^i(X, r)_{\text{red}}$ is a Hodge subvariety which is absolute $\mathbb{Q}$ -closed.*

Every irreducible component of $\Sigma_k^i(X, r)_{\text{red}}$ is a Hodge subvariety and is absolute $\overline{\mathbb{Q}}$ -closed.

Proof. We first verify that $\Sigma_k^i(X, r)$ satisfies the three conditions in [Definition 3.7](#).

(H1). Consider $(\mathbb{V}, \phi) \in R_B(X, x, r)$ semisimple, and let

$$\Lambda = \widehat{\mathcal{O}}_{R_B(X, x, r), (\mathbb{V}, \phi)}, \quad \Lambda_n = \Lambda / \mathfrak{m}_\Lambda^{n+1}$$

Equip $\mathbb{V}$ with the canonical weight 0 $\mathbb{C}$ -VTS. If $\mathbb{V}$ is a $\mathbb{C}^*$ -fixed point, then it underlies a canonical weight 0 $\mathbb{C}$ -VHS. By [Theorem 3.1](#), Λ underlies a pro- $\mathbf{T}(0)$ -MS-algebra $\mathbf{\Lambda}$, and the universal formal local system $(\widehat{\mathbb{V}}, \widehat{\phi})$ underlies a pro- $\mathbf{\Lambda}$ -VMS $(\widehat{\mathcal{V}}, \widehat{\phi})$. Each Λ_n underlies a $\mathbf{T}(0)$ -MS-algebra $\Lambda_n = \mathbf{\Lambda} / \mathfrak{m}_\Lambda^{n+1}$. Write $\mathcal{V}_n = \widehat{\mathcal{V}} \widehat{\otimes}_{\mathbf{\Lambda}} \Lambda_n$, and $\mathbb{V}_n$ the underlying local system of $\mathcal{V}_n$. By [Proposition 2.14](#) we have

$$\mathbf{K}_n = \mathrm{Ra}_{X,*}^{\mathrm{MS}} \mathcal{V}_n \in D^b(\mathbf{\Lambda}_n\text{-MS-mod}), \quad \omega(\mathbf{K}_n) \simeq \mathrm{R}\Gamma(X, \mathbb{V}_n)$$

Let $R = \Gamma(R_B(X, x, r), \mathcal{O}_{R_B(X, x, r)})$, then $R_B(X, x, r) = \mathrm{Spec} R$. By the discussion of universal cellular complex in [Section 2.1](#), we get a complex $C^\bullet(X, \mathbb{V}^{\mathrm{univ}}) \simeq \mathrm{R}\Gamma(X, \mathbb{V}^{\mathrm{univ}})$ of free R -modules which we can think of as a complex of free sheaves on $R_B(X, x, r)$. Its fiber over a point $(\mathbb{W}, \psi) \in R_B(X, x, r)$ gives $\mathrm{R}\Gamma(X, \mathbb{W})$. Notice that in Λ , the jump ideal becomes

$$J_k^i(\mathrm{R}\Gamma(X, \mathbb{V}^{\mathrm{univ}})) \cdot \Lambda = \varprojlim J_k^i(\mathrm{R}\Gamma(X, \mathbb{V}^{\mathrm{univ}})) \cdot \Lambda_n$$

Recall that we have base change

$$\begin{aligned} J_k^i(\mathrm{R}\Gamma(X, \mathbb{V}^{\mathrm{univ}})) \cdot \Lambda_n &\simeq J_k^i(\mathrm{R}\Gamma(X, \mathbb{V}^{\mathrm{univ}}) \otimes_R^L \Lambda_n) \\ &\simeq J_k^i(\mathrm{R}\Gamma(X, \mathbb{V}^{\mathrm{univ}} \otimes_R^L \Lambda_n)) \\ &\simeq J_k^i(\mathrm{R}\Gamma(X, \mathbb{V}_n)) \end{aligned}$$

thus $J_k^i(\mathrm{R}\Gamma(X, \mathbb{V}^{\mathrm{univ}})) \cdot \Lambda \simeq J_k^i(\mathrm{R}\Gamma(X, \widehat{\mathbb{V}}))$. Now, $\omega(\mathbf{K}_n)$ is quasi-isomorphic to a complex of free Λ_n -modules, which is $C^\bullet(X, \mathbb{V}_n)$, hence by [Lemma 3.8](#) we get that $J_k^i(\mathrm{R}\Gamma(X, \widehat{\mathbb{V}}))$ underlies a pro- $\mathbf{T}(0)$ -MS-ideal $J_k^i(\mathbf{K}) \subseteq \mathbf{\Lambda}$. It follows that we get (H1) since $\mathrm{MS}_{\mathbb{C}}$ is abelian.

(H3). We first take a detour. Let $\mathbf{A} \in \mathrm{MTS}_{\mathbb{C}}$ with $A = \omega(\mathbf{A})$, and $\mathcal{W}$ an $\mathbf{A}$ -VMTS with underlying local system $\mathbb{W}$ of free A -modules. By [Remark 2.10](#), this is the same as a family of holomorphic integrable λ -connections. Let $X_\lambda = X \times \mathbb{A}_\lambda^1$ with projection $\pi : X_\lambda \rightarrow \mathbb{A}_\lambda^1$, then the family of λ -connections, over the patch $\mathbb{A}_\lambda^1$, is the same as a holomorphic bundle $\mathcal{W}_\lambda$ on X_λ with an integrable $\nabla_\lambda : \mathcal{W}_\lambda \rightarrow \mathcal{W}_\lambda \otimes \Omega_{X_\lambda/\mathbb{A}_\lambda^1}^1$. The $\mathbf{A}$ -action becomes an $A[\lambda]$ -action. We can then form the λ -De Rham complex

$$\mathrm{DR}_\lambda(\mathcal{W}_\lambda) = \left[\mathcal{W}_\lambda \xrightarrow{\nabla_\lambda} \mathcal{W}_\lambda \otimes \Omega_{X_\lambda/\mathbb{A}_\lambda^1}^1 \rightarrow \cdots \rightarrow \mathcal{W}_\lambda \otimes \Omega_{X_\lambda/\mathbb{A}_\lambda^1}^{\dim X} \right]$$

and $(\mathrm{Ra}_{X,*}^{\mathrm{MS}} \mathcal{W})|_{\mathbb{A}_\lambda^1} \simeq \mathrm{R}\pi_* \mathrm{DR}_\lambda(\mathcal{W}_\lambda) \in D^b(A[\lambda]\text{-mod})$, by [\[Sim97\]](#), proof of Theorem 4.1] and the proof of [Proposition 2.14](#). Denote

$$K_{\mathrm{Hod}}(\mathcal{W}) := \mathrm{R}\pi_* \mathrm{DR}_\lambda(\mathcal{W}_\lambda)$$

Next, let $q : R_{\mathrm{Hod}}(X, x, r) \rightarrow \mathbb{A}_\lambda^1$ be the Hodge moduli space. This carries a universal λ -connection $(\mathcal{V}_\lambda^{\mathrm{univ}}, \nabla_\lambda)$ on $X \times R_{\mathrm{Hod}}$. If the classifying Deligne-Hitchin

section for $\mathcal{W}$ is $s_{\mathcal{W}} : \mathrm{Spec}_{\mathbb{P}^1} \mathbf{A} \rightarrow R_{\mathrm{DH}}(X, x, r)$, denote $s_{\mathcal{W}, \lambda} := s_{\mathcal{W}}|_{\mathbb{A}_{\lambda}^1} : \mathrm{Spec} A[\lambda] \rightarrow R_{\mathrm{Hod}}(X, x, r)$, and

$$(\mathcal{W}_{\lambda}, \nabla_{\lambda}) = s_{\mathcal{W}, \lambda}^*(\mathcal{V}_{\lambda}^{\mathrm{univ}}, \nabla_{\lambda})$$

The complex K_{Hod} satisfies base change (same proof as in [Sta26, Tag 0FM0] works), hence

$$K_{\mathrm{Hod}}(\mathcal{W}) \simeq s_{\mathcal{W}, \lambda}^* K_{\mathrm{Hod}}(\mathcal{V}^{\mathrm{univ}})$$

where $K_{\mathrm{Hod}}(\mathcal{V}^{\mathrm{univ}}) := R\pi_* \mathrm{DR}_{\lambda}(\mathcal{V}_{\lambda}^{\mathrm{univ}}, \nabla_{\lambda})$. We can think of this as a complex of $A[\lambda]$ -modules, hence a notion of λ -jump ideal makes sense. Denote $\Sigma_{k, \mathrm{Hod}}^i(X, r) = V(J_k^i(K_{\mathrm{Hod}}(\mathcal{V}^{\mathrm{univ}})))$. The restriction to $\lambda = 1$ gives the De Rham complex for the flat bundle corresponding to $\mathbb{W}$ via the Riemann-Hilbert correspondence, hence by the comparison theorem this gives $\Sigma_k^i(\mathbb{W})$.

Back to our situation, let $s_n : \mathrm{Spec}_{\mathbb{P}^1} \mathbf{A}_n \rightarrow R_{\mathrm{DH}}(X, x, r)$ be the classifying map for the $\mathbf{A}_n$ -VMTS $\mathcal{V}_n$. The reduction mod $\mathfrak{m}_{\mathbf{A}}$ gives $s_{(\mathbb{V}, \phi)} : \mathbb{P}^1 \rightarrow R_{\mathrm{DH}}(X, x, r)$, the preferred section associated to the canonical weight 0 $\mathbb{C}$ -VTS on semisimple $\mathbb{V}$. Let $\Gamma = \mathbf{A}/J_k^i(\mathbf{K})$ be the pro- $\mathbf{T}(0)$ -MTS-algebra on $\hat{\mathcal{O}}_{\Sigma_k^i, (\mathbb{V}, \phi)}$, and put

$$s_{n, \Sigma} : \mathrm{Spec}_{\mathbb{P}^1} \Gamma_n \rightarrow \mathrm{Spec}_{\mathbb{P}^1} \mathbf{A}_n \xrightarrow{s_n} R_{\mathrm{DH}}(X, x, r)$$

Notice that

$$s_n^* J_k^i(K_{\mathrm{Hod}}(\mathcal{V}^{\mathrm{univ}})) = J_k^i(s_n^* K_{\mathrm{Hod}}(\mathcal{V}^{\mathrm{univ}})) \simeq J_k^i(\mathbf{K}_n|_{\mathbb{A}_{\lambda}^1}) \simeq J_k^i(\mathbf{K}_n)|_{\mathbb{A}_{\lambda}^1}$$

thus the image of $s_{n, \Sigma}|_{\mathbb{G}_m}$ agrees with the $(n+1)$ -th order neighborhood of the image of $s_{(\mathbb{V}, \phi)}|_{\mathbb{G}_m}$ in $\Sigma_{k, \mathrm{Hod}}^i(X, r)$. It follows that the image of $\hat{s}_{\Sigma}|_{\mathbb{G}_m}$ agrees with the completion of $\Sigma_{k, \mathrm{Hod}}^i(X, r)$ along the image of $s_{(\mathbb{V}, \phi)}$, as desired.

(H2). The scaling $\mathbb{G}_m$ -action turns $t \cdot (\mathcal{V}_{\lambda}^{\mathrm{univ}}, \nabla_{\lambda}) \mapsto (\mathcal{V}_{\lambda}^{\mathrm{univ}}, t\nabla_{\lambda})$. Multiplication by t^p in degree p gives a quasi-isomorphism

$$\mathrm{DR}_{\lambda}(\mathcal{V}_{\lambda}^{\mathrm{univ}}, \nabla_{\lambda}) \simeq \mathrm{DR}_{\lambda}(\mathcal{V}_{\lambda}^{\mathrm{univ}}, t\nabla_{\lambda})$$

thus $J_k^i(K_{\mathrm{Hod}}(\mathcal{V}^{\mathrm{univ}}))$ is $\mathbb{G}_m$ -invariant, hence so is $\Sigma_{k, \mathrm{Hod}}^i(X, r)$. It follows that

$$\Theta(J_k^i(K_{\mathrm{Hod}})) \subseteq J_k^i(K_{\mathrm{Hod}})$$

and completing along the image of $s_{(\mathbb{V}, \phi)}$ gives (H2) by the identification in the previous section.

Closed under semisimplification and absoluteness. Closure under semisimplification follows from semicontinuity of local system cohomology. Absoluteness follows from the same proof as in [BW20, Theorem 11.1.1]. Finally, irreducible components and reduction remains Hodge, due to [BBT24, Proposition 8.21(3)], and absolute. $\square$

Corollary 3.10. *Using the notation of Section 2.5, let Z be an irreducible component of $\Sigma_k^i(X, 2) \subseteq R_B(X, x, 2)$. Let C be an irreducible component of $\phi_{\chi}^{-1}(Z) \subset R_B(X, 2, \det = \chi) \times A$, and denote*

$$T := \overline{\mathrm{pr}_2(C)}_{\mathrm{red}}$$

the Zariski closure of the reduced image of the projection onto $A = M_B(X, 1)$. Then T is a torsion-translated subtorus of A .

Furthermore, if $\mathbb{L} \in T$ is a unitary point, then $\hat{\mathcal{O}}_{T,\mathbb{L}}$ underlies a pro- $\mathbf{T}(0)$ -MHS-algebra such that for every $(\mathbb{V}, \mathbb{L}) \in C$ underlying $\mathbb{C}$ -VHS, $\hat{\mathcal{O}}_{T,\mathbb{L}} \rightarrow \hat{\mathcal{O}}_{C,(\mathbb{V},\mathbb{L})}$ is a morphism of pro- $\mathbf{T}(0)$ -MHS-algebras.

Proof. By [BBT24, Proposition 9.10], all the morphisms in

$$\begin{array}{ccc} R_B(X, 2, \det = \chi) \times A & \xrightarrow[\Phi_\chi]{(\mathbb{V}, \mathbb{L}) \mapsto \mathbb{V} \otimes \mathbb{L}} & R_B(X, 2, \det \in \chi A^2) \\ \text{pr}_2 \downarrow & & \downarrow \det \\ A & \xrightarrow{\mathbb{L} \mapsto \chi \otimes \mathbb{L}^{\otimes 2}} & A \end{array}$$

are absolute $\overline{\mathbb{Q}}$ -morphisms [BBT24, Definition 9.9]. By the previous theorem, Z is absolute $\overline{\mathbb{Q}}$ -closed. Thus $\phi_\chi^{-1}(Z)$ is absolute $\overline{\mathbb{Q}}$ -closed, and $\text{pr}_2(C)$ is absolute $\overline{\mathbb{Q}}$ -constructible. By [BBT24, Proposition 9.14], T is absolute $\overline{\mathbb{Q}}$ -closed. Thus by Theorem 2.16, $T = \tau \cdot N$ where N is a subtorus of A , and $\text{Lie}(N)$ is a complex sub Hodge structure of $H^1(X, \mathbb{C})$.

Now, suppose $\mathbb{L} \in T$ is unitary, then $\mathbb{L}$ underlies a canonical weight 0 $\mathbb{C}$ -VHS, and by Theorem 3.1, $\hat{\mathcal{O}}_{A,\mathbb{L}}$ underlies a pro- $\mathbf{T}(0)$ -MHS-algebra. Furthermore, since A and T are smooth, $\hat{\mathcal{O}}_{T,\mathbb{L}}$ and $\hat{\mathcal{O}}_{A,\mathbb{L}}$ are determined by tangent spaces, i.e.,

$$\hat{\mathcal{O}}_{A,\mathbb{L}} \simeq \widehat{\text{Sym}} H^1(X, \mathbb{C})^\vee, \quad \hat{\mathcal{O}}_{T,\mathbb{L}} \simeq \widehat{\text{Sym}} \text{Lie}(N)^\vee$$

and $\hat{\mathcal{O}}_{A,\mathbb{L}} \rightarrow \hat{\mathcal{O}}_{T,\mathbb{L}}$ is a quotient of pro- $\mathbf{T}(0)$ -MHS-algebras. Furthermore, $H^1(X, \mathbb{C})$ is pure of weight 1, hence so is $\text{Lie}(N)$ and we get $\mathfrak{m}_{T,\mathbb{L}} = W_{-1} \hat{\mathcal{O}}_{T,\mathbb{L}}$ with $\mathfrak{m}_{T,\mathbb{L}}/\mathfrak{m}_{T,\mathbb{L}}^2$ is pure of weight -1 . Suppose that $(\mathbb{V}, \mathbb{L}) \in C$ is a $\mathbb{C}$ -VHS point, equivalently $\mathbb{V}$ underlies a weight 0 $\mathbb{C}$ -VHS, then

$$\hat{\mathcal{O}}_{R_B(X, 2, \det = \chi), \mathbb{V}} \hat{\otimes} \hat{\mathcal{O}}_{A,\mathbb{L}} \twoheadrightarrow \hat{\mathcal{O}}_{C,(\mathbb{V}, \mathbb{L})}$$

is a quotient of pro- $\mathbf{T}(0)$ -MHS-algebras. The composition

$$\hat{\mathcal{O}}_{A,\mathbb{L}} \rightarrow \hat{\mathcal{O}}_{R_B(X, 2, \det = \chi), \mathbb{V}} \hat{\otimes} \hat{\mathcal{O}}_{A,\mathbb{L}} \twoheadrightarrow \hat{\mathcal{O}}_{C,(\mathbb{V}, \mathbb{L})}$$

is thus a morphism of pro- $\mathbf{T}(0)$ -MHS-algebras. This factors through $\hat{\mathcal{O}}_{A,\mathbb{L}} \rightarrow \hat{\mathcal{O}}_{T,\mathbb{L}}$ which is a quotient of pro- $\mathbf{T}(0)$ -MHS-algebras, hence the factorization

$$\hat{\mathcal{O}}_{T,\mathbb{L}} \rightarrow \hat{\mathcal{O}}_{C,(\mathbb{V}, \mathbb{L})}$$

is a morphism of pro- $\mathbf{T}(0)$ -MHS-algebras, as desired. $\square$

3.3. Hodge structures on good moduli image. For a Hodge subvariety $W \subseteq R_B(X, x, r)$, at a point $(\mathbb{V}, \phi) \in W$ with $\mathbb{V}$ underlying a canonical weight 0 complex variation of Hodge structures, we have a pro- $\mathbf{T}(0)$ -MHS-algebra structure on $\hat{\mathcal{O}}_{W,(\mathbb{V}, \phi)}$. We prove a similar statement for the reduced image of W under the GIT quotient $q : R_B(X, x, r) \rightarrow M_B(X, r) = R_B(X, x, r)/\text{GL}_2$.

Proposition 3.11. *Let $W \subseteq R_B(X, x, r)$ be a Hodge subvariety, and $Z = q(W)_{\text{red}} \subset M_B(X, r)$. Let $\mathbb{V} \in Z$ be a $\mathbb{C}^*$ -fixed point, hence underlying a weight 0 complex variation of Hodge structures. Then $\hat{\mathcal{O}}_{Z, \mathbb{V}}$ underlies a pro- $\mathbf{T}(0)$ -MHS-algebra, with*

$$\mathfrak{m}_{Z, \mathbb{V}} = W_{-1} \hat{\mathcal{O}}_{Z, \mathbb{V}}$$

Proof. The stabilizer of $(\mathbb{V}, \phi) \in R_B(X, x, r)$ under the GL_r -action is $\text{Aut}(\mathbb{V})$. Denote $H_{\mathbb{V}} = \text{Aut}(\mathbb{V}) \subset \text{GL}_r$. We have an isotypic decomposition

$$\mathbb{V} \simeq \bigoplus \mathbb{V}_{\alpha} \otimes M_{\alpha}$$

where $\{\mathbb{V}_{\alpha}\}$ are pairwise non-isomorphic irreducible local systems, and $\{M_{\alpha}\}$ are complex vector spaces. By equipping each $\mathbb{V}_{\alpha}$ with a weight 0 $\mathbb{C}$ -VHS and M_{α} with a pure complex Hodge structure of type $(0, 0)$, we get an equality of $\mathbb{C}$ -VHS. Then by Schur's lemma, $H_{\mathbb{V}} = \prod \text{GL}(M_{\alpha})$, hence acts by weight 0 automorphisms of the $\mathbb{C}$ -VHS on $\mathbb{V}$.

On the other hand, by picking a framing we get $(\mathbb{V}, \phi) \in W$ lying over $\mathbb{V}$. Recall that included in the definition of Hodge subvariety is the fact that W is GL_r -invariant. Since $\mathbb{V}$ is semisimple, the GL_r -orbit of $(\mathbb{V}, \phi)$ is closed. Then by the Luna étale slice theorem (see [Lun73] or [AHR20, Remark 4.6]) there exists a $H_{\mathbb{V}}$ -invariant affine subscheme $Y \subseteq R_B(X, x, r)$ containing $(\mathbb{V}, \phi)$ such that $\text{GL}_r \times^{H_{\mathbb{V}}} Y \rightarrow R_B(X, x, r)$ is étale; its image $U \subseteq R_B(X, x, r)$ is a GL_r -invariant saturated (i.e., $U = q^{-1} \circ q(U)$) affine open. Here

$$\text{GL}_r \times^{H_{\mathbb{V}}} Y = (\text{GL}_r \times Y) / H_{\mathbb{V}}$$

where the $H_{\mathbb{V}}$ -action is $h \cdot (g, y) = (g \cdot h, h^{-1} \cdot y)$. Furthermore, the diagram

$$\begin{array}{ccc} \text{GL}_r \times^{H_{\mathbb{V}}} Y & \longrightarrow & U \\ \downarrow & & \downarrow \\ Y // H_{\mathbb{V}} & \longrightarrow & U // \text{GL}_r \end{array}$$

is cartesian with two horizontal maps étale. This induces the following cartesian diagram of quotient stacks by restricting to GL_r -invariant W

$$\begin{array}{ccc} [(W \times_{R_B} Y) / H_{\mathbb{V}}] & \longrightarrow & [W / \text{GL}_r] \\ \downarrow & & \downarrow \\ (W \times_{R_B} Y) // H_{\mathbb{V}} & \longrightarrow & Z \end{array}$$

and étaleness of the bottom horizontal map then gives an isomorphism $\hat{\mathcal{O}}_{Z, \mathbb{V}} \simeq \hat{\mathcal{O}}_{W \times_{R_B} Y, (\mathbb{V}, \phi)}^{H_{\mathbb{V}}}$. Now, notice that $\hat{\mathcal{O}}_{Y, (\mathbb{V}, \phi)}$ is a miniversal algebra for $\mathcal{M}_B$ [AHR20, Remark 4.17], and

$$[W / \text{GL}_r] \times_{\mathcal{M}_B} Y \simeq [W / \text{GL}_r] \times_{\mathcal{M}_B} R_B \times_{R_B} Y \simeq W \times_{R_B} Y$$

It follows that the map $\hat{\mathcal{O}}_{Y,(\mathbb{V},\phi)} \rightarrow \hat{\mathcal{O}}_{W \times_{R_B} Y,(\mathbb{V},\phi)}$ is the base change of the miniversal family $\mathrm{Spf} \hat{\mathcal{O}}_{Y,(\mathbb{V},\phi)} \rightarrow \mathcal{M}_B(X, r)$ along $[W/\mathrm{GL}_r] \hookrightarrow \mathcal{M}_B$. Then, by [BBT24, Lemma 4.59] and the fact that W is Hodge at $(\mathbb{V}, \phi)$, $\hat{\mathcal{O}}_{Y,(\mathbb{V},\phi)} \rightarrow \hat{\mathcal{O}}_{W \times_{R_B} Y,(\mathbb{V},\phi)}$ must be a quotient of pro- $\mathbf{T}(0)$ -MHS-algebras; the pro- $\mathbf{T}(0)$ -MHS-algebra on the miniversal $\hat{\mathcal{O}}_{Y,(\mathbb{V},\phi)}$ is given by [BBT24, Theorem 4.3]. Since $H_{\mathbb{V}}$ acts by Hodge morphisms, we get an induced pro- $\mathbf{T}(0)$ -MHS-algebra structure on $\hat{\mathcal{O}}_{Z,\mathbb{V}}$, with

$$\begin{aligned} \mathfrak{m}_{Z,\mathbb{V}} &= \mathfrak{m}_{W \times_{R_B} Y,(\mathbb{V},\phi)}^{H_{\mathbb{V}}} = \left(W_{-1} \hat{\mathcal{O}}_{W \times_{R_B} Y,(\mathbb{V},\phi)} \right)^{H_{\mathbb{V}}} \\ &= W_{-1} \hat{\mathcal{O}}_{W \times_{R_B} Y,(\mathbb{V},\phi)}^{H_{\mathbb{V}}} = W_{-1} \hat{\mathcal{O}}_{Z,\mathbb{V}} \end{aligned}$$

as desired. $\square$

With this pro- $\mathbf{T}(0)$ -MHS-algebra structure on the good moduli space $M_B(X, r)$, we have a version of Corollary 3.10 for $M_B(X, 2)$ instead of the framed space $R_B(X, x, 2)$, as follows.

Corollary 3.12. *Using the notation of Section 2.5, let Z be an irreducible component of the reduced image $\bar{\Sigma}_k^i(X, 2) \subseteq M_B(X, 2)$. Let C be an irreducible component of $\phi_{\chi}^{-1}(Z) \subset M_B(X, 2, \det = \chi) \times A$, and denote*

$$T := \overline{\mathrm{pr}_2(C)}_{\mathrm{red}}$$

the Zariski closure of the reduced image of the projection onto $A = M_B(X, 1)$. Then T is a torsion-translated subtorus of A .

Furthermore, if $\mathbb{L} \in T$ is a unitary point, then $\hat{\mathcal{O}}_{T,\mathbb{L}}$ underlies a pro- $\mathbf{T}(0)$ -MHS-algebra such that for every $(\mathbb{V}, \mathbb{L}) \in C$ underlying $\mathbb{C}$ -VHS, $\hat{\mathcal{O}}_{T,\mathbb{L}} \rightarrow \hat{\mathcal{O}}_{C,(\mathbb{V},\mathbb{L})}$ is a morphism of pro- $\mathbf{T}(0)$ -MHS-algebras.

Proof. Firstly, the image of pr_2 is the same whether we work with $M_B(X, 2)$ or $R_B(X, x, 2)$, hence T is still a torsion-translated subtorus of A . Secondly, every such irreducible component Z is the good moduli image of some irreducible component W of $\Sigma_k^i(X, 2) \subseteq R_B(X, x, 2)$. Next, since the miniversal pro- $\mathbf{T}(0)$ -MHS-algebra structure of [BBT24, Theorem 4.3] is compatible with tensor product, and that the proof of Proposition 3.11 is clearly functorial with respect to the inclusion of Hodge subvarieties $W \subseteq R_B(X, x, r)$, we get that

$$\hat{\mathcal{O}}_{M_B(X, 2, \det=\chi), \mathbb{V}} \hat{\otimes} \hat{\mathcal{O}}_{A, \mathbb{L}} \rightarrow \hat{\mathcal{O}}_{C, (\mathbb{V}, \mathbb{L})}$$

is a quotient of pro- $\mathbf{T}(0)$ -MHS-algebras for when $\mathbb{V}, \mathbb{L}$ underlie weight 0 complex variation of Hodge structures. Then the same proof as in Corollary 3.10 works. $\square$

3.4. Hodge structures on normalization. The goal of this section is to show that if we have a normalization $C^{\nu} \rightarrow C$, and $(\mathbb{V}, \mathbb{L})$ is a $\mathbb{C}$ -VHS point of C , then for every $c \in C^{\nu}$ lying over $(\mathbb{V}, \mathbb{L})$ there is a pro- $\mathbf{T}(0)$ -MHS-algebra structure on $\hat{\mathcal{O}}_{C^{\nu}, c}$ as well. We will start with some algebra facts.

Lemma 3.13. *Let $\phi : (A, \mathfrak{m}_A) \rightarrow (B, \mathfrak{m}_B)$ be a finite homomorphism of pro- $\mathbf{T}(0)$ -MHS-algebras, with $(A, \mathfrak{m}_A)$ and $(B, \mathfrak{m}_B)$ complete local domains. Assume further that*

$$\mathfrak{m}_A = W_{-1}A, \quad W_0A = A$$

then $W_0B = B$ and $\mathfrak{m}_B = W_{-1}B$ as well.

Proof. We have the functorial Deligne splitting [PS08, Lemma-Definition 3.4] for the weight filtrations on A on B , giving $A \simeq \prod_{w \in \mathbb{Z}_{\leq 0}} A_w$, $B \simeq \prod_{w \in \mathbb{Z}} B_w$, and a finite map

$$\bar{\phi} : \prod_{w \in \mathbb{Z}_{\leq 0}} A_w \rightarrow \prod_{w \in \mathbb{Z}} B_w$$

compatible with the weights. Let $0 \neq b \in B_w$ be a homogeneous element, then b is integral, say of degree d , over $\prod_{w \in \mathbb{Z}_{\leq 0}} A_w$. Take the dw -homogeneous part of the monic polynomial for b , we get

$$b^d + a_1 b^{d-1} + \cdots + a_d = 0$$

with $a_j \in A_{jw}$. If $w > 0$ then $A_{jw} = 0$ hence $a_j = 0$ and $b^d = 0$ which gives a contradiction since A is a domain. If $w = 0$ then all $a_j \in A_0 = \mathbb{C}$ hence $b \in \mathbb{C}$. It follows that $W_{-1}B = \mathfrak{m}_B$ and $W_0B = B$. $\square$

Lemma 3.14. *Let B be a complete noetherian local pro- $\mathbf{T}(0)$ -MHS-algebra with $W_{-1}B = \mathfrak{m}_B$. Then the ideal of the reduced singular locus $I(\text{Sing}(\text{Spec } B)_{\text{red}})$ is a pro- $\mathbf{T}(0)$ -MHS-ideal of B .*

Proof. This follows from the proof of [BBT24, Lemma 8.23] which implies that $I(\text{Sing}(\text{Spec } B))$ is a pro- $\mathbf{T}(0)$ -MHS-ideal of B , and [BBT24, Lemma 4.36] which says that taking the reduction preserves the Hodge structure. $\square$

Lemma 3.15. *Let $(B, \mathfrak{m}_B)$ be a pro- $\mathbf{T}(0)$ -MHS-algebra which is a complete noetherian local domain, and $I \subset B$ a pro- $\mathbf{T}(0)$ -MHS-ideal. Then $\text{End}_B(I)$ underlies a local pro- $\mathbf{T}(0)$ -MHS-algebra structure such that the natural map defined by multiplication*

$$B \rightarrow \text{End}_B(I)$$

is an finite injective morphism of pro- $\mathbf{T}(0)$ -MHS-algebras.

Proof. Write

$$B = \varprojlim B/\mathfrak{m}_B^{n+1} = \varprojlim B_n, \quad I = \varprojlim I/\mathfrak{m}_B^{n+1}I = \varprojlim I_n$$

then each $\text{End}_{B_n}(I_n)$ is just the kernel of the morphism of $\mathbb{C}$ -MHS

$$\text{End}_{\mathbb{C}}(I_n) \rightarrow \text{Hom}_{\mathbb{C}}(B_n \otimes_{\mathbb{C}} I_n, I_n)$$

hence carries the structure of an artinian local $\mathbf{T}(0)$ -MHS-algebra. Next, we have tensor-Hom adjunction

$$\text{Hom}_{\text{MHS}_{\mathbb{C}}}(B_n \otimes I_n, I_n) \simeq \text{Hom}_{\text{MHS}_{\mathbb{C}}}(B_n, \text{End}_{\mathbb{C}}(I_n))$$

under which the multiplication map $B_n \rightarrow \text{End}_{\mathbb{C}}(I_n)$ corresponds to the B_n -module structure map $B_n \otimes I_n \rightarrow I_n$ which is a morphism of $\mathbb{C}$ -MHS since I_n is a $\mathbf{T}(0)$ -MHS-ideal of B_n . Hence $B_n \rightarrow \text{End}_{\mathbb{C}}(I_n)$ is a morphism of $\mathbb{C}$ -MHS which factors through $B_n \rightarrow \text{End}_{B_n}(I_n)$ in $\text{MHS}_{\mathbb{C}}$. The kernel is $\text{Ann}_{B_n}(I_n) = 0$ since B is a domain.

The map $\text{End}_B(I) \rightarrow \text{End}_{B_n}(I_n)$ is not surjective, and with kernel

$$K_n = \{f \in \text{End}_B(I) \mid f(I) \subseteq \mathfrak{m}_B^{n+1}I\}$$

Let $\langle x_1, \dots, x_l \rangle$ be a generating set of I as a B -module. Then the map

$$\text{End}_B(I) \rightarrow I^l, \quad f \mapsto (f(x_1, x_2, \dots, x_l))$$

is an injection, and gives $K_n = \text{End}_B(I) \cap \mathfrak{m}_B^{n+1}I^l$. By [AM69, Theorem 10.11], the two filtrations $\mathfrak{m}_B^{\bullet+1}\text{End}_B(I)$ and $K_{\bullet}$ are cofinal. Thus we can take the inverse limit of the first part of the proof to get the desired statements. $\square$

Proposition 3.16. *Let Λ be a reduced complete noetherian local $\mathbb{C}$ -algebra which underlies a pro- $\mathbf{T}(0)$ -MHS-algebra such that $\mathfrak{m}_{\Lambda} = W_{-1}\Lambda$. For every minimal prime $\mathfrak{p} \subset \Lambda$, denote $A = \Lambda/\mathfrak{p}$ and A^{ν} its normalization. Then A^{ν} underlies a pro- $\mathbf{T}(0)$ -MHS-algebra structure with*

$$W_0 A^{\nu} = A^{\nu}, \quad W_{-1} A^{\nu} = \mathfrak{m}_{\nu}$$

such that the normalization map $\nu : A \rightarrow A^{\nu}$ and its composition $\Lambda \rightarrow A^{\nu}$ are morphisms of pro- $\mathbf{T}(0)$ -MHS-algebras.

Proof. We will make use Oka's criterion for normality [GR84, Chapter 6 §5.2], which says that for a reduced ring B , B is normal if and only if the multiplication map

$$B \rightarrow \text{End}_B(I(\text{Sing}(\text{Spec } B)_{\text{red}}))$$

is an isomorphism. Start with $B^0 = A = \Lambda/\mathfrak{p}$ which is a domain, and define recursively

$$B^{j+1} = \text{End}_{B^j}(I(\text{Sing}(\text{Spec } B^j)_{\text{red}}))$$

then each $B^{j+1} \subseteq A^{\nu}$ and is finite. We will prove this by induction. Denote $I^j = I(\text{Sing}(\text{Spec } B^j)_{\text{red}})$. Suppose $B^j \subseteq A^{\nu}$ is a domain and $\text{Frac}(B^j) = \text{Frac}(A)$, which is automatically satisfied for $j = 0$. Pick $0 \neq b \in I^j$, then for

$$f \in \text{End}_{B^j}(I^j)$$

$f(b)/b \in \text{Frac}(A)$. We get that $f(b') = \frac{f(b)}{b}f(b')$ for all $b' \in I^j$, which gives an embedding $\text{End}_{B^j}(I^j) \hookrightarrow \text{Frac}(A)$ by $f \mapsto f(b)/b$. Thus we get

$$B^j \subseteq \text{End}_{B^j}(I^j) = B^{j+1} \subseteq \text{Frac}(A)$$

where the first inclusion is by the multiplication map and the fact that B^j is a domain. Finally B^{j+1} is finite over B^j hence $B^{j+1} \subseteq A^{\nu}$ as well. We have a chain

$$A \subseteq B^1 \subseteq B^2 \subseteq \dots \subseteq A^{\nu}$$

which must stabilize due to noetherian assumption. At the stable stage we get $B^j \simeq \text{End}_{B^j}(I^j)$ hence $B^j = (B^j)^{\nu} = A^{\nu}$.

By [BBT24, Lemma 4.36], $\Lambda \rightarrow A = \Lambda/\mathfrak{p}$ is a morphism of pro- $\mathbf{T}(0)$ -MHS-algebras. By Lemma 3.14 and Lemma 3.15, the inclusions

$$A \subseteq B^1 \subseteq B^2 \subseteq \cdots \subseteq A^\nu$$

are all morphisms of pro- $\mathbf{T}(0)$ -MHS-algebras. It follows that $A \rightarrow A^\nu$ is a morphism of pro- $\mathbf{T}(0)$ -MHS-algebras, which is a finite map hence by Lemma 3.13 we get the weight statements. $\square$

Corollary 3.17. *Using the notation of Section 2.5, let Z be an irreducible component of the reduced image $\bar{\Sigma}_k^i(X, 2) \subseteq M_B(X, 2)$. Let C be an irreducible component of $\phi_\chi^{-1}(Z) \subset M_B(X, 2, \det = \chi) \times A$, and $(\mathbb{V}, \mathbb{L}) \in C$ a $\mathbb{C}$ -VHS point. Consider the normalization $\nu : C^\nu \rightarrow C$ with c lying over $(\mathbb{V}, \mathbb{L})$, then $\hat{\mathcal{O}}_{C^\nu, c}$ underlies a pro- $\mathbf{T}(0)$ -MHS-algebra with*

$$W_0 \hat{\mathcal{O}}_{C^\nu, c} = \hat{\mathcal{O}}_{C^\nu, c}, \quad W_{-1} \hat{\mathcal{O}}_{C^\nu, c} = \mathfrak{m}_{C^\nu, c}$$

such that the map $\hat{\mathcal{O}}_{C, (\mathbb{V}, \mathbb{L})} \rightarrow \hat{\mathcal{O}}_{C^\nu, c}$ is a morphism of pro- $\mathbf{T}(0)$ -MHS-algebras.

Proof. Any such $\hat{\mathcal{O}}_{C^\nu, c}$ would be isomorphic to $\left(\hat{\mathcal{O}}_{C, (\mathbb{V}, \mathbb{L})}/\mathfrak{p}\right)^\nu$ for some minimal prime $\mathfrak{p}$. Proposition 3.16 then gives the desired statements. $\square$

4. VIRTUALLY ABELIAN JUMP COMPONENTS

Let $\Sigma_k^i(X, 2) \subseteq R_B(X, x, 2)$ be a jump locus, and $\bar{\Sigma}_k^i(X, 2) \subseteq M_B(X, 2)$ be the reduced good moduli image. Denote $Z \subseteq \bar{\Sigma}_k^i(X, 2)$ an irreducible component. We first justify labeling Z based on its general point.

4.1. Generic projective type. Consider the following algebraic subgroups of PGL_2 :

$$B = \mathrm{Stab}(\infty) = \left\{ \left[\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \right] \mid ad \neq 0 \right\} \quad (4.1)$$

$$T_{0, \infty} = \mathrm{Stab}(\infty) \cap \mathrm{Stab}(0) = \left\{ \left[\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \right] \mid ad \neq 0 \right\} \quad (4.2)$$

$$D_\infty = T_{0, \infty} \cup \left\{ \left[\begin{pmatrix} 0 & b \\ c & 0 \end{pmatrix} \right] \mid bc \neq 0 \right\} \quad (4.3)$$

where square bracket denotes the projective class. Notice that $T_{0, \infty}$ is an algebraic torus, D_∞ is the normalizer of $T_{0, \infty}$, fixes the set of points $\{0, \infty\} \subset \mathbb{P}^1$ and is $D_\infty \simeq T_{0, \infty} \rtimes \mathbb{Z}/2\mathbb{Z}$.

Let $\rho : \pi_1(X, x) \rightarrow \mathrm{GL}_2(\mathbb{C})$ be a semisimple representation, and $\bar{\rho} : \pi_1(X, x) \rightarrow \mathrm{PGL}_2(\mathbb{C})$ its projective quotient. Denote the Zariski closure

$$G_{\bar{\rho}} := \overline{\rho(\pi_1(X, x))}^{\mathrm{Zar}} \subseteq \mathrm{PGL}_2$$

the algebraic monodromy group of $\bar{\rho}$.

Lemma 4.1. *For each semisimple $\rho : \pi_1(X, x) \rightarrow \mathrm{GL}_2(\mathbb{C})$, exactly one of the following is true:*

(A): ρ is reducible, i.e., $\rho \simeq \chi_1 \oplus \chi_2$;

- (B): ρ is irreducible, and, up to conjugation, $G_{\bar{\rho}} \subseteq D_{\infty}$;
 (C): $G_{\bar{\rho}} \simeq \bar{\rho}(\pi_1(X, x)) \simeq A_4, S_4$ or A_5 ;
 (D): $\bar{\rho}$ has Zariski dense image and is rigid;
 (E): $\bar{\rho}$ has Zariski dense image and is non-rigid.

Proof. ρ is reducible if and only if it fixes a line, or equivalently, $G_{\bar{\rho}}$ fixes a point in $\mathbb{P}^1$. Up to conjugation, this is equivalent to $G_{\bar{\rho}}$ being contained in the Borel B . The fact that this classification is exhaustive follows from [NPT08, Theorem 1].

For mutual exclusivity, it is clear that (A), (B), (D), and (E) are mutually exclusive; same for (C), (D) and (E). For (B), and (C), notice that the only finite subgroups of $D_{\infty} = T_{0,\infty} \rtimes \mathbb{Z}/2\mathbb{Z}$ is either dihedral D_n or a cyclic group. Finally for (A) and (E), a finite subgroup fixing a point must fix another (it acts as an affine transformation on $\mathbb{A}^1$), hence, up to conjugation, is contained in the torus $T_{0,\infty}$. It follows that it must be cyclic. $\square$

We will show that a general point of an irreducible component must be of exactly one of the previous 5 types. In other words, we cannot have two types whose points are both dense in the component. It suffices to show that each type is a locally closed condition.

Definition 4.2. Denote

$$\begin{aligned} \mathfrak{D} &\subset M_B(X, 2) \text{ be } \bigcup_{0 \neq \epsilon \in \text{Hom}(\pi_1, \mathbb{Z}/2\mathbb{Z})} \mathfrak{D}_{\epsilon} \text{ as in Lemma 4.6,} \\ \mathfrak{R} &\subset M_B(X, 2) \text{ the locus of reducible representations,} \\ \mathfrak{F} &\subset M_B(X, 2) \text{ the locus of representations of type (C),} \\ \mathfrak{G} &\subset M_B(X, 2) \text{ the locus of projectively dense representations.} \end{aligned}$$

then $M_B(X, 2) = \mathfrak{R} \cup \mathfrak{D} \cup \mathfrak{F} \cup \mathfrak{G}$.

Proposition 4.3. *Let Z be an irreducible component of $\bar{\Sigma}_k^i(X, 2)$, the reduced good moduli image of $\Sigma_k^i(X, 2) \subseteq R_B(X, x, 2)$. Then Z has a nonempty Zariski open subset on which exactly one of the types in Lemma 4.1 occurs. If the type is (A), (B), or (C), then Z is contained in, respectively, $\mathfrak{R}$, $\mathfrak{D}$ or $\mathfrak{F}$.*

Proof. The locus of type (A) is $\mathfrak{R}$, which is closed by Lemma 4.4. The locus of type (C) is closed by (4.7). The locus of type (B) is open in $\mathfrak{D}$ by Lemma 4.6. Type (D) locus is closed in $\mathfrak{G}$, since it is a finite union of images of θ_{ρ} as defined in Lemma 4.8; this is because there are only finitely many rigid points in $M_B(X, \text{PGL}_2)$. Finally the locus of type (E) is open in $\mathfrak{G}$ hence open. $\square$

4.2. Reducible jump component.

Lemma 4.4. *Let $A = M_B(X, 1)$, then the map*

$$\iota : (A \times A)/S_2 \rightarrow M_B(X, 2), \quad (\chi_1, \chi_2) \mapsto [\chi_1 \oplus \chi_2] \quad (4.4)$$

is a closed embedding, with image the reducible locus $\mathfrak{R}$.

Proof. These are both affine schemes, and the corresponding map of rings is

$$\iota^\# : H^0(R_B(X, x, 2), \mathcal{O}_{R_B})^{\mathrm{GL}_2} \rightarrow (H^0(A, \mathcal{O}_A) \otimes H^0(A, \mathcal{O}_A))^{S_2}$$

mapping

$$f \mapsto \left[(\chi_1, \chi_2) \mapsto f \left(\gamma \mapsto \begin{pmatrix} \chi_1(\gamma) & 0 \\ 0 & \chi_2(\gamma) \end{pmatrix} \right) \right]$$

where f is a function on $\mathrm{Hom}(\pi_1(X, x), \mathrm{GL}_2)$. Notice that $H^0(A, \mathcal{O}_A) \otimes H^0(A, \mathcal{O}_A)$ is generated by the functions

$$\gamma^1 : (\chi_1, \chi_2) \mapsto \chi_1(\gamma), \quad \gamma^2 : (\chi_1, \chi_2) \mapsto \chi_2(\gamma) \quad \gamma \in \pi_1(X, x)$$

i.e., every regular function on $A \times A$ has the form $\sum \gamma^1 \delta^2$ with $\gamma, \delta \in \pi_1(X, x)$. Let $1 \neq \sigma \in S_2$, then $\sigma^*(\gamma^1 \delta^2) = \delta^1 \gamma^2$. The S_2 -invariant functions are then generated by

$$\gamma^1 \delta^2 + \delta^1 \gamma^2 \quad \delta, \gamma \in \pi_1(X, x)$$

Next, notice that the trace functionals

$$\mathrm{tr}_\gamma : R_B(X, x, 2) \rightarrow \mathbb{C}, \quad \rho \mapsto \mathrm{tr}(\rho(\gamma))$$

are GL_2 -invariant and satisfies $\iota^\#(\mathrm{tr}_\gamma) = \gamma^1 + \gamma^2$, hence

$$\iota^\#(\mathrm{tr}_\gamma \mathrm{tr}_\delta - \mathrm{tr}_{\gamma\delta}) = \gamma^1 \delta^2 + \delta^1 \gamma^2$$

thus $\iota^\#$ is a quotient map. As a result, ι is a closed embedding. $\square$

Let $q : A \times A \rightarrow (A \times A)/S_2$ be the quotient. Since $H^i(X, \chi_1 \oplus \chi_2) = H^i(X, \chi_1) \oplus H^i(X, \chi_2)$, we get that

$$\bar{\Sigma}_k^i(X, 2) \cap \mathfrak{R} = \iota \left(\bigcup_{a+b=k} \Sigma_a^i(X, 1) \times \Sigma_b^i(X, 1) \right) \quad (4.5)$$

which is a finite union of images under q of products of torsion-translated subtori, by [Corollary 2.17](#).

Proposition 4.5. *Let Z be an irreducible component of $\bar{\Sigma}_k^i(X, 2)$, the reduced good moduli image of $\Sigma_k^i(X, 2) \subseteq R_B(X, x, 2)$. If the general point of Z is reducible, then there is a product $N = N_1 \times N_2 \subseteq M_B(X, 1) \times M_B(X, 1)$ of torsion-translated subtori such that*

$$\begin{cases} Z^\nu \simeq N_1 \times N_2 & \text{if } N_1 \neq N_2 \\ Z \simeq \mathrm{Sym}^2(N_1) & \text{if } N_1 = N_2 \end{cases}$$

Proof. Since $\mathfrak{R}$ is closed, $Z \subset \mathfrak{R}$. Z is an irreducible component, hence it picks out one component $N = \Sigma_a^i(X, 1) \times \Sigma_b^i(X, 1)$ in (4.5). It follows that $Z = \iota(q(N))$ hence $Z \simeq q(N)$.

If $N_1 \neq N_2$ then $N_1 \cap \sigma N_2$ is closed subset of N_1 , for $1 \neq \sigma \in S_2$. Therefore the map $N_1 \times N_2 \rightarrow q(N_1 \times N_2)$ is finite, surjective with generic degree 1. It follows that it must be the normalization. If $N_1 = N_2$ then $q(N_1 \times N_1) \simeq \mathrm{Sym}^2(N_1)$. $\square$

4.3. Dihedral jump component. If, up to conjugation, $G_{\bar{\rho}} \subseteq D_{\infty}$ then ρ fixes a set of two lines $\{l_1, l_2\}$ with $l_1 \oplus l_2 \simeq \mathbb{C}^2$. Each $\rho(\gamma)(l_i) \in \{l_1, l_2\}$ hence we get a permutation character

$$\epsilon : \pi_1(X, x) \rightarrow \mathbb{Z}/2\mathbb{Z}$$

If $\epsilon \neq 0$ (equivalently, ρ is of type **(B)**), $\rho|_{\ker \epsilon}$ then fixes both line hence $\rho|_{\ker \epsilon} \simeq \chi_1 \oplus \chi_2$. An element $\delta \in \pi_1(X, x) \setminus \ker \epsilon$ has nontrivial $\epsilon(\delta)$. Conjugation by δ on $\ker \epsilon \trianglelefteq \pi_1(X, x)$ then permutes χ_1 and χ_2 .

Now pick any $0 \neq \epsilon \in \text{Hom}(\pi_1(X, x), \mathbb{Z}/2\mathbb{Z})$, and denote

$$A^{\epsilon} := \text{Hom}(\ker \epsilon, \mathbb{G}_m)$$

then conjugation by any $\delta \in \pi_1(X, x) \setminus \ker \epsilon$ on $\ker \epsilon \triangleleft \pi_1(X, x)$, denoted $\tau : \ker \epsilon \rightarrow \ker \epsilon$, induces an involution τ^* on A^{ϵ} . This is because $\delta^2 = 0$; the choice of δ does not matter.

Lemma 4.6. *Induction gives a morphism*

$$\iota_{\epsilon} : A^{\epsilon} / \langle \tau^* \rangle \rightarrow M_B(X, 2), \quad [\chi] \mapsto [\text{Ind}_{\ker \epsilon}^{\pi_1(X, x)} \chi]$$

which is a closed embedding with image $\mathfrak{D}_{\epsilon}$. The irreducible points in the image correspond to representations of type **(B)** with the permutation character ϵ , and form a Zariski open subset of $\mathcal{D}_{\epsilon}$.

Proof. The corresponding map on rings is

$$\iota_{\epsilon}^{\#} : H^0(R_B(X, x, 2), \mathcal{O}_{R_B})^{\text{GL}_2} \rightarrow H^0(A^{\epsilon}, \mathcal{O}_{A^{\epsilon}})^{\tau^*}$$

mapping $f \mapsto \left[\chi \mapsto f \left(\text{Ind}_{\ker \epsilon}^{\pi_1(X, x)} \chi \right) \right]$. Explicitly, with the same δ as before, and $\rho = \text{Ind}_{\ker \epsilon}^{\pi_1(X, x)} \chi$,

$$\begin{aligned} \rho(\gamma) &= \begin{pmatrix} \chi(\gamma) & 0 \\ 0 & \chi(\delta^{-1}\gamma\delta) \end{pmatrix}, \quad \gamma \in \ker(\epsilon) \\ \rho(\delta') &= \begin{pmatrix} 0 & \chi(\delta h \delta) \\ \chi(h) & 0 \end{pmatrix}, \quad \delta' \in \pi_1(X, x) \setminus \ker \epsilon, \delta' = \delta \cdot h \end{aligned}$$

Now, regular functions on A^{ϵ} are generated by those of the form $\gamma : \chi \mapsto \chi(\gamma)$ with $\gamma \in \ker(\epsilon)$. Thus τ^* -invariant functions are generated by

$$\gamma^{\tau} : \chi \mapsto \chi(\gamma) + \chi(\delta^{-1}\gamma\delta), \quad \gamma \in \ker(\epsilon)$$

which are clearly $\iota_{\epsilon}^{\#}(\text{tr}_{\gamma})$ hence $\iota_{\epsilon}^{\#}$ is a quotient map.

Notice that $\iota_{\epsilon}(\chi)|_{\ker \epsilon} = \chi \oplus \tau^* \chi$. If $\chi \neq \tau^* \chi$ then the only possible invariant line must be generated by either basis vectors, i.e., $\mathbb{C} \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ or $\mathbb{C} \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. However, any element in $\pi_1(X, x) \setminus \ker \epsilon$ swaps these two lines, thus there is no invariant line and ι_{ϵ} is of type **(B)**. By the discussion at the beginning of the section, this gives all type **(B)** representations with permutation character ϵ .

Now if $\chi = \tau^* \chi$ then for $\gamma \in \ker \epsilon$ and $\delta' = \delta \cdot h \in \pi_1(X, x) \setminus \ker \epsilon$,

$$\iota_{\epsilon}(\chi)(\gamma) = \chi(\gamma) \cdot \text{Id}, \quad \iota_{\epsilon}(\delta') = \chi(h) \begin{pmatrix} 0 & \chi(\delta^2) \\ 1 & 0 \end{pmatrix}$$

then the eigendecomposition for $\iota_\epsilon(\delta')$, which is independent of h hence independent of $\delta' \in \pi_1(X, x) \setminus \ker \epsilon$, gives a decomposition of $\iota_\epsilon(\chi)$. Finally, the set of χ fixed by $\tau^*(\chi)$ is closed, hence we get the Zariski open statement. $\square$

For $0 \neq \epsilon \in \text{Hom}(\pi_1(X, x), \mathbb{Z}/2\mathbb{Z})$, let $\pi^\epsilon : X^\epsilon \rightarrow X$ be the connected étale double cover coming from $\ker \epsilon \triangleleft \pi_1(X, x)$. For $\chi \in A^\epsilon$ with the corresponding local system $\mathbb{L}$ on X^ϵ , $\text{Ind}_{\ker \epsilon}^{\pi_1(X, x)} \chi$ corresponds to the local system $\pi_*^\epsilon \mathbb{L}$ on X . By a straightforward application of Leray's spectral sequence,

$$H^i(X, \text{Ind}_{\ker \epsilon}^{\pi_1(X, x)} \chi) \simeq H^i(X^\epsilon, \chi) \quad (4.6)$$

and we get the following consequence for a type **(B)** jump component.

Proposition 4.7. *Let Z be an irreducible component of $\bar{\Sigma}_k^i(X, 2)$, the reduced good moduli image of $\Sigma_k^i(X, 2) \subseteq R_B(X, x, 2)$. If the general point of Z is of type **(B)**, then there exists a torsion-translated subtorus $N \subseteq M_B(X^\epsilon, 1)$ such that either $Z^\nu \simeq N$ or, if $\tau^* N = N$, $Z \simeq N / \langle \tau^* \rangle$.*

Proof. Since Z is irreducible, and its general point is of type **(B)**, it must lie in some $\mathfrak{D}_\epsilon$ in Lemma 4.6. By (4.6), it must be the image of an irreducible component N of $\Sigma_k^i(X^\epsilon, 1)$, hence N is a torsion-translated subtorus by Corollary 2.17. If N is τ^* -invariant, then since ι_ϵ is a closed embedding, $Z \simeq N / \langle \tau^* \rangle$. If N is not τ^* -invariant, then $N \cap \tau^* N$ is closed in N , thus the composition $N \rightarrow A^\epsilon / \langle \tau^* \rangle \xrightarrow{\iota_\epsilon} Z$ is a finite map of generic degree 1, hence $N \simeq Z^\nu$. $\square$

4.4. Projectively finite and rigid jump components. Let $\mathbb{V}_\rho \in M_B(X, 2)$ be an irreducible semisimple representation with corresponding representation $\rho : \pi_1(X, x) \rightarrow \text{GL}_2(\mathbb{C})$. Let $A = M_B(X, 1)$.

Lemma 4.8. *The morphism*

$$\theta_\rho : A \rightarrow M_B(X, 2), \quad \chi \mapsto [\rho \otimes \chi]$$

is finite, with fibers being the orbits of the finite group

$$K(\rho) = \{ \chi \in A \mid \rho \otimes \chi \simeq \rho \} \subseteq A[2]$$

Proof. Consider the diagram

$$\begin{array}{ccc} A & \xrightarrow{\chi \mapsto \rho \otimes \chi} & M_B(X, 2) \\ \chi \mapsto (\rho \otimes \chi, \chi) \searrow & & \uparrow \\ M_B(X, 2) \times_A A & \xrightarrow{\Phi} & M_B(X, 2) \\ \downarrow & & \downarrow \det \\ A & \xrightarrow{\chi \mapsto \det(\rho) \otimes \chi^{\otimes 2}} & A \end{array}$$

then $A \rightarrow M_B(X, 2) \times_A A$ is a closed embedding, and Φ is finite since the horizontal $A \rightarrow A$ is finite. Thus θ_ρ , being the composition, is finite. For the fiber identification, notice that $\rho \otimes \chi \simeq \rho \otimes \chi'$ if and only if $\chi' \otimes \chi^{-1} \in K(\rho)$. $\square$

Let $\Gamma \in \{A_4, S_4, A_5\}$. Since $\pi_1(X, x)$ is finitely generated and Γ is finite, there are only finitely many homomorphism $\varphi : \pi_1(X, x) \rightarrow \Gamma \subset \mathrm{PGL}_2(\mathbb{C})$. Since every subgroup of PGL_2 isomorphic to Γ is conjugate to the standard embedding, and we are working with M_B , there is no ambiguity with φ . Suppose φ has a GL_2 -lift $\mathbb{V}_{\Gamma, \varphi}$. Since any two lifts differ by a rank one twist, we get that

$$\mathfrak{F} = \bigcup_{\Gamma, \varphi} \theta_{\mathbb{V}_{\Gamma, \varphi}}(A) \subset M_B(X, 2) \quad (4.7)$$

with $\mathfrak{F}$ the locus of type **(C)** representations.

Proposition 4.9. *Let Z be an irreducible component of $\overline{\Sigma}_k^i(X, 2)$, the reduced good moduli image of $\Sigma_k^i(X, 2)$. If the general point of Z is of type **(C)**, then there is a torsion-translated $N \subset A$ such that Z has a normalization*

$$Z^\nu = N/H$$

where $H = \{\chi \in K(\rho) \mid \chi N = N\}$. Consequently, Z^ν is a torsion-translated subtorus of A/H .

Proof. Since Z is irreducible, it must lie in some $\theta_{\mathbb{V}_{\Gamma, \varphi}}(A)$, with $\mathbb{V}_{\Gamma, \varphi}$ a GL_2 -lift of some $\varphi : \pi_1(X, x) \rightarrow \mathrm{PGL}_2(\mathbb{C})$ with image Γ . We first claim that it is possible to pick $\mathbb{V}_{\Gamma, \varphi}$ with finite monodromy. Equivalently, there is some $\chi \in A$ such that $\chi \otimes \mathbb{V}_{\Gamma, \varphi}$ has finite monodromy. Recall that every class in A/A^2 has a torsion lift in A , and pick one such lift ϵ for $[\det \mathbb{V}_{\Gamma, \varphi}] \in A/A^2$. We get that for some $\mathbb{L} \in A$, $\epsilon = \det(\mathbb{V}_{\Gamma, \varphi}) \otimes \mathbb{L}$. Then

$$\det(\mathbb{V}_{\Gamma, \varphi} \otimes \mathbb{L}) = e$$

is torsion. It follows that $\mathbb{V}_{\Gamma, \varphi} \otimes \mathbb{L}$ has finite monodromy projective quotient, and finite monodromy determinant. Since the map

$$\mathrm{GL}_2 \rightarrow \mathrm{PGL}_2 \times \mathbb{G}_m, \quad g \mapsto ([g], \det g)$$

is finite of degree 2, we get that $\mathbb{V}_{\Gamma, \varphi} \otimes \mathbb{L}$ has finite monodromy. In particular, $\mathbb{V}_{\Gamma, \varphi} \otimes \mathbb{L}$ is of geometric origin.

We can now assume that $\mathbb{V}_{\Gamma, \varphi}$ is of geometric origin. Then $\theta_{\mathbb{V}_{\Gamma, \varphi}}^{-1}(Z)$ is an irreducible component of the twisted jump loci

$$\Sigma_k^i(X, \mathbb{V}_{\Gamma, \varphi}, 1) = \{\chi \in A \mid \dim H^i(X, \mathbb{V}_{\Gamma, \varphi} \otimes \chi) \geq k\}$$

which is then a torsion-translated subtorus $N \subset A$ by [Corollary 2.17](#). It follows that $Z = \theta_{\mathbb{V}_{\Gamma, \varphi}}(N)$. Next, denote $H = \{\chi \in K(\rho) \mid \chi N = N\}$, then since H acts by translation, N/H is a translated subtorus of A/H . Finally, $N/H \rightarrow \theta_{\mathbb{V}_{\Gamma, \varphi}}(N)$ is finite of degree 1 (since the extra identifications come from the fixed loci of $\chi \in K(\rho)$ with $\chi N \neq N$; these fixed loci are closed proper subset of N). As a result, $N/H \rightarrow Z$ is the normalization map. $\square$

Proposition 4.10. *Let Z be an irreducible component of $\overline{\Sigma}_k^i(X, 2)$, the reduced good moduli image of $\Sigma_k^i(X, 2)$. If the general point of Z is of type **(D)**. Then there exists a translated subtorus $N \subseteq A$ such that $Z^\nu \simeq N/H$ with $\{\chi \in K(\rho) \mid \chi N = N\}$.*

Proof. Here we must have $Z \subseteq \theta_{\mathbb{V}_{\Gamma, \varphi}}(A)$, with $\mathbb{V}_{\Gamma, \varphi}$ a GL_2 -lift of an isolated point in $M_B(X, \mathrm{PGL}_2)$. Hence it is the image under $\theta_{\mathbb{V}_{\Gamma, \varphi}}(A)$ of an irreducible component of

$$\Sigma_k^i(X, \mathbb{V}_{\Gamma, \varphi}, 1) = \{\chi \in A \mid \dim H^i(X, \mathbb{V}_{\Gamma, \varphi} \otimes \chi) \geq k\}$$

hence it is $\theta_{\mathbb{V}_{\Gamma, \varphi}}(N)$ with $N \subseteq A$ a translated subtorus by [Corollary 2.17](#) (we lose that the translation is torsion). The remaining statements are exactly as in the proof of [Proposition 4.9](#). $\square$

5. NONABELIAN JUMP COMPONENT

This section will handle the remaining case from the previous section, i.e., Z is an irreducible component of $\bar{\Sigma}_k^i(X, 2)$ with general point of type **(E)** (see [Lemma 4.1](#)). Let $A = M_B(X, 1)$. Following [Section 2.5](#), Z is contained in some $M_B(X, 2, \det \in \epsilon A^2)$ with torsion $\epsilon \in A$. Pick an irreducible component

$$\begin{array}{ccc} C \subseteq \Phi_\epsilon^{-1}(Z) \subseteq M_B(X, 2, \det = \epsilon) \times A & & \\ \text{pr}_1 \downarrow & \searrow \text{pr}_2 & \\ M_B(X, 2, \det = \epsilon) & & A \end{array} \quad (5.1)$$

then by [Proposition 2.19](#), the normalized $C^\nu \rightarrow Z^\nu$ is a finite étale torsor.

First we take a detour to talk about local systems and cohomology on a Deligne-Mumford quotient stack.

5.1. Local systems on finite quotient curves. We are following the exposition in [\[CS08, §2.1 and 2.2\]](#). For more details, see [\[Noo04\]](#).

Definition 5.1. A *DM curve* is a connected smooth one-dimensional complex Deligne-Mumford stack, and an *orbicurve* is a DM curve with trivial generic stabilizer.

In this paper, we work only with DM curves $\mathcal{Y}$ of the form $\mathcal{Y} = [Y/G]$ where Y is a connected smooth projective curve and G is a finite group acting on Y .

Definition 5.2. Let $\bar{y} \in \mathcal{Y}$ be a point, then we have the fundamental group $\pi_1(\mathcal{Y}, \bar{y})$. For $y \in Y$ lying over $\bar{y} \in \mathcal{Y}$ in the étale map

$$\pi : Y \rightarrow \mathcal{Y} = [Y/G]$$

an element of $\pi_1(\mathcal{Y}, \bar{y})$ is a pair (g, γ) with $g \in G$ and γ a path from y to $g \cdot y$. Thus we have an exact sequence

$$1 \rightarrow \pi_1(Y, y) \rightarrow \pi_1(\mathcal{Y}, \bar{y}) \rightarrow G \rightarrow 1$$

Definition 5.3. A complex local system on $\mathcal{Y}$ is a G -equivariant local system on Y , i.e. a pair $(\mathbb{V}, \alpha)$ where $\mathbb{V}$ is a local system on Y and α is an action associating to each $g \in G$ an isomorphism $\alpha_g : g^* \mathbb{V} \xrightarrow{\sim} \mathbb{V}$ such that

$$\alpha_g \circ g^* \alpha_h = \alpha_{gh}, \quad g, h \in G$$

Then we recover the familiar equivalence [\[Noo05, Theorem 18.19\]](#)

$$\mathrm{Loc}^G(Y) \simeq \mathrm{Loc}_{\mathbb{C}}(\mathcal{Y}) \simeq \mathrm{Rep}_{\mathbb{C}}(\pi_1(\mathcal{Y}, \bar{y}))$$

between complex local systems on $\mathcal{Y}$ and complex representations of $\pi_1(\mathcal{Y}, \bar{y})$.

We have similar definitions for constructible complexes and perverse sheaves, i.e. $D_c^b(\mathcal{Y}, \mathbb{C}) = D_c^{b,G}(Y, \mathbb{C})$ and $\text{Perv}(\mathcal{Y}) = \text{Perv}^G(Y)$. For $K \in D_c^b(\mathcal{Y}, \mathbb{C})$, $\text{R}\Gamma(\mathcal{Y}, K) \simeq \text{R}\Gamma(Y, \pi^* K)^G$.

Definition 5.4. Since $\text{Loc}_{\mathbb{C}}(\mathcal{Y}) \simeq \text{Rep}_{\mathbb{C}}(\pi_1(\mathcal{Y}, \bar{y}))$, we have framed character varieties of $\mathcal{Y}$,

$$R_B(\mathcal{Y}, \bar{y}, r) := \text{Hom}(\pi_1(\mathcal{Y}, \bar{y}), \text{GL}_r)$$

which are affine schemes. Its affine good moduli space is the GIT quotient

$$M_B(\mathcal{Y}, r) := R_B(\mathcal{Y}, \bar{y}, r) // \text{GL}_r = \text{Spec } H^0(R_B(\mathcal{Y}, \bar{y}, r), \mathcal{O}_{R_B})^{\text{GL}_r}$$

by the conjugation action.

For $y \in Y$ let $G_y = \{g \in G \mid g \cdot y = y\}$ be the stabilizer group. Notice that replacing y by $g \cdot y$ identifies G_y with G_{gy} by conjugation, which then gives a canonical isomorphism

$$M_B(BG_y, r) \simeq M_B(BG_{gy}, r)$$

where $BH = [\text{Spec } \mathbb{C}/H]$. Denote $I_\eta = \ker(G \rightarrow \text{Aut}(Y))$ the generic inertia, then for general $y \in Y$, $G_y = I_\eta$. There are finitely many exceptional points $y \in Y$ with $I_\eta \subsetneq G_y$; pick their G -orbit representatives $y_1, \dots, y_l$ with $G_j = G_{y_j}$. Pulling back along the residual gerbes $\iota_j : BG_j \rightarrow \mathcal{Y}$ gives

$$\iota_j^* : M_B(\mathcal{Y}, r) \rightarrow M_B(BG_j, r), \quad 0 \leq j \leq l$$

where $\iota_0 : BI_\eta \rightarrow \mathcal{Y}$ be the residual gerbe of a generic $y \in Y$. Since I_η acts trivially on Y , ι_0^* does not depend on the choice of general point.

Definition 5.5. For $\tau = (\tau_j) \in \prod_{1 \leq j \leq l} M_B(BG_j, r)$, we define the *fixed-inertia locus* to be

$$M_B(\mathcal{Y}, r, \tau) = \bigcap_{0 \leq j \leq l} \iota_j^*(\tau_j)$$

Since $M_B(BG_j, r)$ is finite and reduced for G_j finite, we get that every irreducible subvariety of $M_B(\mathcal{Y}, r)$ lies in one such fixed-inertia locus.

Proposition 5.6. For $\pi : Y \rightarrow \mathcal{Y} = [Y/G]$ we have the following

- (1) If $j : \mathcal{U} \rightarrow \mathcal{Y}$ is a nonempty open substack, then $j_* : \pi_1(\mathcal{U}, \bar{u}) \rightarrow \pi_1(\mathcal{Y}, \bar{u})$ is surjective. Consequently restriction of local systems from $\mathcal{Y}$ to $\mathcal{U}$ is fully faithful and preserves irreducibility.
- (2) If $f : \mathcal{X} \rightarrow \mathcal{Y}$ induces surjection on fundamental groups, then

$$f^* : M_B(\mathcal{Y}, r) \rightarrow M_B(\mathcal{X}, r)$$

is a closed immersion.

Proof. Let $U = \mathcal{U} \times_{\mathcal{Y}} Y$ then U is a G -invariant subset of Y , and we have a commutative diagram

$$\begin{array}{ccccccc} 1 & \longrightarrow & \pi_1(U, u) & \longrightarrow & \pi_1(\mathcal{U}, \bar{u}) & \longrightarrow & G \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \parallel \\ 1 & \longrightarrow & \pi_1(Y, u) & \longrightarrow & \pi_1(\mathcal{Y}, \bar{u}) & \longrightarrow & G \longrightarrow 1 \end{array}$$

where the left vertical surjection comes from $Y \setminus U$ being finitely many points. Thus we get a surjection $\pi_1(\mathcal{U}, \bar{u}) \twoheadrightarrow \pi_1(\mathcal{Y}, \bar{u})$. The second statement (2) is exactly the same as for character varieties of varieties. $\square$

5.2. A factorization theorem following Corlette-Simpson. With the notations in (5.1), let

$$D = \overline{\text{pr}_1(C)}_{\text{red}} \subset M_B(X, 2, \det = \epsilon), \quad T = \overline{\text{pr}_2(C)}_{\text{red}} \subset A$$

with T a torsion-translated subtorus, and $\epsilon \in A$ torsion. Notice that since a general point of Z has dense PGL_2 -image, the same is true for D .

Remark 5.7. We can look at the image of D under $\mathbb{P} : M_B(X, 2) \rightarrow M_B(X, \text{PGL}_2)$. If the image is zero-dimensional, then D itself must be finite since two fixed-determinant lifts of a PGL_2 -representation differs by an element of $A[2]$ which is finite. Since C is irreducible, we must have $D = \{\mathbb{V}\}$ and $C = \{\mathbb{V}\} \times T$. From now on, we assume that $\mathbb{P}(D)$ is positive-dimensional in $M_B(X, \text{PGL}_2)$.

The goal of this section is the following proposition:

Proposition 5.8. *There is a proper hyperbolic DM curve $\mathcal{Y} = [Y/G]$ and a morphism $s : X \rightarrow \mathcal{Y}$ and an irreducible component Q of a fixed-determinant, fixed-inertia locus $M_B(\mathcal{Y}, 2, \det = \delta, \tau)$ such that*

- (1) $s_* : \pi_1(X, x) \rightarrow \pi_1(\mathcal{Y}, \bar{y})$ is a surjection.
- (2) with the closed embedding $s^* : M_B(\mathcal{Y}, 2) \hookrightarrow M_B(X, 2)$, $D \subseteq s^*(Q)$.

Let $p : X^\epsilon \rightarrow X$ be the connected finite étale cover corresponding to $\ker \epsilon \triangleleft \pi_1(X, x)$. Then for every $\rho \in D \subset M_B(X, 2, \det = \epsilon)$, the restriction $\rho|_{X^\epsilon} \in M_B(X^\epsilon, \text{SL}_2)$. Call this map $\text{res}_\epsilon : M_B(X, 2, \det = \epsilon) \rightarrow M_B(X^\epsilon, \text{SL}_2)$.

Lemma 5.9. *The general point of $\text{res}_\epsilon(D)$ has Zariski-dense image in SL_2 , and is non-rigid.*

Proof. We start with some claims.

Claim. *If $\bar{\rho} : \pi_1(X, x) \rightarrow \text{PGL}_2(\mathbb{C})$ has Zariski-dense image in PGL_2 and $\Gamma \triangleleft \pi_1(X, x)$ a normal finite-index subgroup, then $\bar{\rho}(\Gamma)$ is Zariski-dense in PGL_2 .*

Proof of Claim. Let $H = \overline{\bar{\rho}(\Gamma)}$ be the Zariski closure. Then H is a normal finite-index subgroup of PGL_2 . However, PGL_2 is connected, hence $H = \text{PGL}_2$. $\blacksquare$

Claim. *On the locus of Zariski-dense representations, the restriction*

$$\text{res} : M_B(X, \text{PGL}_2) \rightarrow M_B(X^\epsilon, \text{PGL}_2)$$

is injective on closed points. Consequently, $\text{res}_\epsilon(D)$ has positive-dimensional image in $M_B(X^\epsilon, \text{SL}_2)$.

Proof of Claim. Suppose otherwise, and $\bar{\rho}_1, \bar{\rho}_2$ are Zariski-dense projective representations such that their restrictions to X^ϵ are conjugate. Since $\bar{\rho}_2 \in M_B(X, \text{PGL}_2)$ only depends on conjugacy class, we might assume that $\bar{\rho}_1|_{X^\epsilon} = \bar{\rho}_2|_{X^\epsilon} = \bar{\rho}_\epsilon$. For $\gamma \in \pi_1(X, x)$ and $h \in \ker \epsilon$, denote

$$c_\gamma = \bar{\rho}_2(\gamma)^{-1} \circ \bar{\rho}_1(\gamma) \in \text{PGL}_2$$

and we have, since $\gamma h \gamma^{-1} \in \ker(\epsilon)$,

$$c_\gamma \circ \bar{\rho}_\epsilon(h) \circ c_\gamma^{-1} = \bar{\rho}_\epsilon(h)$$

Now, by the previous claim, $\bar{\rho}_\epsilon$ has Zariski-dense image. Hence the centralizer of $\bar{\rho}_\epsilon(\ker \epsilon)$ agrees with the center $Z(\text{PGL}_2)$ which is trivial. Thus $c_\gamma = \text{Id}$ for all $\gamma \in \pi_1(X, x)$. It follows that

$$\text{res} : M_B(X, \text{PGL}_2) \rightarrow M_B(X^\epsilon, \text{PGL}_2)$$

is injective on closed points. We have a diagram

$$\begin{array}{ccc} M_B(X, 2, \det = \epsilon) & \xrightarrow{\text{res}_\epsilon} & M_B(X^\epsilon, \text{SL}_2) \\ \mathbb{P} \downarrow & & \downarrow \mathbb{P} \\ M_B(X, \text{PGL}_2) & \xrightarrow{\text{res}} & M_B(X^\epsilon, \text{PGL}_2) \end{array}$$

where both vertical maps are finite, $\mathbb{P}(D)$ is positive dimensional and its general point has Zariski-dense image, hence by the second claim $\text{res} \circ \mathbb{P}(D)$ has positive dimensional image. It follows that $\text{res}_\epsilon(D)$ has positive dimensional image. $\blacksquare$

Thus the general point $\rho_\epsilon \in \text{res}_\epsilon(D)$ is non-rigid. By the diagram, such a general point has $\mathbb{P}(\rho_\epsilon)$ with Zariski-dense image in PGL_2 . On the other hand, let $\overline{\rho_\epsilon(\ker \epsilon)} \subseteq \text{SL}_2$ be the Zariski closure of the image of ρ_ϵ , then $\overline{\rho_\epsilon(\ker \epsilon)} \twoheadrightarrow \text{PGL}_2$. As a result, $\dim \overline{\rho_\epsilon(\ker \epsilon)} \geq 3$ hence $\overline{\rho_\epsilon(\ker \epsilon)} = \text{SL}_2$. $\square$

For a general $\rho \in D$, by [Lemma 5.9](#), the restriction $\rho|_{X^\epsilon}$ has Zariski-dense image and is non-rigid in $M_B(X^\epsilon, \text{SL}_2)$. Hence we have the following as immediate consequences of [\[CS08\]](#); see also [\[Sim91, Theorem 10\]](#).

Lemma 5.10 ([\[CS08, Corollary 3.3 and Theorem 6.8\]](#)). *For a general $\rho \in D$, the restricted projective representation $\bar{\rho}|_{X^\epsilon}$ factors through an algebraic map $X^\epsilon \rightarrow Y$ where Y is a proper hyperbolic orbicurve.*

To get a factorization for $\bar{\rho}$ instead of $\bar{\rho}|_{X^\epsilon}$ we have the following lemma.

Lemma 5.11 ([\[CS08, Lemma 3.6\]](#)+ ϵ). *Let $f : X' \rightarrow X$ be a finite étale map of smooth projective varieties, and $\bar{\rho} : \pi_1(X, x) \rightarrow \text{PGL}_2(\mathbb{C})$ is a Zariski-dense representation such that $\bar{\rho}|_{X'}$ factors through a proper hyperbolic orbicurve. Then $\bar{\rho}$ itself factors through a proper hyperbolic orbicurve.*

Proof. By taking the Galois closure, we might assume $f : X' \rightarrow X$ is a Galois cover and $X = [X'/G]$. Let $s : X' \rightarrow Y$ be the factorization map for $\bar{\rho}|_{X'}$ with Y proper hyperbolic orbicurve. Let $Y' \rightarrow Y$ be the finite cover by a smooth curve. Replacing X' with $X' \times_Y Y'$ we can assume that the factorization map $s' : X' \rightarrow Y$ has smooth curve target. Take the Stein factorization, and remove from X the closure of the image of all fibers of s' which are reducible or not smooth (passing to an open subset of X is fine by [CS08, Lemma 3.5]). Thus we may assume that all fibers of s' are irreducible, and smooth.

The fibers of s' are maximal subvarieties W of X' such that $\bar{\rho}|_W$ is trivial. For $g \in G = \text{Gal}(X'/X)$, $f \circ g = f$ hence g permutes the fibers of $s' : X' \rightarrow Y$. Hence the G -action descends to Y making s' a G -equivariant map. Thus it descends to $s : X \rightarrow [Y/G]$.

$[Y/G]$ is an orbicurve if and only if the G -action is faithful. Let $I_\eta = \ker(G \rightarrow \text{Aut}(Y))$, then I_η is finite, and a normal subgroup of $\pi_1([Y/G], \bar{y})$. By the previous paragraphs, $\bar{\rho}$ descends to some $\bar{\theta} : \pi_1([Y/G], \bar{y}) \rightarrow \text{PGL}_2(\mathbb{C})$ which must also have Zariski-dense image. Then the normalizer $N_{\text{PGL}_2}(\bar{\theta}(I_\eta))$ contains $\bar{\theta}(\pi_1([Y/G], \bar{y}))$, thus must be PGL_2 . Hence $\bar{\theta}(I_\eta)$ is finite and normal in PGL_2 , which gives that $\bar{\theta}(I_\eta)$ is trivial. We can then replace $[Y/G]$ by $[Y/(G/K)]$ which has trivial generic inertia. $\square$

Corollary 5.12. *For a general $\rho \in D$, the projective representation $\bar{\rho}$ factors through a nonconstant map $s_{\bar{\rho}} : X \rightarrow Y_{\bar{\rho}}$ where $Y_{\bar{\rho}}$ is a proper hyperbolic orbicurve.*

Proof. Apply Lemma 5.11 to the cover $X^\epsilon \rightarrow X$, and use Lemma 5.10 to get factorization on X^ϵ . $\square$

Notice that if we have a finite cover $\pi : Y \rightarrow Y_{\bar{\rho}}$, and a lift $s : X \rightarrow Y$ of $s_{\bar{\rho}}$, then $s_{\bar{\rho}}^* = s^* \circ \pi^*$. Hence $s : X \rightarrow Y$ also satisfies the factorization property. By the following lemma, we can assume that $(s_{\bar{\rho}})_* : \pi_1(X, x) \rightarrow \pi_1(Y_{\bar{\rho}}, y)$ is a surjection.

Lemma 5.13. *Let $s : X \rightarrow Y$ be a nonconstant algebraic morphism from projective connected variety X to a proper hyperbolic orbicurve Y . Then there is a finite étale cover $\pi : Y' \rightarrow Y$ and an algebraic lift $s' : X \rightarrow Y'$ such that s' is surjective and $s'_* : \pi_1(X, x) \rightarrow \pi_1(Y', y')$ is surjective.*

Proof. By [Noo05, Proposition 18.18 and Corollary 18.20] there is a cover $\pi : Y' \rightarrow Y$ corresponding to $\text{im}(s_*) \subseteq \pi_1(Y, y)$, and an analytic lift $s' : X \rightarrow Y'$ with s'_* surjective. Since X is a projective variety and s' is nonconstant, we get that $\text{im}(s') = Y'$, and Y' is also proper. Thus $Y' \rightarrow Y$ is a finite analytic covering map. By [Noo05, Theorem 20.4] this is algebraizable. $\square$

Corollary 5.14. *For a general $\rho \in D$, the projective representation $\bar{\rho}$ factors through a map $s_{\bar{\rho}} : X \rightarrow Y_{\bar{\rho}}$ where $Y_{\bar{\rho}}$ is a proper hyperbolic orbicurve, $s_{\bar{\rho}}$ is surjective, and the induced map $(s_{\bar{\rho}})_* : \pi_1(X, x) \rightarrow \pi_1(Y_{\bar{\rho}}, y)$ is surjective.*

Proposition 5.15. *There is a proper hyperbolic orbicurve Y and an algebraic $s : X \rightarrow Y$ such that $s_* : \pi_1(X, x) \rightarrow \pi_1(Y, y)$ is surjective, and for every $\rho \in D$, the projective representation $\bar{\rho}$ factors through s .*

Proof. By [CS08, Proposition 2.8], there are only finitely many maps $s_j : X \rightarrow Y_j$ with Y_j a hyperbolic orbicurve, and s_j is surjective. By Lemma 5.13, we can assume all $(s_j)_*$ are surjective, thus by Proposition 5.6, we have closed embeddings

$$s_j^* : M_B(Y_j, \mathrm{PGL}_2) \hookrightarrow M_B(X, \mathrm{PGL}_2)$$

Let $\overline{\mathbb{P}(D)}$ be the closure of the image of D under

$$\mathbb{P} : M_B(X, 2) \rightarrow M_B(X, \mathrm{PGL}_2)$$

then by Corollary 5.14, a dense open subset of $\overline{\mathbb{P}(D)}$ must be contained in the finite union $\bigcup_j \mathrm{im}(s_j^*)$ which is Zariski-closed. Hence $\overline{\mathbb{P}(D)} \subseteq \bigcup_j \mathrm{im}(s_j^*)$. Since D is irreducible, so is $\overline{\mathbb{P}(D)}$ and thus must lie entirely in some $\mathrm{im}(s_j^*)$, and we are done. $\square$

Next we recover a factorization for every $\rho \in D$ through the projective factorization. See also [CS08, Lemma 3.1] for the strategy in the case ϵ is trivial, which we will adapt.

Lemma 5.16. *Let Y and s be as in Proposition 5.15. There is a finite gerbe $\mathcal{Y} \rightarrow Y$ and a lift $s_D : X \rightarrow \mathcal{Y}$ such that $s_{D,*} : \pi_1(X, x) \rightarrow \pi_1(\mathcal{Y}, \bar{y})$ is surjective, and every $\rho \in D$ factors through $s_D : X \rightarrow \mathcal{Y}$.*

Proof. For $\rho \in D$, notice that $\bar{\rho}$ factors through $s : X \rightarrow Y$ hence $\ker(s_*) \subseteq \ker(\bar{\rho})$. It follows that for every $k \in \ker(s_*)$, $\rho(k)$ is scalar, and we can write

$$\rho(k) = \lambda_\rho(k) \cdot \mathrm{Id}, \quad \lambda_\rho(k)^2 = \det \rho(k) = \epsilon(k)$$

This λ_ρ is independent of $\rho \in D$.

Claim. *There is a $\pi_1(X, x)$ -invariant character $\lambda : \ker(s_*) \rightarrow \mu_{2m}$ such that $\rho(k) = \lambda(k) \cdot \mathrm{Id}$ for every $\rho \in D$ and $k \in \ker(s_*)$.*

Proof of Claim. Indeed, for each $k \in \ker(s_*)$ consider the regular function $f_k \in \mathcal{O}_D(D)$ by

$$f_k(\rho) = \frac{1}{2} \mathrm{tr}(\rho(k))$$

then $f_k(\rho) = \lambda_\rho(k)$, $f_k^2(\rho) = \epsilon(k)$ for all $\rho \in D$. Since $\mathcal{O}_D(D)$ is integral, $f_k(\rho)$ must be constant and equal to one of the two square roots of $\epsilon(k)$; denote this square root $\lambda(k) = f_k$. For $k_1, k_2 \in \ker(s_*)$ and $\rho \in D$,

$$\lambda(k_1 k_2) \cdot \mathrm{Id} = \lambda_\rho(k) \cdot \mathrm{Id} = \rho(k_1 k_2) = \rho(k_1) \rho(k_2) = \lambda(k_1) \lambda(k_2) \cdot \mathrm{Id}$$

hence λ is multiplicative. Similarly, one can check that λ is invariant under conjugation by $\pi_1(X, x)$. Finally, since ϵ is torsion, there is $m \in \mathbb{Z}_+$ such that $\epsilon^m = 1$ hence $\lambda^{2m} = 1$. $\blacksquare$

We thus have a central (since $[\pi_1(X, x), \ker(s_*)] \subseteq \ker(\lambda)$) extension

$$1 \rightarrow \ker(s_*) / \ker(\lambda) \rightarrow \pi_1(X, x) / \ker(\lambda) \xrightarrow{s_*} \pi_1(Y, y) \rightarrow 1$$

with $\ker(s_*) / \ker(\lambda) \simeq \mathrm{im}(\lambda)$ finite. By [CS08, Lemma 2.5], this determines a finite $\ker(s_*) / \ker(\lambda)$ -gerbe $\mathcal{Y} \rightarrow Y$ with $\pi_1(\mathcal{Y}, \bar{y}) = \pi_1(X, x) / \ker(\lambda)$. Furthermore, the quotient $\pi_1(X, x) \rightarrow \pi_1(\mathcal{Y}, \bar{y})$ is a lift of $s_* : \pi_1(X, x) \rightarrow \pi_1(Y, y)$ which gives a lift

$s_D : X \rightarrow \mathcal{Y}$ of $s : X \rightarrow Y$. By construction, $s_{D,*}$ is surjective. Notice that every $\rho \in D$ has $\rho(\ker(\lambda)) = 1$ hence ρ factors through $\pi_1(\mathcal{Y}, \bar{y})$ as desired. $\square$

Proof of Proposition 5.8. By Lemma 5.16 we get a proper hyperbolic DM curve $\mathcal{Y}$ and $s_D : X \rightarrow \mathcal{Y}$ such that every $\rho \in D$ factors through s_D , i.e. D lies in the image of the closed embedding

$$s_D^* : M_B(\mathcal{Y}, 2) \rightarrow M_B(X, 2)$$

The preimage $(s_D^*)^{-1}(D)$ is irreducible. Furthermore, for every $\psi \in (s_D^*)^{-1}(D)$,

$$s_D^*(\det \psi) = \det(s_D^* \psi) = \epsilon$$

and $s_D^* : M_B(\mathcal{Y}, 1) \hookrightarrow M_B(X, 1)$ is also injective hence $\det(\psi)$ is constant for all $\psi \in (s_D^*)^{-1}(D)$. It follows that $\psi \in (s_D^*)^{-1}(D)$ is irreducible and has fixed-determinant, and thus must lie in a fixed-determinant, fixed-inertia locus by Definition 5.5. Finally, the construction of Lemma 5.11 gives an proper hyperbolic orbicurve of the form $[Y/G]$. The finite gerbe $\mathcal{Y} \rightarrow [Y/G]$ from Lemma 5.16 remains a proper hyperbolic finite quotient curve. $\square$

5.3. $C \rightarrow T$ is either a product or generically finite. Recall (5.1) that we have C an irreducible component of $\Phi_\epsilon^{-1}(Z) \subseteq M_B(X, 2, \det = \epsilon) \times A$ with Z an irreducible component of the reduced jump loci $\bar{\Sigma}_k^i(X, 2) \subseteq M_B(X, 2)$. And the two projections give

$$D = \overline{\text{pr}_1(C)}_{\text{red}} \subseteq M_B(X, 2), \quad T = \overline{\text{pr}_2(C)}_{\text{red}} \subseteq A$$

and by Proposition 5.8, there is a proper hyperbolic DM curve $\mathcal{Y} = [Y/G]$ (with Y smooth and G finite), an irreducible component Q of a fixed-determinant, fixed-inertia locus of $M_B(\mathcal{Y}, 2)$ such that, and a map $s : X \rightarrow \mathcal{Y}$ such that $D \subseteq s^*(Q)$. Since $C \subseteq D \times T$, and

$$(s^*, \text{id}) : M_B(\mathcal{Y}, 2) \times A \rightarrow M_B(X, 2) \times A$$

is a closed embedding, there is some $\tilde{C} \subseteq Q \times T$ isomorphic to C . We might as well work with $C \subseteq Q \times T$. Denote

$$\tilde{\Sigma}_k^i = \{(\mathbb{W}, \mathbb{L}) \in Q \times T \mid \dim H^i(s^* \mathbb{W} \otimes \mathbb{L}) \geq k\}$$

then $C \subseteq \tilde{\Sigma}_k^i$. We start with a decomposition lemma for the proper map $s : X \rightarrow \mathcal{Y}$. We have the following diagram

$$\begin{array}{ccc} X' & \xrightarrow{\pi'} & X \\ s' \downarrow & & \downarrow s \\ Y & \xrightarrow{\pi} & \mathcal{Y} \end{array} \tag{5.2}$$

where π and π' are both finite étale G -torsor. $X = [X'/G]$, and $s' : X' \rightarrow Y$ is G -equivariant. We have base change for $K \in D_c^b(X, \mathbb{C})$,

$$\pi^* \text{R} s_* K \simeq \text{R} s'_*(\pi')^* K$$

Lemma 5.17. *Let $\mathbb{L}$ be a rank one local system on X , then there is G -invariant dense open $j : U \hookrightarrow Y$ and a G -equivariant decomposition*

$$\mathrm{Rs}'_*(\pi')^*\mathbb{L} \simeq \bigoplus_{\alpha} j_* \mathbb{U}_{\alpha}[-a_{\alpha}] \oplus \bigoplus_{\beta} \iota_{\beta,*} \mathcal{V}_{\beta}[-b_{\beta}]$$

where $a_{\alpha}, b_{\beta} \in \mathbb{Z}$, $\mathbb{U}_{\alpha}$ is an irreducible G -equivariant local system on U , and $\mathcal{V}_{\beta}$ is an irreducible local system on the orbit $G \cdot y_{\beta}$ of a point $y_{\beta} \in Y \setminus U$ with G -equivariant inclusion $\iota_{\beta} : G \cdot y_{\beta} \hookrightarrow Y$.

For $\mathbb{W}$ a local system on $\mathcal{Y}$, let $V_{\beta} = (\mathcal{V}_{\beta})_{y_{\beta}}$ and G_{β} the stabilizer of y_{β} . There is a decomposition of cohomology

$$\begin{aligned} H^i(X, s^*\mathbb{W} \otimes \mathbb{L}) &\simeq \bigoplus_{\alpha} H^{i-a_{\alpha}}(Y, (\pi^*\mathbb{W}) \otimes j_* \mathbb{U}_{\alpha})^G \\ &\quad \oplus \bigoplus_{\beta, b_{\beta}=i} ((\pi^*\mathbb{W})_{y_{\beta}} \otimes V_{\beta})^{G_{\beta}} \end{aligned}$$

Proof. We give a sketch here. $\mathbb{L}' = (\pi')^*\mathbb{L}$ has rank one, and thus is irreducible. By [Sab05, Main Theorem 1], we have a decomposition

$$\mathrm{Rs}'_*(\mathbb{L}'[-\dim X]) \simeq \bigoplus_i {}^p\mathcal{H}^i(\mathrm{Rs}'_*(\mathbb{L}'[-\dim X]))[-i] \quad (5.3)$$

where each ${}^p\mathcal{H}^i(\mathrm{Rs}'_*(\mathbb{L}'[-\dim X]))$ further decomposes into intersection complexes supported on closed irreducible analytic subset of Y . Since Y is a smooth projective curve here, these are either of the form $j_*\mathbb{U}$ where $\mathbb{U}$ is a local system on a Zariski open subset of Y , or skyscraper sheaves.

Notice that $\mathbb{L}' = (\pi')^*\mathbb{L}$ is G -equivariant, and s' is G -equivariant hence the same is true for

$$K := \mathrm{Rs}'_*(\mathbb{L}'[-\dim X])$$

For $g \in G$, $g : Y \rightarrow Y$ is an isomorphism hence g^* preserves perverse t-structure. Thus the truncation triangle

$${}^p\tau_{\leq i-1}K \rightarrow {}^p\tau_{\leq i}K \rightarrow {}^p\mathcal{H}^i(K) \rightarrow {}^p\tau_{\leq i-1}K[1] \quad (5.4)$$

is also G -equivariant. Sabbah's decomposition (5.3) gives a splitting of this triangle. Since G is finite, by averaging we get a G -equivariant splitting. Thus we might assume that (5.3) is G -equivariant. Each ${}^p\mathcal{H}^i(K)$ is G -equivariant, hence the G -action will permute the skyscraper sheaves. Thus the skyscraper sheaves come in G -orbits, as stated in the lemma.

Next, the full-support part of ${}^p\mathcal{H}^i(K)$ has to look like $j_{0,*}(\mathbb{U}_0[1])$ for some semisimple local system $\mathbb{U}_0$ on Zariski open $j_0 : U_0 \hookrightarrow Y$. Let $U \subseteq U_0$ be a G -invariant Zariski open subset of Y with G -equivariant inclusion $j : U \hookrightarrow Y$, and put

$$\mathbb{U} = \mathcal{H}^{-1}(j^{-1}j_{0,*}(\mathbb{U}_0[1])) \simeq \mathbb{U}_0|_U$$

then $\mathbb{U}$ is a G -equivariant local system on U . It is semisimple since $\pi_1(U_0) \twoheadrightarrow \pi_1(U)$. By uniqueness of intermediate extension, we get a G -equivariant isomorphism

$$j_*(\mathbb{U}[1]) \simeq j_{0,*}(\mathbb{U}_0[1])$$

For the cohomology statement, by projection formula we have

$$\begin{aligned} H^i(X, s^*\mathbb{W} \otimes \mathbb{L}) &\simeq H^i(X', (\pi')^* s^*\mathbb{W} \otimes (\pi')^*\mathbb{L})^G \\ &\simeq H^i(X', (s')^* \pi^*\mathbb{W} \otimes (\pi')^*\mathbb{L})^G \\ &\simeq \mathbb{H}^i(Y, \pi^*\mathbb{W} \otimes R(s')_*(\pi')^*\mathbb{L})^G \end{aligned}$$

and we can apply the decomposition of $R(s')_*(\pi')^*\mathbb{L}$ to conclude. $\square$

Proposition 5.18. *C is an irreducible component of $\tilde{\Sigma}_k^i$. For every $\mathbb{L} \in T$, the fiber $\tilde{\Sigma}_k^i|_{\mathbb{L}}$ is either Q , a finite set, or empty. Consequently, $C = Q \times T$, or $C \rightarrow T$ is generically finite.*

Proof. If an irreducible component C' of $\tilde{\Sigma}_k^i$ contains C , then

$$Z = \Phi_\epsilon(C) \subseteq \Phi_\epsilon(C') \subseteq \bar{\Sigma}_k^i(X, 2)$$

and Φ_ϵ is finite hence $\Phi_\epsilon(C')$ is irreducible. But Z is an irreducible component, hence $Z = \Phi_\epsilon(C')$. Finiteness then gives $\dim C = \dim C'$ thus $C = C'$.

Next, recall that the fiber $\tilde{\Sigma}_k^i|_{\mathbb{L}}$ is the locus

$$\{\mathbb{W} \in Q \mid \dim H^i(s^*\mathbb{W} \otimes \mathbb{L}) \geq k\}$$

and by Lemma 5.17, it suffices to look at the jumping behavior of each

$$H^{i-a}(Y, (\pi^*\mathbb{W}) \otimes j_*\mathbb{U})^G, \quad ((\pi^*\mathbb{W})_y \otimes V)^{G_y}$$

with $j : U \hookrightarrow Y$ the inclusion of a G -invariant open, $\mathbb{U}$ an irreducible G -equivariant local system on U , $y \in Y \setminus U$ with G_y the stabilizer, and V an irreducible G_y -representation. Notice that $j : U \rightarrow Y$, $\mathbb{U}$, G_y and V do not depend on $\mathbb{W}$. Also, since Y is a smooth curve, $H^{i-a} = 0$ unless $0 \leq i - a \leq 2$.

Claim. *The G_y -representation $(\pi^*\mathbb{W})_y$ does not depend on $\mathbb{W}$. Consequently,*

$$\dim((\pi^*\mathbb{W})_y \otimes V)^{G_y}$$

is constant as $\mathbb{W}$ varies in Q .

Proof of Claim. We have injective $G_y \hookrightarrow \pi_1(\mathcal{Y}, \bar{y})$ hence each $g \in G_y$ determines an element of $\gamma_g \in \pi_1(\mathcal{Y}, \bar{y})$. Thus it induces a regular function on Q

$$\mathbb{W} \mapsto \text{tr}(g|_{(\pi^*\mathbb{W})_y}) = \text{tr}(\rho_{\mathbb{W}}(\gamma_g))$$

where $\rho_{\mathbb{W}}$ is the $\pi_1(\mathcal{Y}, \bar{y})$ -representation corresponding to $\mathbb{W}$. Since G_y is finite, the function defined above has discrete image. Since Q is irreducible, it must be constant. It follows that all G_y -representations $(\pi^*\mathbb{W})_y$, as $\mathbb{W}$ varies in Q , have the same characters hence must be isomorphic. $\blacksquare$

Moving on, notice that $(\pi^*\mathbb{W}) \otimes j_*\mathbb{U} = j_*((\pi^*\mathbb{W})|_U \otimes \mathbb{U})$, and denote

$$h_G^l(\mathbb{W}, \mathbb{U}) = \dim H^l(Y, j_*((\pi^*\mathbb{W})|_U \otimes \mathbb{U}))^G$$

and $\chi_G(\mathbb{W}, \mathbb{U}) = h_G^0(\mathbb{W}, \mathbb{U}) - h_G^1(\mathbb{W}, \mathbb{U}) + h_G^2(\mathbb{W}, \mathbb{U})$.

Claim. *$\chi_G(\mathbb{W}, \mathbb{U})$ is constant as $\mathbb{W}$ varies in Q .*

Proof of Claim. Let $\bar{Y} = Y/G$ be the coarse space for $[Y/G]$, with the quotient $q : Y \rightarrow \bar{Y}$ which is a finite map. Denote the constructible sheaf

$$\mathcal{L}(\mathbb{W}, \mathbb{U}) := j_*((\pi^*\mathbb{W})|_U \otimes \mathbb{U}), \quad \bar{\mathcal{L}}(\mathbb{W}, \mathbb{U}) := (q_*\mathcal{L}(\mathbb{W}, \mathbb{U}))^G$$

then $H^l(Y, \mathcal{L}(\mathbb{W}, \mathbb{U}))^G \simeq H^l(\bar{Y}, \bar{\mathcal{L}}(\mathbb{W}, \mathbb{U}))$. Denote $I_\eta = \ker(G \rightarrow \text{Aut}(Y))$ the generic inertia. Denote $E \subset \bar{Y}$ a finite set of points, consisting of $q(Y \setminus U)$ and images of points on Y with larger than generic stabilizer. Then $\bar{\mathcal{L}}(\mathbb{W}, \mathbb{U})|_{\bar{Y} \setminus E}$ is a local system of dimension

$$r(\mathbb{W}, \mathbb{U}) = \dim((\pi^*\mathbb{W})_y \otimes \mathbb{U}_y)^{G_y} = \dim((\pi^*\mathbb{W})_y \otimes \mathbb{U}_y)^{I_\eta}, \quad y \in q^{-1}(\bar{Y} \setminus E)$$

and for $z \in E$, $\bar{\mathcal{L}}(\mathbb{W}, \mathbb{U})|_z$ is a vector space of dimension

$$\dim((\pi^*\mathbb{W})_y \otimes \mathbb{U}_y)^{G_y}, \quad y \in q^{-1}(z)$$

By the same argument as in previous claim, both of these values are constant as $\mathbb{W}$ varies in Q . On the other hand, since compactly supported Euler characteristic is additive for the stratification $\bar{Y} = E \sqcup (\bar{Y} \setminus E)$ we get that

$$\begin{aligned} \chi_G(\mathbb{W}, \mathbb{U}) &= \chi(\bar{Y}, \bar{\mathcal{L}}(\mathbb{W}, \mathbb{U})) \\ &= \chi_c(\bar{Y} \setminus E, \bar{\mathcal{L}}(\mathbb{W}, \mathbb{U})|_{\bar{Y} \setminus E}) + \sum_{z \in E} \dim \bar{\mathcal{L}}(\mathbb{W}, \mathbb{U})|_z \\ &= \chi_c(\bar{Y} \setminus E) \cdot r(\mathbb{W}, \mathbb{U}) + \sum_{z \in E} \dim \bar{\mathcal{L}}(\mathbb{W}, \mathbb{U})|_z \\ &= (2 - 2g(\bar{Y}) - |E|) \cdot r(\mathbb{W}, \mathbb{U}) + \sum_{z \in E} \dim \bar{\mathcal{L}}(\mathbb{W}, \mathbb{U})|_z \end{aligned}$$

which is constant as $\mathbb{W}$ varies in Q , as desired. ■

Next, we examine these cohomology groups more explicitly:

$$\begin{aligned} H^0(Y, j_*((\pi^*\mathbb{W})|_U \otimes \mathbb{U}))^G &\simeq H^0(U, (\pi^*\mathbb{W})|_U \otimes \mathbb{U})^G \\ &\simeq \text{Hom}_{\text{Loc}^G(U)}(\mathbb{U}^\vee, (\pi^*\mathbb{W})|_U) \end{aligned}$$

and by Poincaré duality [Dim04, Corollary 3.3.12]

$$H^2(Y, j_*((\pi^*\mathbb{W})|_U \otimes \mathbb{U}))^G \simeq \text{Hom}_{\text{Loc}^G(U)}((\pi^*\mathbb{W})|_U, \mathbb{U}^\vee)^\vee$$

Since $\mathbb{W}$ is semisimple on $\mathcal{Y}$, $\pi^*\mathbb{W}$ is semisimple in $\text{Loc}^G(Y)$. Thus $(\pi^*\mathbb{W})|_U$ is semisimple in $\text{Loc}^G(U)$. It follows that the two cohomology groups above both count the multiplicities of $\mathbb{U}^\vee$ in $(\pi^*\mathbb{W})|_U$ and $h_G^0(\mathbb{W}, \mathbb{U}) = h_G^2(\mathbb{W}, \mathbb{U})$.

Claim. $h_G^0(\mathbb{W}, \mathbb{U}) \neq 0$ for only finitely many $\mathbb{W} \in Q$.

Proof of Claim. The observation which will be used throughout is that the functor

$$\text{Loc}(\mathcal{Y}) \rightarrow \text{Loc}_G(U), \quad \mathbb{W} \mapsto (\pi^*\mathbb{W})|_U$$

is fully faithful. If $\mathbb{U}$ has rank ≥ 2 , then if $h_G^0(\mathbb{W}, \mathbb{U}) \neq 0$ we must have $(\pi^*\mathbb{W})|_U \simeq \mathbb{U}^\vee$. There is at most one such $\mathbb{W} \in Q$.

Now suppose $\mathbb{U}$ has rank 1. If $h_G^0(\mathbb{W}, \mathbb{U}) \neq 0$ then $\mathbb{U}^\vee$ must be a summand of $(\pi^*\mathbb{W})|_U$ and $(\pi^*\mathbb{W})|_U$ must be reducible in $\text{Loc}^G(U)$. It follows that $(\pi^*\mathbb{W})$

must be reducible in $\text{Loc}^G(Y)$. Since $\mathbb{W} \in Q$, $s^*\mathbb{W} \in D \subset M_B(X, 2, \det = \epsilon)$, with $s^* : M_B(\mathcal{Y}, 1) \hookrightarrow M_B(X, 1)$ a closed embedding, we get that $\pi^*\mathbb{W}$ has fixed determinant $s^*\epsilon$. It follows that

$$\pi^*\mathbb{W} \simeq \mathbb{M} \oplus (\mathbb{M}^\vee \otimes s^*\epsilon)$$

and $\mathbb{U}^\vee = \mathbb{M}$ or $\mathbb{M}^\vee \otimes s^*\epsilon$. Again, finitely many $\mathbb{W}$ giving this. $\blacksquare$

Each $\mathbb{U}$ in Lemma 5.17 gives a finite set of $\mathbb{W} \in Q$ where $h_G^0(\mathbb{W}, \mathbb{U}) \neq 0$. Take $F(\mathbb{L})$ to be the union as $\mathbb{U}$ runs over all the finitely many summands in the decomposition of $\text{Rs}'_*(\pi')^*\mathbb{L}$, then $F(\mathbb{L})$ is also finite. To summarize, for a fixed $\mathbb{L}$

$$\dim H^0(Y, (\pi^*\mathbb{W}) \otimes j_*\mathbb{U})^G = h_G^0(\mathbb{W}, \mathbb{U}) = 0 \quad \forall \mathbb{W} \in Q \setminus F(\mathbb{L})$$

$$\dim H^2(Y, (\pi^*\mathbb{W}) \otimes j_*\mathbb{U})^G = h_G^2(\mathbb{W}, \mathbb{U}) = h_G^0(\mathbb{W}, \mathbb{U}) \quad \forall \mathbb{W} \in Q \setminus F(\mathbb{L})$$

and thus for all $\mathbb{W} \in Q \setminus F(\mathbb{L})$,

$$\begin{aligned} \dim H^1(Y, (\pi^*\mathbb{W}) \otimes j_*\mathbb{U})^G &= -\chi_G(\mathbb{W}, \mathbb{U}) + 2h_G^0(\mathbb{W}, \mathbb{U}) \\ &= -\chi_G(\mathbb{W}, \mathbb{U}) \end{aligned}$$

which is constant for $\mathbb{W} \in Q$. It follows that there is a constant $k_{\mathbb{L}}$ such that

$$\dim H^i(X, s^*\mathbb{W} \otimes \mathbb{L}) = k_{\mathbb{L}} \quad \forall \mathbb{W} \in Q \setminus F(\mathbb{L}) \quad (5.5)$$

$$\dim H^i(X, s^*\mathbb{W} \otimes \mathbb{L}) \geq k_{\mathbb{L}} \quad \forall \mathbb{W} \in Q \quad (5.6)$$

and consequently we have proven:

Claim. *For every $\mathbb{L} \in T$, the fiber $\tilde{\Sigma}_k^i|_{\mathbb{L}}$ is either Q , finite, or empty.*

Back to $C \subseteq \tilde{\Sigma}_k^i \subseteq Q \times T$. The map $C \rightarrow T$ is dominant by definition. Here we can assume $\dim Q > 0$ by Remark 5.7. Suppose that the general fiber of $C \rightarrow T$ has positive dimension, then on a Zariski open subset $T^\circ \subseteq T$, since $\tilde{\Sigma}_k^i \supset C$, $\tilde{\Sigma}_k^i|_{\mathbb{L}}$ has fiber Q . Hence,

$$Q \times T^\circ \subset \tilde{\Sigma}_k^i \subset Q \times T$$

and since $\tilde{\Sigma}_k^i$ is closed, we get $\tilde{\Sigma}_k^i = Q \times T$. This is irreducible, so $C = Q \times T$. The other case is that $C \rightarrow T$ is generically finite. $\square$

5.4. Finiteness of $C \rightarrow T$. In this section, we will prove that if $C \xrightarrow{\text{pf}_2} T$ is generically finite, then it must in fact be finite. To differentiate, let the G -invariant dense open in Lemma 5.17 be denoted $j_{\mathbb{L}} : U_{\mathbb{L}} \hookrightarrow Y$ for the decomposition of $\text{Rs}'_*(\pi')^*\mathbb{L}$. Also denote $\mathbb{L}' = (\pi')^*\mathbb{L}$.

For $U \subseteq Y$ a G -equivariant Zariski open subset, we have a diagram by restricting (5.2)

$$\begin{array}{ccc} X'_U & \xrightarrow{\pi'_U} & X_U \\ s'_U \downarrow & & \downarrow s_U \\ U & \xrightarrow{\pi_U} & [U/G] \end{array}$$

and we can pick U so that the map $s'_U : X'_U \rightarrow U$ is smooth and proper. Then

$$\mathcal{H}_{\mathbb{L}}^m = R^m(s'_U)_*(\mathbb{L}'|_{X'_U}) \quad (5.7)$$

is a G -equivariant local system on U .

Lemma 5.19. *Let $\mathbb{L} \in T$, and suppose that the fiber $\tilde{\Sigma}_k^i|_{\mathbb{L}}$ is finite. Let $(\mathbb{W}, \mathbb{L}) \in C \subseteq \tilde{\Sigma}_k^i$ with $\mathbb{W}$ irreducible in $\text{Loc}(\mathcal{Y})$, then, for some m , there is a quotient*

$$\mathcal{H}_{\mathbb{L}}^m \rightarrow ((\pi^*\mathbb{W})|_U)^\vee$$

in $\text{Loc}^G(U)$. In fact, $(\pi^*\mathbb{W})|_U^\vee$ is a direct summand of $\mathcal{H}_{\mathbb{L}}^m$.

Proof. Look at the G -equivariant decomposition of Lemma 5.17,

$$R(s'_*)\mathbb{L}' \simeq \bigoplus_{\alpha} j_{\mathbb{L},*}\mathbb{U}_{\alpha}[-a_{\alpha}] \oplus \bigoplus_{\beta} \iota_{\beta,*}\mathcal{V}_{\beta}[-b_{\beta}] \quad (5.8)$$

with $\mathbb{U}_{\alpha}$ irreducible in $\text{Loc}^G(U_{\mathbb{L}})$. By the proof of Proposition 5.18 (see in particular the paragraphs around (5.5)), if $\tilde{\Sigma}_k^i|_{\mathbb{L}}$ is finite, then it consists of points $(\mathbb{W}, \mathbb{L})$ such that

$$\dim H^0(Y, (\pi^*\mathbb{W}) \otimes j_{\mathbb{L},*}\mathbb{U}_{\alpha})^G > 0$$

which implies that $\text{Hom}_{\text{Loc}^G(U_{\mathbb{L}})}(\mathbb{U}_{\alpha}^\vee, (\pi^*\mathbb{W})|_{U_{\mathbb{L}}}) > 0$. Since $\mathbb{W} \mapsto (\pi^*\mathbb{W})|_{U_{\mathbb{L}}}$ is a fully faithful functor which preserves irreducibility, we have that

$$\mathbb{U}_{\alpha} \simeq ((\pi^*\mathbb{W})|_{U_{\mathbb{L}}})^\vee \quad \text{in } \text{Loc}^G(U_{\mathbb{L}})$$

Next, by restricting the decomposition (5.8) to U we get that $(j_{\mathbb{L},*}\mathbb{U}_{\alpha})|_U$ is a direct summand of $\mathcal{H}_{\mathbb{L}}^{a_{\alpha}}$ in $\text{Loc}^G(U)$; in particular, $(j_{\mathbb{L},*}\mathbb{U}_{\alpha})|_U$ is a local system on U . The previous argument show that

$$(j_{\mathbb{L},*}\mathbb{U}_{\alpha})|_{U \cap U_{\mathbb{L}}} \simeq \mathbb{U}_{\alpha}|_{U \cap U_{\mathbb{L}}} \simeq ((\pi^*\mathbb{W})|_{U \cap U_{\mathbb{L}}})^\vee$$

and since restriction $\text{Loc}^G(U) \rightarrow \text{Loc}^G(U \cap U_{\mathbb{L}})$ is fully faithful, we get that

$$(j_{\mathbb{L},*}\mathbb{U}_{\alpha})|_U \simeq ((\pi^*\mathbb{W})|_U)^\vee \quad \text{in } \text{Loc}^G(U)$$

and thus $((\pi^*\mathbb{W})|_U)^\vee$ is a direct summand of $\mathcal{H}_{\mathbb{L}}^{a_{\alpha}}$. Since $\mathcal{H}_{\mathbb{L}}^{a_{\alpha}}$ is semisimple we get the quotient statement as well. $\square$

Let $\mathcal{O}(T) = \Gamma(T, \mathcal{O}_T)$. Pulling back the universal rank 1 local system along $T \subseteq M_B(X, 1)$ we get $\mathbb{L}_{\text{univ}}$ a local system of free $\mathcal{O}(T)$ -modules on X . For $\mathbb{L} \in T$,

$$\mathbb{L}_{\text{univ}} \otimes_{\mathcal{O}(T)} \kappa(\mathbb{L}) \simeq \mathbb{L}$$

where we are taking tensor products with, and over, constant local systems on X . Again, denote $\mathbb{L}'_{\text{univ}} = (\pi')^*\mathbb{L}_{\text{univ}}$, then $\mathbb{L}'_{\text{univ}}$ is a G -equivariant local system of $\mathcal{O}(T)$ -modules on X' (the G -action on T is trivial). We also have that

$$\mathcal{H}_{\text{univ}}^m = R^m(s'_U)_*(\mathbb{L}'_{\text{univ}}|_{X'_U})$$

is a G -equivariant local system of $\mathcal{O}(T)$ -modules on U , since s'_U is a smooth and proper G -equivariant map. Equivalently, fixing $u \in U$ with $\bar{u}$ the image in $[U/G]$, $(\mathcal{H}_{\text{univ}}^m)_u$ is a $\mathcal{O}(T)$ -module, and we have a $\mathcal{O}(T)$ -linear representation

$$\pi_1([U/G], \bar{u}) \rightarrow \text{GL}_{\mathcal{O}(T)}((\mathcal{H}_{\text{univ}}^m)_u)$$

On the other hand, we can also think of $(\mathcal{H}_{\text{univ}}^m)_u$ as a $\mathcal{O}_T$ -module on T . There is then a Zariski dense open $T^{\text{bc}} \subseteq T$ where cohomology and base change is available, i.e., for $\mathbb{L} \in T^{\text{bc}}$, there is an isomorphisms of $\pi_1([U/G], \bar{u})$ -representations

$$(\mathcal{H}_{\text{univ}}^m)_u \otimes \kappa(\mathbb{L}) \simeq (\mathcal{H}_{\mathbb{L}}^m)_u$$

with $\mathcal{H}_{\mathbb{L}}^m$ defined in (5.7).

Remark 5.20. By Lemma 5.19, a point $(\mathbb{W}, \mathbb{L})$ in a finite fiber of $C \rightarrow T$ with $\mathbb{L} \in T^{\text{bc}}$ and $\mathbb{W}$ irreducible correspond to a rank two quotient

$$(\mathcal{H}_{\text{univ}}^m)_u \otimes \kappa(\mathbb{L}) \rightarrow ((\pi^*\mathbb{W})|_U)_u^\vee$$

as $\pi_1([U/G], \bar{u})$ -representations.

Proposition 5.21 ([HS04, Corollary 2.4]). *Let T be an affine noetherian variety, $\mathcal{H}$ a coherent $\mathcal{O}_T$ -module, and Π a finitely generated group. Suppose that there is a $\mathcal{O}_T$ -linear action of Π on $\mathcal{H}$. The functor assigning to an affine T -scheme $\phi : S \rightarrow T$ the set*

$$\{\Pi - \text{invariant locally free rank 2 quotients of } \phi^*\mathcal{H}\}$$

is representable by a projective T -scheme.

Proof. Since we are on affine schemes, we will abuse notation between sheaves and modules. Denote the $\mathcal{O}(T)$ -linear action of Π on $\mathcal{H}$ by

$$\rho : \Pi \rightarrow \text{GL}_{\mathcal{O}(T)}(\mathcal{H})$$

then our functor assigns to a commutative $\mathcal{O}(T)$ -algebra R the set of $\rho(\Pi)$ -invariant locally free R -module quotients of $\mathcal{H} \otimes_{\mathcal{O}(T)} R$. To apply the cited result, let $k = \mathcal{O}(T)$, A be the singleton set, $T = \mathcal{H}$, and $F = \rho(\Pi)$. $\square$

Apply the previous proposition to $\Pi = \pi_1([U/G], \bar{u})$ and $\mathcal{H} = (\mathcal{H}_{\text{univ}}^m)_u$. Let the resulting projective T -scheme be denoted $\mathcal{G}_m$. Notice that a closed point of $\mathcal{G}_m$, over $\mathbb{L} \in T^{\text{bc}}$, is a $\pi_1([U/G], \bar{u})$ -invariant quotient of $(\mathcal{H}_{\text{univ}}^m)_u \otimes \kappa(\mathbb{L}) \simeq (\mathcal{H}_{\mathbb{L}}^m)_u$. If $\mathbb{L} \notin T^{\text{bc}}$ then the quotient is still a rank 2 $\pi_1([U/G], \bar{u})$ -representation. We have a natural map over T

$$\Phi_m : \mathcal{G}_m \rightarrow M_B([U/G], 2) \times T, \quad [(\mathcal{H}_{\text{univ}}^m)_u \otimes \kappa(\mathbb{L}) \rightarrow \mathbb{V}] \mapsto ([\mathbb{V}^\vee], \mathbb{L})$$

which has image finite over T since $M_B([U/G], 2)$ is affine.

Proposition 5.22. *There is a closed subscheme $\mathcal{F}_Q \subset Q \times T$, finite over T , with the following property: if $\mathbb{W} \in Q$ and $\mathbb{L} \in T^{\text{bc}}$, and $((\pi^*\mathbb{W})|_U)^\vee$ is a rank two G -equivariant quotient of $\mathcal{H}_{\mathbb{L}}^m$ for some m , then $(\mathbb{W}, \mathbb{L}) \in \mathcal{F}_Q$.*

Proof. Denote the disjoint union

$$\mathcal{G} = \bigsqcup_{m=0}^{2 \dim X - 1} \mathcal{G}_m$$

and let $\mathcal{F}$ be the image of $\Phi : \mathcal{G} \rightarrow M_B([U/G], 2) \times T$. Then $\mathcal{F}$ is finite over T . We have a closed embedding

$$M_B(\mathcal{Y}, 2) \hookrightarrow M_B([U/G], 2), \quad \mathbb{W} \mapsto (\pi^*\mathbb{W})|_U$$

hence we can view $Q \times T$ as a closed subscheme of $M_B([U/G], 2) \times T$, and define

$$\mathcal{F}_Q := \mathcal{F} \times_{M_B([U/G], 2) \times T} (Q \times T)$$

which is finite over T . On the other hand, a point $(\mathbb{W}, \mathbb{L})$ with $\mathbb{L} \in T^{\text{bc}}$ and $((\pi^*\mathbb{W})|_U)^\vee$ is a rank two G -equivariant quotient of $\mathcal{H}_{\mathbb{L}}^m$ for some m must be in the image of Φ_m by Remark 5.20. As a result it must be in $\mathcal{F}_Q$. $\square$

Proposition 5.23. *If $\text{pr}_2 : C \rightarrow T$ is generically finite, then it must be finite and surjective.*

Proof. There is an Zariski open subset $T^\circ \subseteq T$ such that

- $T^\circ \subseteq T^{\text{bc}}$;
- $\tilde{\Sigma}_k^i|_{\mathbb{L}}$ is finite or empty for every $\mathbb{L} \in T^\circ$;
- every $(\mathbb{W}, \mathbb{L}) \in C$ with $\mathbb{L} \in T^\circ$ has $\mathbb{W}$ irreducible.

Let $(\mathbb{W}, \mathbb{L})$ be a closed point of $C \times_T T^\circ$. Then $\tilde{\Sigma}_k^i|_{\mathbb{L}}$ is a finite fiber. By Lemma 5.19 we get a quotient

$$\mathcal{H}_{\mathbb{L}}^m \twoheadrightarrow ((\pi^*\mathbb{W})|_U)^\vee$$

for some m . By Proposition 5.22, $(\mathbb{W}, \mathbb{L}) \in \mathcal{F}_Q$. It follows that

$$C \times_T T^\circ \subseteq \mathcal{F}_Q$$

On the other hand, since pr_2 is dominant, $C \times_T T^\circ$ is dense in C . Taking closures in $Q \times T$ then gives C is a closed subscheme of $\mathcal{F}_Q$. Since $\mathcal{F}_Q$ is finite over T , we get that $C \rightarrow T$ is finite. Since it is dominant, it must be surjective as well. $\square$

5.5. Étaleness of $C^\nu \rightarrow T$. We have that $\text{pr}_2 : C \rightarrow T$ is finite. Let $\nu : C^\nu \rightarrow C$ be the normalization, and $\text{pr}_2' := \text{pr}_2 \circ \nu : C^\nu \rightarrow T$. Then pr_2' is also finite and surjective. Let $C^{\text{ét}} \subseteq C$ be Zariski open locus where pr_2 is étale, and denote

$$\Delta(\text{pr}_2) := \text{pr}_2(C \setminus C^{\text{ét}})_{\text{red}} \subset T$$

which is closed in T .

Lemma 5.24. *The set $\Delta(\text{pr}_2)$ is absolute $\overline{\mathbb{Q}}$ -closed in A .*

Proof. See (5.1) for the setup. By the first paragraph of the proof of Corollary 3.10, but with M_B , we have that $C \subseteq M_B(X, 2, \det = \epsilon) \times A$ and $T \subseteq A$ are both absolute $\overline{\mathbb{Q}}$ -closed. In the notation of Definition 2.15, C^σ and T^σ are both $\overline{\mathbb{Q}}$ -closed for any $\sigma \in \text{Aut}(\mathbb{C}/\mathbb{Q})$. By [Sim94, Proposition 7.8] we have an analytic isomorphism

$$\text{RH} : M_{\text{DR}}(X, r)^{\text{an}} \xrightarrow{\sim} M_B(X, r)^{\text{an}}$$

and $C_{\text{DR}} = \text{RH}^{-1}(C)$, $T_{\text{DR}} = \text{RH}^{-1}(T)$ are both algebraic by [BW20, Proposition 7.4.5]. We also have diagrams

$$\begin{array}{ccc} (C_{\text{DR}})^{\text{an}} & \xrightarrow{(\text{pr}_{\text{DR}, 2})^{\text{an}}} & (T_{\text{DR}})^{\text{an}} \\ \downarrow \text{RH} & & \downarrow \text{RH} \\ C^{\text{an}} & \xrightarrow{(\text{pr}_2)^{\text{an}}} & T^{\text{an}} \end{array} \quad \begin{array}{ccc} (C_{\text{DR}}^\sigma)^{\text{an}} & \xrightarrow{(\text{pr}_{\text{DR}, 2}^\sigma)^{\text{an}}} & (T_{\text{DR}}^\sigma)^{\text{an}} \\ \downarrow \text{RH} & & \downarrow \text{RH} \\ (C^\sigma)^{\text{an}} & \xrightarrow{(\text{pr}_2^\sigma)^{\text{an}}} & (T^\sigma)^{\text{an}} \end{array} \quad (5.9)$$

where $(\mathrm{pr}_{\mathrm{DR},2}^\sigma)^{\mathrm{an}}$ is induced by $(\mathrm{pr}_{\mathrm{DR},2})^{\mathrm{an}}$ through an isomorphism of schemes

$$\sigma : M_{\mathrm{DR}}(X, r) \xrightarrow{\sim} M_{\mathrm{DR}}(X^\sigma, r)$$

which is itself the pullback along the base change $X^\sigma \rightarrow X$.

We have that analytification detects étaleness [Sta26, Last paragraph of Tag 025L], i.e. an algebraic map is étale if and only if its analytification is a local isomorphism. We also have that $(\mathrm{pr}_{\mathrm{DR},2}^\sigma)^{\mathrm{an}}$ is a local isomorphism if and only if $(\mathrm{pr}_{\mathrm{DR},2})^{\mathrm{an}}$ is one. It follows that $\Delta(\mathrm{pr}_2)^\sigma = \Delta(\mathrm{pr}_2^\sigma)$. Thus $\Delta(\mathrm{pr}_2)$ is absolute $\overline{\mathbb{Q}}$ -closed. $\square$

Lemma 5.25. *If pr_2^ν is not étale, then there is a prime divisor $E \subset C^\nu$ such that pr_2^ν is ramified at the generic point η_E of E . Furthermore, $\mathrm{pr}_2^\nu(E)$ is an irreducible component of $\Delta(\mathrm{pr}_2)$.*

Proof. The existence of η_E follows from (5) of [Sta26, Tag 0BMB] since pr_2^ν satisfies all (1-4). Since pr_2^ν is ramified at η_E , pr_2 must not be étale at $\nu(\eta_E)$ because otherwise C is smooth at $\nu(\eta_E)$ (since T is smooth) and ν is an isomorphism at η_E . Thus $\mathrm{pr}_2^\nu(E) \subseteq \Delta(\mathrm{pr}_2)$. A finite map preserves dimensions, so $\mathrm{pr}_2^\nu(E)$ is a prime divisor in T hence must be an irreducible component of $\Delta(\mathrm{pr}_2)$. $\square$

Proposition 5.26. *The finite map $\mathrm{pr}_2^\nu : C^\nu \rightarrow T$ is étale.*

Proof. Suppose otherwise, then by Lemma 5.25 we get a prime divisor $E \subset C^\nu$ such that pr_2^ν is ramified at every point of E , and $\mathrm{pr}_2^\nu(E) = T_0$ is an irreducible component of $\Delta(\mathrm{pr}_2)$. Since $\Delta(\mathrm{pr}_2)$ is absolute $\overline{\mathbb{Q}}$ -closed, the same is true for T_0 and by Theorem 2.16, T_0 is a torsion-translated subtorus of A .

Since C^ν is normal, the singular locus is codimension 2, and C^ν is smooth on a Zariski open subset of E . T_0 is a torsion-translated subtorus, hence unitary points are Zariski dense in T_0 . It follows that there exists a point $c \in E \subset C^\nu$ lying over $(\mathbb{V}, \mathbb{L})$ such that $\mathrm{pr}_2^\nu(c) = \mathbb{L} \in T_0$ is unitary. We have a $\mathbb{C}^*$ -action on $M_B(X, 2) \times A \supset C$. Now, C is the good moduli image of a Hodge substack of $M_B(X, 2) \times A$, hence by [BBT24, Corollary 8.24], C is $\mathbb{C}^*$ -stable. Now look at the $\mathbb{C}^*$ -orbit of $(\mathbb{V}, \mathbb{L})$. Since $\mathbb{L}$ is unitary, it is $\mathbb{C}^*$ -fixed, hence the image under pr_2 of this orbit is just $\mathbb{L}$. But $\mathrm{pr}_2 : C \rightarrow T$ is finite, so the $\mathbb{C}^*$ -orbit is finite, thus $(\mathbb{V}, \mathbb{L})$ must be $\mathbb{C}^*$ -fixed to begin with. To summarize, we have $c \in C^\nu$ a smooth point lying over $(\mathbb{V}, \mathbb{L})$ with both $\mathbb{V}, \mathbb{L}$ underlying weight 0 variation of complex Hodge structures. By Corollary 3.17 and Corollary 3.12 we get a morphism of pro- $\mathbf{T}(0)$ -MHS-algebras

$$\mathrm{pr}_{2,c}^{\nu,\#} : \hat{\mathcal{O}}_{T,\mathbb{L}} \rightarrow \hat{\mathcal{O}}_{C,(\mathbb{V},\mathbb{L})} \rightarrow \hat{\mathcal{O}}_{C^\nu,c}$$

which is a finite local map between regular complete local rings of the same dimension. By [Sta26, Tag 00R4] it must be flat, hence faithfully flat. In particular $\mathrm{pr}_{2,c}^{\nu,\#}$ is injective. Next look at the map on cotangent spaces

$$d(\mathrm{pr}_2^\nu)_c^\vee : \mathfrak{m}_{T,\mathbb{L}}/\mathfrak{m}_{T,\mathbb{L}}^2 \rightarrow \mathfrak{m}_{C^\nu,c}/\mathfrak{m}_{C^\nu,c}^2$$

which must have nontrivial kernel since pr_2^ν is ramified at c . Take the functorial Deligne weight splitting (similar to the proof of Lemma 3.13) to talk about homogeneous elements. Notice that $\mathfrak{m}_{T,\mathbb{L}}/\mathfrak{m}_{T,\mathbb{L}}^2$ is pure of weight -1 . Pick a homogeneous weight -1 element in the kernel of $d(\mathrm{pr}_2^\nu)_c^\vee$. Its image must either be 0 or also

homogeneous of weight -1 . The latter is impossible since $\mathfrak{m}_{C^\nu, c}^2 \subseteq W_{-2}\hat{\mathcal{O}}_{C^\nu, c}$, and the former is also impossible since $\hat{\mathcal{O}}_{T, \mathbb{L}} \rightarrow \hat{\mathcal{O}}_{C^\nu, c}$ is injective. Thus we reach a contradiction, and $\mathrm{pr}_2^\nu : C^\nu \rightarrow T$ is étale. $\square$

Corollary 5.27. *C^ν is an algebraic torus.*

REFERENCES

- [AHR20] Jarod Alper, Jack Hall, and David Rydh. A Luna étale slice theorem for algebraic stacks. In: *Annals of mathematics* 191.3 (2020), pp. 675–738.
- [AM69] Michael F. Atiyah and Ian G. Macdonald. *Introduction to Commutative Algebra*. Addison-Wesley Publishing Co., 1969.
- [BBT24] Benjamin Bakker, Yohan Brunebarbe, and Jacob Tsimerman. *The linear Shafarevich conjecture for quasiprojective varieties and algebraicity of Shafarevich morphisms*. 2024. arXiv: [2408.16441 \[math.AG\]](#).
- [BW15] Nero Budur and Botong Wang. Cohomology jump loci of differential graded Lie algebras. In: *Compositio Mathematica* 151.8 (2015), pp. 1499–1528.
- [BW20] Nero Budur and Botong Wang. Absolute sets and the decomposition theorem. In: *Annales scientifiques de l’Ecole normale supérieure* 53.2 (2020), pp. 469–536.
- [Cor88] Kevin Corlette. Flat G -bundles with canonical metrics. In: *Journal of differential geometry* 28.3 (1988), pp. 361–382.
- [CS08] Kevin Corlette and Carlos Simpson. On the classification of rank-two representations of quasiprojective fundamental groups. In: *Compositio Mathematica* 144.5 (2008), pp. 1271–1331.
- [Dim04] Alexandru Dimca. *Sheaves in Topology*. Universitext. Springer Berlin, Heidelberg, 2004.
- [GR84] Hans Grauert and Reinhold Remmert. *Coherent Analytic Sheaves*. Vol. 265. Grundlehren der mathematischen Wissenschaften. Springer-Verlag, 1984.
- [Har77] Robin Hartshorne. *Algebraic Geometry*. Vol. 52. Graduate Texts in Mathematics. Springer, 1977. ISBN: 978-0-387-90244-9.
- [HS04] Mark Haiman and Bernd Sturmfels. Multigraded Hilbert schemes. In: *Journal of Algebraic Geometry* 13 (2004), pp. 725–769.
- [Lun73] Domingo Luna. Slices étales. In: *Mémoires de la Société mathématique de France* 1 (1973), pp. 81–105.
- [Moc15] Takuro Mochizuki. *Mixed Twistor-Modules*. Springer, 2015.
- [Noo04] Behrang Noohi. Fundamental groups of algebraic stacks. In: *Journal of the Institute of Mathematics of Jussieu* 3.1 (2004), pp. 69–103.
- [Noo05] Behrang Noohi. *Foundations of Topological Stacks I*. 2005. arXiv: [math/0503247 \[math.AG\]](#).
- [NPT08] Khuong An Nguyen, Marius van der Put, and Jakob Top. Algebraic subgroups of $\mathrm{GL}(2)(\mathbb{C})$. In: *Indagationes mathematicae-New series* 19.2 (2008), pp. 287–297.
- [PS08] Chris A. M. Peters and Joseph H. M. Steenbrink. *Mixed Hodge Structures*. Vol. 52. Springer Berlin Heidelberg, 2008.

- [Sab05] Claude Sabbah. Polarizable twistor $\mathcal{D}$ -modules. In: *Astérisque* 300 (2005).
- [Sai90] Morihiko Saito. Mixed hodge modules. In: *Publications of the Research Institute for Mathematical Sciences* 26.2 (1990), pp. 221–333.
- [Sim91] C Simpson. The ubiquity of variations of Hodge structure. In: *Proc. Symp. Pure Math.* Vol. 53. 1991, pp. 329–348.
- [Sim92] Carlos T Simpson. Higgs bundles and local systems. In: *Publications Mathématiques de l’IHÉS* 75 (1992), pp. 5–95.
- [Sim93] Carlos Simpson. Subspaces of moduli spaces of rank one local systems. In: *Annales scientifiques de l’Ecole normale supérieure*. Vol. 26. 3. 1993, pp. 361–401.
- [Sim94] Carlos T. Simpson. Moduli of representations of the fundamental group of a smooth projective variety II. In: *Publications Mathématiques de l’IHÉS* 80 (1994), pp. 5–79.
- [Sim96] Carlos Simpson. *The Hodge filtration on nonabelian cohomology*. 1996. arXiv: [alg-geom/9604005](https://arxiv.org/abs/alg-geom/9604005) [[alg-geom](#)].
- [Sim97] Carlos Simpson. *Mixed twistor structures*. 1997. arXiv: [alg-geom/9705006](https://arxiv.org/abs/alg-geom/9705006) [[alg-geom](#)].
- [Sta26] The Stacks project authors. *The Stacks project*. <https://stacks.math.columbia.edu>. 2026.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO, ILLINOIS, USA

Email address: quocanh@uchicago.edu

URL: <https://quocanh-tran.github.io/>