



[Home](#) → [Medical Encyclopedia](#) → Doxepin overdose

URL of this page: [//medlineplus.gov/ency/article/002547.htm](https://medlineplus.gov/ency/article/002547.htm)

## Doxepin overdose

Doxepin is a type of medicine called a tricyclic antidepressant (TCA). It is prescribed to treat depression and anxiety. Doxepin overdose occurs when someone takes more than the normal or recommended amount of this medicine, either by accident or on purpose. Toxic level of a TCA can build up in the body if the TCA and other medicines interact. This interaction can affect how well the body can break down the TCA.

This article is for information only. DO NOT use it to treat or manage an actual overdose. If you or someone you are with overdoses, call your local emergency number (such as 911), or your local poison control center can be reached directly by calling the national toll-free Poison Help hotline (1-800-222-1222) from anywhere in the United States.

### Poisonous Ingredient

Doxepin

### Where Found

These medicines contain doxepin:

- Silenor
- Sinequan
- Zonalon

Other medicines may also contain doxepin.

### Symptoms

Below are symptoms of an overdose of doxepin in different parts of the body:

#### AIRWAYS AND LUNGS

- Slow breathing
- Difficulty breathing

#### BLADDER AND KIDNEYS

- Hard to start urinating

- Hard to empty bladder

## EYES, EARS, NOSE, AND THROAT

- Blurred vision
- Ringing in the ears

## HEART AND BLOOD

- Irregular heartbeat (may be fatal)
- Low blood pressure
- Shock

## MOUTH, STOMACH, AND INTESTINAL TRACT

- Constipation
- Dry mouth
- Nausea and vomiting
- Unpleasant taste in mouth

## NERVOUS SYSTEM

- Agitation, confusion
- Drowsiness, decreased alertness, coma
- Headache
- Muscle rigidity, lack of coordination
- Restlessness
- Seizures

## SKIN

- Very sensitive to sunlight

## Home Care

Get medical help right away. DO NOT make the person throw up unless poison control or a health care provider tells you to.

## Before Calling Emergency

Have this information ready:

- Person's age, weight, and condition
- Name of the medicine and the strength of the medicine, if known
- Time it was swallowed
- Amount swallowed

- If the medicine was prescribed for the person

## Poison Control

Your local poison control center can be reached directly by calling the national toll-free Poison Help hotline (1-800-222-1222) from anywhere in the United States. This national hotline number will let you talk to experts in poisoning. They will give you further instructions.

This is a free and confidential service. All local poison control centers in the United States use this national number. You should call if you have any questions about poisoning or poison prevention. It does NOT need to be an emergency. You can call for any reason, 24 hours a day, 7 days a week.

## What to Expect at the Emergency Room

Take the container with you to the hospital, if possible.

The provider will measure and monitor the person's vital signs, including temperature, pulse, breathing rate, and blood pressure. Symptoms will be treated. The person may receive:

- Activated charcoal
- Blood and urine tests
- Breathing support, including oxygen and a tube through the mouth into the lungs
- Chest x-ray
- CT scan of the brain
- ECG (electrocardiogram, or heart tracing)
- Intravenous fluids (given through a vein)
- Laxative
- Medicine to treat symptoms
- Sodium bicarbonate, to counteract the effects of TCA overdose
- Catheter (thin, flexible tube) into the bladder if the person cannot urinate on their own

## Outlook (Prognosis)

How well a person does depends on the amount of medicine they swallowed and how quickly the treatment is received. The faster a person gets medical help, the better the chance for recovery.

Tricyclic depressant overdoses are very toxic and difficult to treat. Many people have died from TCA overdose, even with aggressive medical treatment.

## Alternative Names

Doxepin hydrochloride overdose

## References

Aronson JK. Doxepin. In: Aronson JK, ed. *Meyler's Side Effects of Drugs*. 16th ed. Waltham, MA: Elsevier; 2016:1084.

Levine MD, Ruha AM. Antidepressants. In: Walls RM, ed. *Rosen's Emergency Medicine: Concepts and Clinical Practice*. 10th ed. Philadelphia, PA: Elsevier; 2023:chap 141.

## Review Date 4/1/2023

Updated by: Jesse Borke, MD, CPE, FAAEM, FACEP, Attending Physician at Kaiser Permanente, Orange County, CA. Also reviewed by David C. Dugdale, MD, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team.

[Learn how to cite this page](#)



Health Content  
Provider  
06/01/2028

A.D.A.M., Inc. is accredited by [URAC](#), for Health Content Provider ([www.urac.org](#)). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#), and [privacy policy](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Links to other sites are provided for information only – they do not constitute endorsements of those other sites. No warranty of any kind, either expressed or implied, is made as to the accuracy, reliability, timeliness, or correctness of any translations made by a third-party service of the information provided herein into any other language. © 1997-2025 A.D.A.M., a business unit of Ebix, Inc. Any duplication or distribution of the information contained herein is strictly prohibited.



National Library of Medicine 8600 Rockville Pike, Bethesda, MD 20894 U.S. Department of Health and Human Services  
National Institutes of Health