



[Home](#) → [Medical Encyclopedia](#) → C1 esterase inhibitor

URL of this page: //medlineplus.gov/ency/article/003353.htm

C1 esterase inhibitor

C1 esterase inhibitor (C1-INH) is a protein found in the fluid part of your blood. It controls a protein called C1, which is part of the complement system.

The complement system is a group of nearly 60 proteins in blood plasma or on the surface of some cells. The complement proteins work with your immune system to protect the body from infections. They also help remove dead cells and foreign material. There are nine major complement proteins. They are labeled C1 through C9. Rarely, people may inherit deficiency of some complement proteins. These people are prone to certain infections or autoimmune disorders.

This article discusses the test that is done to measure the amount of C1-INH in your blood.

How the Test is Performed

A blood sample is needed. This is most often taken through a vein. The procedure is called a venipuncture.

How to Prepare for the Test

No special preparation is needed.

How the Test will Feel

When the needle is inserted to draw blood, some people feel moderate pain. Others may feel only a prick or stinging sensation. Afterward, there may be some throbbing.

Why the Test is Performed

You may have this test if you have signs of hereditary or acquired angioedema, which is a type of tissue swelling. Both forms of angioedema are caused by low levels of C1-INH.

Complement factors may also be important in testing for autoimmune diseases, such as systemic lupus erythematosus.

Normal Results

Normal value ranges may vary slightly among different laboratories. Your health care provider will also measure the functional activity level of your C1 esterase inhibitor. Talk to your provider about the meaning of your specific test results.

What Abnormal Results Mean

Low levels of C1-INH may cause certain types of angioedema. Angioedema results in sudden swelling of the tissues of the face, upper throat and tongue. It may also cause difficulty breathing. Swelling in the intestine and abdominal pain may also occur. There are two types of angioedema that result from decreased levels of C1-INH. Hereditary angioedema affects children and young adults under age 20. Acquired angioedema is seen in adults older than age 40. Adults with acquired angioedema are much more likely to also have other conditions, such as cancer or autoimmune disease.

Risks

Risks associated with having blood drawn are slight, but may include:

- Excessive bleeding
- Fainting or feeling lightheaded
- Hematoma (blood accumulating under the skin)
- Infection (a slight risk any time the skin is broken)

Alternative Names

C1 inhibiting factor; C1-INH

References

Atkinson JP, Liszewski MK, Java A, Kulkarni HS. Complement and complement disorders. In: Rich RR, Fleisher TA, Schroeder HW, Weyand CM, Corry DB, Puck JM, eds. *Clinical Immunology: Principles and Practice*. 6th ed. Philadelphia, PA: Elsevier; 2023:chap 40.

Leslie TA. Hereditary angioedema. In: Lebwohl MG, Heymann WR, Coulson IH, Murrell DF, eds. *Treatment of Skin Disease: Comprehensive Therapeutic Strategies*. 6th ed. Philadelphia, PA: Elsevier; 2022:chap 101.

Saini SS. Urticaria and angioedema. In: Goldman L, Cooney KA, eds. *Goldman-Cecil Medicine*. 27th ed. Philadelphia, PA: Elsevier; 2024:chap 232.

Review Date 1/28/2025

Updated by: Diane M. Horowitz, MD, Rheumatology and Internal Medicine, Northwell Health, Great Neck, NY. Review provided by VeriMed Healthcare Network. Also reviewed by David C. Dugdale, MD, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team.

[Learn how to cite this page](#)



A.D.A.M., Inc. is accredited by URAC, for Health Content Provider (www.urac.org). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#), and [privacy policy](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Links to other sites are provided for information only – they do not constitute endorsements of those other sites. No warranty of any kind, either expressed or implied, is made as to the accuracy, reliability, timeliness, or correctness of any translations made by a third-party service of the information provided herein into any other language. © 1997-2025 A.D.A.M., a business unit of Ebix, Inc. Any duplication or distribution of the information contained herein is strictly prohibited.



National Library of Medicine 8600 Rockville Pike, Bethesda, MD 20894 U.S. Department of Health and Human Services

National Institutes of Health