



[Home](#) → [Medical Encyclopedia](#) → Costochondritis

URL of this page: [//medlineplus.gov/ency/article/000164.htm](https://medlineplus.gov/ency/article/000164.htm)

## Costochondritis

All but your lowest 2 ribs are connected to your breastbone by cartilage. This cartilage can become inflamed and cause pain. This condition is called costochondritis. It is a common cause of chest pain.

### Causes

There is often no known cause of costochondritis. But it may be caused by:

- Chest injury
- Hard exercise or heavy lifting
- Viral infections, such as respiratory infections
- Strain from coughing
- Infections after surgery or from intravenous (IV) drug use
- Some types of arthritis

### Symptoms

The most common symptoms of costochondritis are pain and tenderness in the chest. You may feel:

- Sharp pain at the front of your chest wall, which may move to your back or stomach
- Increased pain when you take a deep breath or cough
- Tenderness when you press the area where the rib joins the breastbone
- Less pain when you stop moving and breathe quietly

### Exams and Tests

Your health care provider will take your medical history and do a physical exam. The area where the ribs meet the breastbone is checked. If this area is tender and sore, costochondritis is the most likely cause of your chest pain.

A chest x-ray may be done if your symptoms are severe or do not improve with treatment.

Your provider may also order tests to check for other conditions, such as a heart attack.

## Treatment

Costochondritis most often goes away on its own in a few days or weeks. It can also take up to a few months. Treatment focuses on relieving the pain.

- Apply hot or cold compresses.
- Avoid activities that make the pain worse.

Pain medicines, such as ibuprofen (Advil, Motrin) or naproxen (Aleve), may help to ease pain and swelling. You can buy these without a prescription.

- Talk with your provider before using these medicines if you have heart disease, high blood pressure, kidney disease, liver disease, or have had stomach ulcers or internal bleeding in the past.
- Take the dose as advised by the provider. Do not take more than the amount recommended on the bottle. Carefully read the warnings on the label before taking any medicine.

You may also take acetaminophen (Tylenol) instead, if your provider tells you it is safe to do so. People with liver disease should not take this medicine.

If your pain is severe, your provider may prescribe stronger pain medicine.

In some cases, your provider may recommend physical therapy.

## Outlook (Prognosis)

Costochondritis pain often goes away in a few days or weeks.

## When to Contact a Medical Professional

Call 911 or the local emergency number, or go to your local emergency room right away if you have chest pain. The pain of costochondritis can be similar to the pain of a heart attack.

If you have already been diagnosed with costochondritis, contact your provider if you have any of the following symptoms:

- Trouble breathing
- A high fever
- Any signs of infection such as pus, redness, or swelling around your ribs
- Pain that continues or gets worse after taking pain medicine
- Sharp pain with every breath

## Prevention

Because the cause is often unknown, there is no known way to prevent costochondritis.

## Alternative Names

Chest wall pain; Costosternal syndrome; Costosternal chondrodynia; Chest pain - costochondritis

# References

Crasto JA, Vaswani RS, Pauyo T, Musahl V. Overview of sport-specific injuries. In: Miller MD, Thompson SR. eds. *DeLee, Drez, & Miller's Orthopaedic Sports Medicine*. 5th ed. Philadelphia, PA: Elsevier; 2020:chap 9.

Hanak JA. Tietze syndrome. In: Frontera WR, Silver JK, Rizzo TD Jr, eds. *Essentials of Physical Medicine and Rehabilitation: Musculoskeletal Disorders, Pain, and Rehabilitation*. 4th ed. Philadelphia, PA: Elsevier; 2019:chap 117.

Kurz J. Costosternal syndrome. In: Frontera WR, Silver JK, Rizzo TD Jr, eds. *Essentials of Physical Medicine and Rehabilitation: Musculoskeletal Disorders, Pain, and Rehabilitation*. 4th ed. Philadelphia, PA: Elsevier; 2019:chap 101.

## Review Date 10/9/2024

Updated by: Linda J. Vorvick, MD, Clinical Professor, Department of Family Medicine, UW Medicine, School of Medicine, University of Washington, Seattle, WA. Also reviewed by David C. Dugdale, MD, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team.

Learn how to cite this page



Health Content  
Provider  
06/01/2028

A.D.A.M., Inc. is accredited by [URAC](#), for Health Content Provider ([www.urac.org](#)). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#), and [privacy policy](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Links to other sites are provided for information only – they do not constitute endorsements of those other sites. No warranty of any kind, either expressed or implied, is made as to the accuracy, reliability, timeliness, or correctness of any translations made by a third-party service of the information provided herein into any other language. © 1997-2025 A.D.A.M., a business unit of Ebix, Inc. Any duplication or distribution of the information contained herein is strictly prohibited.

