



[Home](#) → [Medical Encyclopedia](#) → Excessive crying in infants

URL of this page: //medlineplus.gov/ency/article/003023.htm

Excessive crying in infants

Crying is an important way for infants to communicate. But, when a baby cries a lot, it may be a sign of something that needs treatment.

Considerations

Infants normally cry about 1 to 3 hours a day. It is perfectly normal for an infant to cry when hungry, thirsty, tired, lonely, or in pain. It is also normal for a baby to have a fussy period in the evening.

But, if an infant cries too often, there might be a health problem that needs attention.

Causes

Infants may cry because of any of the following:

- Boredom or loneliness
- Colic
- Discomfort or irritation from a wet or dirty diaper, excessive gas, or feeling cold
- Hunger or thirst
- Illness
- Infection (a likely cause if the crying is accompanied by irritability, lethargy, poor appetite, or fever. You should contact your baby's health care provider)
- Medicines
- Normal muscle jerks and twitches that disturb the sleep
- Pain
- Teething

Home Care

Home care depends on the causes. Follow your provider's advice.

If the infant seems constantly hungry despite short, frequent feedings, talk to your provider about normal growth and feeding times.

If crying is due to boredom or loneliness, it may be helpful to touch, hold, and talk to the infant more and place the infant within sight. Place baby-safe toys where the child can see them. If crying is due to sleep disturbance, wrap the baby firmly in a blanket before putting the infant to bed.

For excessive crying in infants due to cold, dress the infant warmly or adjust the temperature of the room. If adults are cold, the baby is also likely cold.

Always check for possible causes of pain or discomfort in a crying baby. When cloth diapers are used, look for diaper pins that have become loose or loose threads that have become tightly wrapped around fingers or toes. Diaper rashes also can be uncomfortable.

Take your baby's temperature to check for fever. Check your baby head-to-toe for any injuries. Pay particular attention to the fingers, toes, and genitalia. It is not uncommon for a hair to get wrapped around part of your baby, such as a toe, creating pain.

When to Contact a Medical Professional

Contact your child's provider if:

- A baby's excessive crying remains unexplained and does not go away in 1 day, despite attempts at home treatment
- The baby has other symptoms, such as fever, along with the excessive crying

What to Expect at Your Office Visit

Your provider will examine your baby and ask about the child's medical history and symptoms. Questions may include:

- Is the child teething?
- Is the child bored, lonely, hungry, thirsty?
- Does the child seem to have a lot of gas?
- What other symptoms does the child have? Such as, difficulty waking up, fever, irritability, poor appetite, or vomiting?

Your provider will check the infant's growth and development. Antibiotics may be prescribed if the baby has a bacterial infection.

Alternative Names

Infants - excessive crying; Well child - excessive crying

References

Marcdante KJ, Kliegman RM, Schuh AM. Crying and colic. In: Marcante KJ, Kliegman RM, Schuh AM, eds. *Nelson Essentials of Pediatrics*. 9th ed. Philadelphia, PA: Elsevier; 2023:chap 11.

Onigbanjo MT, Feigelman S. The first year. In: Kliegman RM, St. Geme JW, Blum NJ, et al, eds. *Nelson Textbook of Pediatrics*. 22nd ed. Philadelphia, PA: Elsevier; 2025:chap 23.

Pomeranz AJ, Sabnis S, Busey SL, Kliegman RM. Irritable infant (fussy or excessively crying infant). In: Pomeranz AJ, Sabnis S, Busey SL, Kliegman RM, eds. *Pediatric Decision-Making Strategies*. 2nd ed. Philadelphia, PA: Elsevier Saunders; 2016:chap 79.

Review Date 10/20/2024

Updated by: Neil K. Kaneshiro, MD, MHA, Clinical Professor of Pediatrics, University of Washington School of Medicine, Seattle, WA. Also reviewed by David C. Dugdale, MD, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team.


Learn how to cite this page



CERTIFIED
Health Content
Provider
06/01/2028

A.D.A.M., Inc. is accredited by URAC, for Health Content Provider (www.urac.org). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#), and [privacy policy](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Links to other sites are provided for information only – they do not constitute endorsements of those other sites. No warranty of any kind, either expressed or implied, is made as to the accuracy, reliability, timeliness, or correctness of any translations made by a third-party service of the information provided herein into any other language. © 1997-2025 A.D.A.M., a business unit of Ebix, Inc. Any duplication or distribution of the information contained herein is strictly prohibited.



National Library of Medicine 8600 Rockville Pike, Bethesda, MD 20894 U.S. Department of Health and Human Services

National Institutes of Health