

Combined Hormonal Birth Control: Pill, Patch, and Ring

Overview

What are

Birth control
hormonal
progestin

How do

Combined



We Use Cookies

ACOG uses cookies, pixels and similar technologies to personalize your website experience. By clicking “continue” or continuing to use our site, you agree to our [Privacy Policy](#).

Continue

whole body. These hormones prevent pregnancy mainly by stopping [ovulation](#) (the release of an egg from one of the [ovaries](#)). They also cause other changes in the body that help prevent pregnancy. The mucus in the [cervix](#) thickens, making it hard for sperm to enter the [uterus](#). The lining of the uterus also thins.

How effective are combined hormonal birth control methods?

With typical use—meaning that the method may not always be used consistently or correctly—9 in 100 women will get pregnant during the first year of using these methods. With perfect use—meaning that the method is used consistently and correctly each time—fewer than 1 in 100 women will get pregnant during the first year.

What are the benefits of combined hormonal methods?

Combined hormonal methods have several benefits in addition to protecting against pregnancy:

- They may make your period more regular, lighter, and shorter.
- They help reduce menstrual cramps.
- They decrease the risk of cancer of the uterus, ovary, and colon.
- They may improve acne and reduce unwanted hair growth.
- They can be used to treat certain disorders that cause heavy bleeding and menstrual pain, such as [fibroids](#) and [endometriosis](#).
- Used continuously, they can reduce the frequency of migraines associated with menstruation (although they should not be used if you have migraines with [aura](#)).

The
per

We Use Cookies

ACOG uses cookies, pixels and similar technologies to personalize your website experience. By clicking "continue" or continuing to use our site, you agree to our [Privacy Policy](#).

What a

Combi
small i
higher

- are
- have multiple risk factors for [cardiovascular disease](#), such as high cholesterol, high blood pressure, and diabetes
- have a history of stroke, heart attack, or DVT
- have a history of migraine headaches with aura

You should not use combined hormonal methods during the first 3 weeks after delivery because the risk of DVT is higher in the weeks after childbirth. If you have additional risk factors for DVT, you should wait to use combined hormonal methods until after the first 4 to 6 weeks following delivery.

The risk of DVT may also be slightly higher if you are taking pills containing a progestin called drospirenone or if you are using the patch. But the risk of DVT is higher during

pregnancy and in the weeks after childbirth than when taking drospirenone-containing pills or using the patch.

Can I use combined hormonal birth control methods while I am breastfeeding?

If you are breastfeeding, estrogen may affect your milk supply. It is recommended that you wait until the fifth week after delivery to start using these methods, when breastfeeding has been well established.

Combined Hormonal Birth Control Pills

How can I get combined hormonal birth control pills?

In the
only.

What a
are the

- 21-day
start
per
- 28-day
bra
rem

We Use Cookies

ACOG uses cookies, pixels and similar technologies to personalize your website experience. By clicking “continue” or continuing to use our site, you agree to our [Privacy Policy](#).

such as iron, but no hormones; or inactive pills (containing no hormones or supplements). During the days you are taking the hormone-free pills, you will have your period.

- 90-day pills—Take one pill at the same time each day for 84 days. Depending on the brand, the last seven pills either contain no hormones or contain estrogen only. With both brands, you will have your period on the last 7 days every 3 months.
- 365-day pills—Take one pill at the same time each day for a year. In time, your bleeding may become lighter and may even stop.

What are possible side effects of using the combined hormonal birth control pill?

Most side effects are minor and often go away after a few months of use. Possible side effects include

- headache
- nausea
- breast tenderness
- [breakthrough bleeding](#)

[Breakthrough bleeding](#) usually is a temporary side effect as the body adjusts to a change in hormone levels. It may last longer than a few months with continuous-dose pills.

Vaginal

What is

The vaginal
estrogen

We Use Cookies

ACOG uses cookies, pixels and similar technologies to personalize your website experience. By clicking “continue” or continuing to use our site, you agree to our [Privacy Policy](#).

How can I get the vaginal ring?

A health care professional must prescribe the vaginal ring, but you insert it yourself.

How do I use the vaginal ring?

You fold the ring and insert it into the vagina. It stays there for 21 days. You then remove it for 7 days. During the week the ring is not used, you will have your period.

There are two types of vaginal rings. One type of vaginal ring requires inserting a new ring each month. Another type of vaginal ring can be reused for up to a year. The reusable ring should be washed and stored in its case when not in use.

If you use the monthly ring, you can choose to skip the ring-free week. Just insert a new ring every 21 days with no ring-free week in between. This is called continuous use. With continuous use, your period may become lighter and even stop. There are not yet studies on continuous use of the reusable ring.

What are possible side effects of using the vaginal ring?

Most side effects are minor and often go away after a few months of use. Possible side effects include

- headache
- nausea
- breast tenderness
- vaginal irritation
- vaginal discharge
- breast tenderness

We Use Cookies

ACOG uses cookies, pixels and similar technologies to personalize your website experience. By clicking “continue” or continuing to use our site, you agree to our [Privacy Policy](#).

Birth

What is

The birth

skin to

absorbed through the skin into the body.

How can I get the birth control patch?

A health care professional must prescribe the patch, but you do not need to visit a health care professional to apply or remove the patch. The patch is less effective in women who weigh more than 198 pounds.

How do I use the birth control patch?

The patch can be worn on the buttocks, chest (except the breasts), upper back or arm, or abdomen. You wear a patch for a week at a time for a total of 3 weeks in a row. During the fourth week, a patch is not worn, and you will have your period.

After week 4, a new patch is applied and the cycle is repeated. You apply the patch on the same day of the week even if you still are bleeding.

To use the patch as a continuous-dose form of birth control, apply a new patch every week on the same day without skipping a week.

What are possible side effects of using the birth control patch?

Most side effects are minor and often go away after a few months of use. Possible side effects include

- skin irritation
- breast tenderness
- headache
- breast tenderness

We Use Cookies

ACOG uses cookies, pixels and similar technologies to personalize your website experience. By clicking “continue” or continuing to use our site, you agree to our [Privacy Policy](#).

Glossary

Aura: A
migraine
particu

Breakt

Cardiovascular Disease: Disease of the heart and blood vessels.

Cervix: The lower, narrow end of the uterus at the top of the vagina.

Deep Vein Thrombosis (DVT): A condition in which a blood clot forms in veins in the leg or other areas of the body.

Endometriosis: A condition in which tissue that lines the uterus is found outside of the uterus, usually on the ovaries, fallopian tubes, and other pelvic structures.

Estrogen: A female hormone produced in the ovaries.

Fibroids: Growths that form in the muscle of the uterus. Fibroids usually are noncancerous.

Hormones: Substances made in the body that control the function of cells or organs.

Ovaries: Organs in women that contain the eggs necessary to get pregnant and make important hormones, such as estrogen, progesterone, and testosterone.

Ovulation: The time when an ovary releases an egg.

Progestin: A synthetic form of progesterone that is similar to the hormone made naturally by the body.

Uterus

nouris

If you

Don't h

FAQ185

Last up

Last reviewed: May 2020

We Use Cookies

ACOG uses cookies, pixels and similar technologies to personalize your website experience. By clicking “continue” or continuing to use our site, you agree to our [Privacy Policy](#).

Copyright 2025 by the American College of Obstetricians and Gynecologists. All rights reserved. Read [copyright and permissions information](#).

This information is designed as an educational aid for the public. It offers current information and opinions related to women's health. It is not intended as a statement of the standard of care. It does not explain all of the proper treatments or methods of care. It is not a substitute for the advice of a physician. Read [ACOG's complete disclaimer](#).

About ACOG

Disclaimer

Contact Us

How to Find an Ob-Gyn



We Use Cookies

ACOG uses cookies, pixels and similar technologies to personalize your website experience. By clicking “continue” or continuing to use our site, you agree to our [Privacy Policy](#).