



[Home](#) → [Medical Encyclopedia](#) → Folic acid - test

URL of this page: //medlineplus.gov/ency/article/003686.htm

## Folic acid - test

Folic acid is also called vitamin B9. This article discusses the test to measure the amount of folic acid in the blood. It is often done together with a vitamin B12 test of the blood.

### How the Test is Performed

A blood sample is needed.

### How to Prepare for the Test

You should not eat or drink for 6 hours before the test. Your health care provider may tell you to stop taking any drugs that may interfere with test results, including folic acid supplements.

Drugs that can decrease folic acid measurements include:

- Alcohol
- Aminosalicylic acid
- Birth control pills
- Estrogens
- Tetracyclines
- Ampicillin
- Chloramphenicol
- Erythromycin
- Methotrexate
- Penicillin
- Aminopterin
- Phenobarbital
- Phenytoin
- Drugs to treat malaria

### How the Test will Feel

You may feel slight pain or a little sting when the needle is inserted. There may be some throbbing at the site.

# Why the Test is Performed

This test is done to check for folic acid deficiency.

Folic acid helps form red blood cells and produce DNA that stores genetic codes. Taking the right amount of folic acid before and during pregnancy helps prevent neural tube defects, such as spina bifida.

Women who are pregnant or planning to become pregnant should take at least 600 micrograms (mcg) of folic acid every day. It may be provided by prenatal vitamins and certain foods, such as enriched bread, flour, and cereal. Some women may need to take more if they have a history of neural tube defects in earlier pregnancies. Ask your provider how much you need.

## Normal Results

The normal range is 2.7 to 17.0 nanograms per milliliter (ng/mL) or 6.12 to 38.52 nanomoles per liter (nmol/L).

Normal value ranges may vary slightly among different labs. Talk to your provider about the meaning of your test results.

The examples above show the common measurements for results for these tests. Some labs use different measurements or may test different specimens.

## What Abnormal Results Mean

Lower-than-normal folic acid levels may indicate:

- Poor diet
- Malabsorption syndrome (for example, celiac sprue)
- Malnutrition

The test may also be done in cases of:

- Anemia that may be due to folate deficiency
- Megaloblastic anemia

## Risks

There is very little risk involved with having your blood taken. Veins and arteries vary in size from one person to another and from one side of the body to the other. Taking blood from some people may be more difficult than from others.

Other slight risks from having blood drawn may include:

- Excessive bleeding
- Fainting or feeling lightheaded
- Hematoma (blood accumulating under the skin)
- Infection (a slight risk any time the skin is broken)

# Alternative Names

Folate - test

## References

Antony AC. Megaloblastic anemias. In: Hoffman R, Benz EJ, Silberstein LE, et al, eds. *Hematology: Basic Principles and Practice*. 8th ed. Philadelphia, PA: Elsevier; 2023:chap 40.

Elghetany MT, Banki K. Erythrocytic disorders. In: McPherson RA, Pincus MR, eds. *Henry's Clinical Diagnosis and Management by Laboratory Methods*. 24th ed. Philadelphia, PA: Elsevier; 2022:chap 33.

Mason JB, Booth SL. Vitamins, trace minerals, and other micronutrients. In: Goldman L, Schafer AI, eds. *Goldman-Cecil Medicine*. 26th ed. Philadelphia, PA: Elsevier; 2020:chap 205.

## Review Date 2/2/2023

Updated by: Mark Levin, MD, Hematologist and Oncologist, Monsey, NY. Review provided by VeriMed Healthcare Network. Also reviewed by David C. Dugdale, MD, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team.

### Learn how to cite this page



Health Content  
Provider  
06/01/2028

A.D.A.M., Inc. is accredited by URAC, for Health Content Provider ([www.urac.org](http://www.urac.org)). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#), and [privacy policy](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Links to other sites are provided for information only – they do not constitute endorsements of those other sites. No warranty of any kind, either expressed or implied, is made as to the accuracy, reliability, timeliness, or correctness of any translations made by a third-party service of the information provided herein into any other language. © 1997-2025 A.D.A.M., a business unit of Ebix, Inc. Any duplication or distribution of the information contained herein is strictly prohibited.



National Library of Medicine 8600 Rockville Pike, Bethesda, MD 20894 U.S. Department of Health and Human Services

National Institutes of Health