



[Home](#) → [Medical Encyclopedia](#) → Cord blood testing

URL of this page: [//medlineplus.gov/ency/article/003403.htm](https://medlineplus.gov/ency/article/003403.htm)

Cord blood testing

Cord blood refers to a sample of blood collected from the umbilical cord when a baby is born. The umbilical cord is the cord connecting the baby to the mother's womb (uterus).

Cord blood testing can be done to evaluate a newborn's health.

How the Test is Performed

Right after the birth of your baby, the umbilical cord is clamped and cut. If cord blood is to be drawn, another clamp is placed 8 to 10 inches (20 to 25 centimeters) away from the first. The section between the clamps is cut and a blood sample is collected into a specimen tube.

How to Prepare for the Test

No special steps are needed to prepare for this test.

How the Test will Feel

You will not feel anything beyond the normal birthing process.

Why the Test is Performed

Cord blood testing is done to measure the following in your baby's blood:

- Bilirubin level
- Blood culture (if an infection is suspected)
- Blood gases (including oxygen, carbon dioxide, and pH levels)
- Blood sugar level
- Blood type and Rh
- Complete blood count (CBC)
- Platelet count

Normal Results

Normal values mean that all items checked are within normal range.

What Abnormal Results Mean

A low pH (less than 7.04 to 7.10) means there are higher levels of acids in the baby's blood. This might occur when the baby does not get enough oxygen during labor. One reason for this could be that the umbilical cord was compressed during labor or delivery.

A blood culture that is positive for bacteria means your baby has a blood infection.

High level of blood sugar (glucose) in the cord blood may be found if the mother has diabetes. The newborn will be watched for hypoglycemia (low blood sugar) after delivery.

High level of bilirubin in the newborn has many causes, which could be due to infections the baby gets.

Note: Normal value ranges may vary slightly among different laboratories. Talk to your health care provider about the meaning of your specific test results.

Considerations

Most hospitals routinely collect cord blood for testing at birth. The process is fairly easy and this is the only time when this type of blood sample can be collected.

You can also decide to bank or donate cord blood at the time of your delivery. Cord blood can be used to treat certain types of bone marrow-related cancers. Some parents may choose to save (bank) their child's cord blood for this and other future medical purposes.

Cord blood banking for personal use is done by both cord blood banks and private companies. There is a charge for the service if you use a private service. If you choose to bank your infant's cord blood, you should talk to your health care provider about the pros and cons of different options.

References

American College of Obstetricians and Gynecologists. ACOG committee opinion no. 771: umbilical cord blood banking. *Obstet Gynecol*. 2019;133(3):e249-e253. PMID: 30801478
pubmed.ncbi.nlm.nih.gov/30801478/ [<https://pubmed.ncbi.nlm.nih.gov/30801478/>].

Hsu YMS, Elkins MB. Tissue banking and progenitor cells. In: McPherson RA, Pincus MR, eds. *Henry's Clinical Diagnosis and Management by Laboratory Methods*. 24th ed. Philadelphia, PA: Elsevier; 2022:chap 39.

Waldorf KMA. Maternal-fetal immunology. In: Landon MB, Galan HL, Jauniaux ERM, et al, eds. *Gabbe's Obstetrics: Normal and Problem Pregnancies*. 8th ed. Philadelphia, PA: Elsevier; 2021:chap 4.

Review Date 4/16/2024

Updated by: John D. Jacobson, MD, Professor Emeritus, Department of Obstetrics and Gynecology, Loma Linda University School of Medicine, Loma Linda, CA. Also reviewed by David C. Dugdale, MD, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team.



Health Content
Provider
06/01/2028

A.D.A.M., Inc. is accredited by [URAC](http://www.urac.org), for Health Content Provider (www.urac.org). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#), and [privacy policy](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Links to other sites are provided for information only – they do not constitute endorsements of those other sites. No warranty of any kind, either expressed or implied, is made as to the accuracy, reliability, timeliness, or correctness of any translations made by a third-party service of the information provided herein into any other language. © 1997-2025 A.D.A.M., a business unit of Ebix, Inc. Any duplication or distribution of the information contained herein is strictly prohibited.



National Library of Medicine 8600 Rockville Pike, Bethesda, MD 20894 U.S. Department of Health and Human Services
National Institutes of Health