



[Home](#) → [Medical Encyclopedia](#) → Exchange transfusion

URL of this page: [//medlineplus.gov/ency/article/002923.htm](https://medlineplus.gov/ency/article/002923.htm)

Exchange transfusion

Exchange transfusion is a potentially life-saving procedure that is done to counteract the effects of serious jaundice or changes in the blood due to diseases such as sickle cell anemia.

The procedure involves slowly removing the person's blood and replacing it with fresh donor blood or plasma.

Description

An exchange transfusion requires that the person's blood be removed and replaced. In most cases, this involves placing one or more thin tubes, called catheters, into a blood vessel. The exchange transfusion is done in cycles, each one most often lasts a few minutes.

The person's blood is slowly withdrawn (most often about 5 to 20 mL at a time, depending on the person's size and the severity of illness). An equal amount of fresh, prewarmed blood or plasma flows into the person's body. This cycle is repeated until the correct volume of blood has been replaced.

After the exchange transfusion, catheters may be left in place in case the procedure needs to be repeated.

In diseases such as sickle cell anemia, blood is removed and replaced with donor blood.

In conditions such as neonatal polycythemia, a specific amount of the child's blood is removed and replaced with a normal saline solution, plasma (the clear liquid part of blood), or albumin (a normal blood protein). This decreases the total number of red blood cells in the body and makes it easier for blood to flow through the body.

Why the Procedure is Performed

An exchange transfusion may be needed to treat the following conditions:

- Dangerously high red blood cell count in a newborn (neonatal polycythemia)
- Rh-induced hemolytic disease of the newborn
- Severe disturbances in body chemistry
- Severe newborn jaundice that does not respond to phototherapy with bili lights
- Severe sickle cell crisis
- Toxic effects of certain drugs

Risks

General risks are the same as with any transfusion. Other possible complications include:

- Blood clots
- Changes in blood chemistry (high or low potassium, low calcium, low glucose, change in acid-base balance in the blood)
- Heart and lung problems
- Infection (very low risk due to careful screening of blood)
- Shock if not enough blood is replaced

After the Procedure

The person may need to be monitored for several days in the hospital after the transfusion. The length of stay depends on what condition the exchange transfusion was performed to treat.

Alternative Names

Hemolytic disease - exchange transfusion

References

Golden WC, Watchko JF. Neonatal hyperbilirubinemia and kernicterus. In: Gleason CA, Sawyer T, eds. *Avery's Diseases of the Newborn*. 11th ed. Philadelphia, PA: Elsevier; 2024:chap 72.

Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM. Blood disorders. In: Kliegman RM, St. Geme JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, eds. *Nelson Textbook of Pediatrics*. 21st ed. Philadelphia, PA: Elsevier; 2020:chap 124.

Park J. Hematology. In: Anderson CC, Kapoor S, Mark TE, eds. *Harriet Lane Handbook, The: The Johns Hopkins Hospital*. 23rd ed. Philadelphia, PA: Elsevier; 2021:chap 14.

Rodrigue BB, Sloan SR. Pediatric transfusion medicine. In: Hoffman R, Benz EJ, Silberstein LE, et al, eds. *Hematology: Basic Principles and Practice*. 8th ed. Philadelphia, PA: Elsevier; 2023:chap 119.

Review Date 11/6/2023

Updated by: Neil K. Kaneshiro, MD, MHA, Clinical Professor of Pediatrics, University of Washington School of Medicine, Seattle, WA. Also reviewed by David C. Dugdale, MD, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team.

Learn how to cite this page



A.D.A.M., Inc. is accredited by [URAC](http://www.urac.org), for Health Content Provider (www.urac.org). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#), and [privacy policy](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Links to other sites are provided for information only – they do not constitute endorsements of those other sites. No warranty of any kind, either expressed or implied, is made as to the accuracy, reliability, timeliness, or correctness of any translations made by a third-party service of the information provided herein into any other language. © 1997-2025 A.D.A.M., a business unit of Ebix, Inc. Any duplication or distribution of the information contained herein is strictly prohibited.



National Library of Medicine 8600 Rockville Pike, Bethesda, MD 20894 U.S. Department of Health and Human Services
National Institutes of Health