

hemopneumothorax

hemopoiesis

hemoprecipitin

hemoprotein

hemopsonin

hemoptysis

hemorrhage

hemorrhage of the knee

hemorrhagenic

hemorrhagic

hemorrhagic abscess

hemorrhagic ascites

hemorrhagic disease of the newborn

(hem'(-)răi )

## hemorrhage

To hear audio pronunciation of this topic, purchase a subscription or log in.

[ hem- + -rrhage ]  
Loss of blood. The term is usually used for episodes of bleeding that last more than a few minutes, compromise organ or tissue perfusion, or threaten life. The most hazardous forms of blood loss result from arterial bleeding, internal bleeding, or bleeding into the cranium. The risk of uncontrolled bleeding is greatest in people who have suffered major trauma, or in those who take anticoagulant drugs or who have other coagulation defects.  
hemorrhagic (hem-ŏ-raj'ik )

To hear audio pronunciation of this topic, purchase a subscription or log in.

, adj. SEE TABLE: [Common Sites of Bleeding](#)

### SYMPTOMS AND SIGNS

Orthostatic dizziness, weakness, fatigue, shortness of breath, and palpitations are common symptoms of hemorrhage. Signs of hemorrhage include tachycardia, hypotension, pallor, and cold, moist skin.

### TREATMENT

Pressure should be applied directly to any obviously bleeding body part, and the part should be elevated. Cautery may be

Search PRIME  
PubMed

### Related Topics

- T fetomaternal hemorrhage
- T splinter hemorrhage
- T nosebleed
- T Terson syndrome
- T ICH
- T myelitis
- T pleurisy
- T weeping
- T antihemorrhagic,...
- T headache

[more...](#)

## Want to read the entire topic?

Access up-to-date medical information for less than \$1 a week

[Purchase a subscription](#)

[I'm already a subscriber](#)

[Browse sample topics](#)

For trauma patients with massive bleeding, the experienced nurse or emergency care provider may apply pneumatic splints or antishock garments during patient transportation to the hospital. These devices may prevent hemorrhagic shock.



**Standard precautions should be used for all procedures involving contact with blood or wounds.**

### Common Sites of Bleeding

Lower GI tract	Hematochezia; melena
Upper GI tract	Hematemesis
Joints	Hemarthrosis
Lungs/bronchi (coughed up)	Hemoptysis
Nasal passages	Epistaxis
Skin	Ecchymosis
Urinary tract	Hematuria



There's more to see -- the rest of this topic is available only to subscribers.



Home

Contact Us

Privacy / Disclaimer

Terms of Service

Log in

© 2000–2025 Unbound Medicine, Inc. All rights reserved

CONNECT WITH US

