

Determination of factors governing fibrillogenesis of polypeptide chains using lattice models

Mai Suan Li¹, Nguyen Truong Co², Govardhan Reddy³, C-K. Hu^{5,6}, and D. Thirumalai^{3,4}

¹*Institute of Physics, Polish Academy of Sciences, Al. Lotnikow 32/46, 02-668 Warsaw, Poland*

²*Saigon Institute for Computational Science and Technology, 6 Quarter,
Linh Trung Ward, Thu Duc District, Ho Chi Minh City, Vietnam*

³*Biophysics Program, Institute for Physical Science and Technology,
University of Maryland, College Park, MD 20742*

⁴*Department of Chemistry and Biochemistry, University of Maryland, College Park, MD 20742*

⁵*Institute of Physics, Academia Sinica, Nankang, Taipei 11529, Taiwan*

⁶*Center for Nonlinear and Complex Systems and Department of Physics,
Chung Yuan Christian University, Chungli 32023, Taiwan*

Using lattice models we explore the factors that determine the tendencies of polypeptide chains to aggregate by exhaustively sampling the sequence and conformational space. The morphologies of the fibril-like structures and the time scales (τ_{fib}) for their formation depend on a subtle balance between hydrophobic and coulomb interactions. The extent of population of \mathbf{N}^* , which is a fibril-prone structure in the spectrum of monomer conformations, is the major determinant of τ_{fib} . This observation is used to determine the aggregation-prone consensus sequences by exhaustively exploring the sequence space. Our results provide a basis for genome wide search of fragments that are aggregation prone.

PACS numbers: 87.15.A,87.14.E