Title

Estimation of the binding free energy of AC1NX476 to HIV-1 protease wild-type and mutations using free energy perturbation method

Authors

Son Tung Ngo^{1,2}, Binh Khanh Mai³, Dinh Minh Hiep⁴, and Mai Suan Li²

Institutions

Short Title: Binding free energy of AC1NX476 to HIV-1 protease

Corresponding Author: Son Tung Ngo (email: nstung@ifpan.edu.pl) and Mai Suan Li (email: masli@ifpan.edu.pl)

Key words: HIV-1 protease, AC1NX476, Darunavir, Ritonavir, Free energy perturbation method, binding free energy, docking, binding cavity.

Abstract

The binding mechanism of AC1NX476 to HIV-1 protease wild-type and mutations was studied by the docking and molecular dynamics simulations. The binding free energy was calculated using the double-annihilation binding-free energy method. It is shown that the binding affinity of AC1NX476 to wild-type is higher than not only ritonavir but also darunavir making AC1NX476 become attractive candidate for HIV treatment. Our theoretical results are in excellent agreement with the experiment data as the correlation coefficient between calculated and experimentally measured binding free energies R=0.993. Residues Asp25-A, Asp29-A, Asp30-A, Ile47-A,

Gly48-A, and Val50-A from chain A, and Asp25-B from chain B play a crucial role in the ligand binding. The mutations were found to reduce the receptor-ligand interaction by widening the binding cavity and the binding propensity is mainly driven by the van der Waals interaction. Our finding may be useful for designing potential drugs to combat with HIV.

¹Institute for Computational Science and Technology, Quang Trung Software City, Tan Chanh Hiep Ward, District 12, Ho Chi Minh City, Vietnam

²Institute of Physics, Polish Academy of Sciences, Al. Lotnikow 32/46, 02-668 Warsaw, Poland ³Department of Applied Chemistry, College of Applied Sciences, Kyung Hee University, Yongin 446-701, Korea

⁴Department of Sciences and Technology, Ho Chi Minh City, Vietnam