

Introduction to Particles and Nuclear Physics

Class Exercise 4

Matan Parnas

Tel Aviv University

20/06/2024

Today's Topics:

1 Feynman Diagrams

Feynman Diagrams

Feynman diagrams are an important tool in particle physics which allow us to visualize and understand the processes in the SM.

Feynman Diagrams

Feynman diagrams are an important tool in particle physics which allow us to visualize and understand the processes in the SM.

To calculate the probabilities for relativistic scattering processes we need to find out the Lorentz-invariant *scattering amplitude*, \mathcal{M}_{fi} , which connects an initial state $|\Psi_i\rangle$ to a final state $|\Psi_f\rangle$.

Feynman Diagrams

Feynman diagrams are an important tool in particle physics which allow us to visualize and understand the processes in the SM.

To calculate the probabilities for relativistic scattering processes we need to find out the Lorentz-invariant *scattering amplitude*, \mathcal{M}_{fi} , which connects an initial state $|\Psi_i\rangle$ to a final state $|\Psi_f\rangle$.

$$A \sim M_{fi} = \langle \Psi_f | \mathcal{O} | \Psi_i \rangle \rightarrow P \propto A^2 \sim |M_{fi}|^2$$

Feynman diagrams are graphs that give pictorial descriptions of a way for the state $|\Psi_i\rangle$ to propagate into state $|\Psi_f\rangle$. The full expression of \mathcal{M}_{fi} is an expansion obtained using perturbation theory. Each diagram represents a contribution to \mathcal{M}_{fi} and can be translated to a complex number.

$$M_{fi} = \sum_{i=0}^{\infty} M_{fi}^{(i)} \quad ; \quad M_{fi}^{(i)} \in \mathbb{C}$$

In this course we will not learn how to properly calculate the probabilities using Feynman diagrams. But even without extracting the quantitative expressions, Feynman diagrams are still of great importance, and they'll help us determine which processes are allowed, and will give us ways to *qualitatively* compare between different processes and their probabilities.

Feynman Diagrams


Let's start with a simple example of $q\bar{q} \rightarrow q\bar{q}$

(Electromagnetic scattering between a quark and an anti-quark)

Feynman Diagrams

Let's start with a simple example of $q\bar{q} \rightarrow q\bar{q}$
(Electromagnetic scattering between a quark and an anti-quark)

This process can be described with the following Feynman diagram:




Feynman Diagrams

Let's start with a simple example of $q\bar{q} \rightarrow q\bar{q}$

(Electromagnetic scattering between a quark and an anti-quark)

This process can be described with the following Feynman diagram:



General Feynman diagrams rules:



fermion



anti - fermion



EM/Weak boson ($\gamma/W, Z$)



gluon

In our notation time goes from left to right

External lines: Left side = initial state particles = $|\Psi_i\rangle$, Right side = final state particles = $|\Psi_f\rangle$

Internal lines are called *propagators*

Feynman Diagrams

External lines:

- Represents the initial state (left) and final state particles (right)
- Each such line can be associated with "well-defined" (measurable) 4-momenta p_i^μ .
- These states are often referred to as "on-shell", "asymptotic" or "free" states and represent measurable particles.

Feynman Diagrams

External lines:

- Represents the initial state (left) and final state particles (right)
- Each such line can be associated with "well-defined" (measurable) 4-momenta p_i^μ .
- These states are often referred to as "on-shell", "asymptotic" or "free" states and represent measurable particles.

Internal lines:

- Also called propagators, or "virtual" particles. They represent "intermediate states".
- We often use their 4-momentum q^μ to perform calculations, but in principle we can not measure it.
- Since these particles exist for very short times before they are destroyed, they can have different mass/energy than the real particles
 - ▶ according to the uncertainty principle $\Delta E \Delta t \geq \frac{\hbar}{2}$


Feynman Diagrams

External lines:

- Represents the initial state (left) and final state particles (right)
- Each such line can be associated with "well-defined" (measurable) 4-momenta p_i^μ .
- These states are often referred to as "on-shell", "asymptotic" or "free" states and represent measureable particles.

Internal lines:


- Also called propagators, or "virtual" particles. They represents "intermediate states".
- We often use their 4-momentum q^μ to perform calculations, but in principle we can not measure it.
- Since these particles exists for very short times before they are destroyed, they can have different mass/energy than the real particles
 - ▶ according to the uncertainty principle $\Delta E \Delta t \geq \frac{\hbar}{2}$
- This can explain how can we create a pair of $q\bar{q}$ from a "massless" photon



Feynman Diagrams

The Vertex:


- This is where particles are created or annihilated.
- Describes the type of interaction, so vertices are interaction-specific.
- It enforces all the conservation laws relevant for that interaction ("what goes in must come out").
 - ▶ ("Quantum numbers that enter the vertex = Quantum numbers that come out of the vertex")



Feynman Diagrams

The Vertex:


- This is where particles are created or annihilated.
- Describes the type of interaction, so vertices are interaction-specific.
- It enforces all the conservation laws relevant for that interaction ("what goes in must come out").
 - ▶ ("Quantum numbers that enter the vertex = Quantum numbers that come out of the vertex")
- Since we are discussing perturbative processes, there is an infinite number of Feynman diagrams for each process (just like orders in perturbation theory).
- The "order" of the diagram is determined by the number (and type) of vertices.
- Each vertex gives a factor of the "*coupling constant*" - dimensionless measure of the strength of the interaction it represents.



Feynman Diagrams


The Vertex:

- This is where particles are created or annihilated.
- Describes the type of interaction, so vertices are interaction-specific.
- It enforces all the conservation laws relevant for that interaction ("what goes in must come out").
 - ▶ ("Quantum numbers that enter the vertex = Quantum numbers that come out of the vertex")
- Since we are discussing perturbative processes, there is an infinite number of Feynman diagrams for each process (just like orders in perturbation theory).
- The "order" of the diagram is determined by the number (and type) of vertices.
- Each vertex gives a factor of the "*coupling constant*" - dimensionless measure of the strength of the interaction it represents.




For example for the diagram we saw before, which has two "EM" vertices (that connect to a photon), we can write:

$$|\mathcal{M}| \propto e^2$$




Question 1: ee-scattering

For the following diagrams, which describe an electromagnetic scattering between two electrons, what is the ratio between the contributions to the amplitude? What is the ratio between their "probabilities"?




First order / Leading order ("tree-level")




Second order / Next-to-Leading order ("one-loop")

Question 1: ee-scattering

For the following diagrams, which describe an electromagnetic scattering between two electrons, what is the ratio between the contributions to the amplitude? What is the ratio between their "probabilities"?



First order / Leading order ("tree-level")



Second order / Next-to-Leading order ("one-loop")


Solution:

The leading order diagram (left) has two EM vertices, so we denote $|\mathcal{M}_\gamma| \propto e^2$. The loop diagram (right) has four EM vertices so we denote $|\mathcal{M}_{\gamma\gamma}| \propto e^4$. Hence the ratio of amplitudes is


$$\frac{|\mathcal{M}_{\gamma\gamma}|}{|\mathcal{M}_\gamma|} \approx e^2 \sim \alpha = \frac{1}{137}$$

Question 1: ee-scattering

For the following diagrams, which describe an electromagnetic scattering between two electrons, what is the ratio between the contributions to the amplitude? What is the ratio between their "probabilities"?



First order / Leading order ("tree-level")



Second order / Next-to-Leading order ("one-loop")

Solution:

The leading order diagram (left) has two EM vertices, so we denote $|\mathcal{M}_\gamma| \propto e^2$. The loop diagram (right) has four EM vertices so we denote $|\mathcal{M}_{\gamma\gamma}| \propto e^4$. Hence the ratio of amplitudes is $\frac{|\mathcal{M}_{\gamma\gamma}|}{|\mathcal{M}_\gamma|} \approx e^2 \sim \alpha = \frac{1}{137}$

The probability of a process is proportional to the square of the amplitude. This means that the second order diagram is suppressed by a factor of $\sim \alpha^2 \approx 10^{-4}$ w.r.t leading order. In principle one should sum all of the diagrams to evaluate the probability of a process. But this result shows it's sometimes enough to consider only the first order(s) in perturbation theory.


Feynman Diagrams

For $2 \rightarrow 2$ processes, there are cases where there is more than one **leading order** diagrams.


Feynman Diagrams

For $2 \rightarrow 2$ processes, there are cases where there is more than one **leading order** diagrams.

For example, for the $q\bar{q} \rightarrow q\bar{q}$ we saw before, we can have:



"s-channel"




"t-channel"


Feynman Diagrams

For $2 \rightarrow 2$ processes, there are cases where there is more than one **leading order** diagrams.

For example, for the $q\bar{q} \rightarrow q\bar{q}$ we saw before, we can have:



"s-channel"



"t-channel"

Quantum Mechanics tells us that the two diagrams are coherent, and in order to evaluate the total amplitude we must sum them.



In general this is true for all orders of perturbation theory (diagrams of all orders), so actually:

$$\mathcal{M}_{fi} = \sum_{l \in \text{Order}}^{\infty} \sum_{m_l=0}^{N_l} M_{fi}^{(l,m)} = \sum_{i \in \text{order } 1} \mathcal{M}_i + \sum_{j \in \text{order } 2} \mathcal{M}_j + \sum_{k \in \text{order } 3} \mathcal{M}_k + \dots$$

Feynman Diagrams

Mandelstam variables

The difference between the two types of processes can be more easily understood in terms of the *Mandelstam variables*.





For particles with 4-momenta p_1^μ, p_2^μ in the initial state, p_3^μ, p_4^μ in the final state, and propagator momenta q^μ , we define:

Feynman Diagrams

Mandelstam variables

The difference between the two types of processes can be more easily understood in terms of the *Mandelstam variables*.





For particles with 4-momenta p_1^μ, p_2^μ in the initial state, p_3^μ, p_4^μ in the final state, and propagator momenta q^μ , we define:

- $s \equiv q^2 = (p_1 + p_2)^2 = (p_3 + p_4)^2$
- $t \equiv q^2 = (p_1 - p_3)^2 = (p_2 - p_4)^2$
- $u \equiv q^2 = (p_1 - p_4)^2 = (p_2 - p_3)^2$

Feynman Diagrams

Mandelstam variables

The difference between the two types of processes can be more easily understood in terms of the *Mandelstam variables*.



For particles with 4-momenta p_1^μ, p_2^μ in the initial state, p_3^μ, p_4^μ in the final state, and propagator momenta q^μ , we define:

- $s \equiv q^2 = (p_1 + p_2)^2 = (p_3 + p_4)^2$
- $t \equiv q^2 = (p_1 - p_3)^2 = (p_2 - p_4)^2$
- $u \equiv q^2 = (p_1 - p_4)^2 = (p_2 - p_3)^2$

Note: For s-channel process the total amount of energy available for creating new particles is higher than in t- or u- channels

Feynman Diagrams

A small remark regarding t- and u- channels:


u-channel diagrams are relevant only when we have in the final state two "indistinguishable" particles.

Feynman Diagrams

A small remark regarding t- and u- channels:

u-channel diagrams are relevant only when we have in the final state two "indistinguishable" particles.

In the previous example, for the t-channel diagram the vertex can only connect a quark with a quark, and anti-quark with an anti-quark.




Feynman Diagrams


A small remark regarding t- and u- channels:

u-channel diagrams are relevant only when we have in the final state two "indistinguishable" particles.


In the previous example, for the t-channel diagram the vertex can only connect a quark with a quark, and anti-quark with an anti-quark.



But if we look for example at $e^- e^- \rightarrow e^- e^-$ we need to consider both options:



t - channel



u - channel

Question 2:

Draw the leading order diagrams contributing to the following processes:



- $e^+e^- \rightarrow e^+e^-$

Question 2:

Draw the leading order diagrams contributing to the following processes:

• $e^+e^- \rightarrow e^+e^-$

In the leading order there are two s-channel diagrams, and two t-channel diagrams (Two diagrams for each channel because the propagator can be either a γ or a Z).





Question 2:

Draw the leading order diagrams contributing to the following processes:

- $e^+e^- \rightarrow e^+e^-$

In the leading order there are two s-channel diagrams, and two t-channel diagrams (Two diagrams for each channel because the propagator can be either a γ or a Z).




- $e^+e^- \rightarrow \mu^+\mu^-$

Question 2:

Draw the leading order diagrams contributing to the following processes:


• $e^+e^- \rightarrow e^+e^-$

In the leading order there are two s-channel diagrams, and two t-channel diagrams (Two diagrams for each channel because the propagator can be either a γ or a Z).



• $e^+e^- \rightarrow \mu^+\mu^-$

Here we only have the two s-channel diagrams (via a γ or a Z). a t-channel diagram like in the previous example is not allowed as it would require a vertex connecting e and μ , and this would violate conservation of Lepton family (generation) number in each of the vertices.




Question 3:

Draw the leading order diagram for the process $K^- \rightarrow \pi^0 e^- \bar{\nu}_e$. Show which Quantum numbers are conserved and which are not

Question 3:

Draw the leading order diagram for the process $K^- \rightarrow \pi^0 e^- \bar{\nu}_e$. Show which Quantum numbers are conserved and which are not

K^- is a meson comprised of \bar{u}, s . π^0 is a meson comprised of \bar{u}, u ¹. This means that for the process to take place a s quark must transform into an u quark, and this can only happen via the charged weak interaction (with a W^-). So the total flavor is not conserved (which is always the case for processes involving a W). In addition, since this decay involves quarks from different generations u, s , this means the Quark family number is also not conserved.



¹Actually π^0 is a state of superposition of $\frac{\bar{u}u - \bar{d}d}{\sqrt{2}}$