

Linear Algebra

Star

Contents

Chapter 1	Vector Spaces	5
1	Definitions	5
2	Bases	8
3	Dimension	9
Chapter 2	Matrices	11
1	Definitions	11
2	Linear Equations	13
Chapter 3	Linear Mappings	15
1	Definitions	15
2	Kernel and Image	16
3	Isomorphism	18
Chapter 4	Linear Maps and Matrices	19
1	The Linear Map associated with a matrix	19
2	The Matrix Representation of a linear map	19
Chapter 5	Scalar Products and Orthogonality	25
1	Scalar Products	25
2	Hermitian Products	29
3	Bilinearity and Quadratic Form	30
4	Determinants	30
5	Symmetric, Hermitian and Unitary Operators	34
6	Eigenvectors and eigenvalues	40
7	Polynomials and Triangulation of Matrices and Operators	46
Appendix A	Chinese-English glossary	49

Chapter 1 Vector Spaces

1 Definitions

Definition 1.1.1 A **field** $\mathcal{K}$ is a set equipped with addition and multiplication, denoted by $(\mathcal{K}, +, \cdot)$, satisfying the following axioms.

1. **Associative Laws:** let $a, b, c \in \mathcal{K}$,

$$a + b + c = a + (b + c), \quad a \cdot b \cdot c = a \cdot (b \cdot c).$$

2. **Existence of Identity Elements:** let $a \in \mathcal{K}$,

$$a + 0_{\mathcal{K}} = a, \quad a \cdot 1_{\mathcal{K}} = a,$$

where $0_{\mathcal{K}}$ and $1_{\mathcal{K}}$ is called the additive identity and multiplicative identity.

3. **Existence of Negatives:** for any $a \in \mathcal{K}$, there exists one element in $\mathcal{K}$ denoted by $(-a)$ such that

$$a + (-a) = 0_{\mathcal{K}}.$$

4. **Existence of Reciprocal:** for any $a \in \mathcal{K}$ and $a \neq 0_{\mathcal{K}}$, there exists one element in $\mathcal{K}$ denoted by a^{-1} such that

$$a \cdot a^{-1} = 1_{\mathcal{K}}.$$

5. **Commutative Laws:** let $a, b \in \mathcal{K}$,

$$a + b = b + a, \quad a \cdot b = b \cdot a.$$

6. **Attributive Law:** let $a, b, c \in \mathcal{K}$,

$$a \cdot (b + c) = a \cdot b + a \cdot c.$$

Remark 1.1.2 Closure is contained in the defined addition and multiplication.

Theorem 1.1.3 $0_{\mathcal{K}}$ and $1_{\mathcal{K}}$ defined above is unique.

Example 1.1.4 Note that $\mathbb{C}, \mathbb{R}, \mathbb{Q}$ are fields, but $\mathbb{N}$ is not a field because $2 \in \mathbb{N}$ but its reciprocal $\frac{1}{2} \notin \mathbb{N}$. This is the main difference between $\mathbb{Q}$ and $\mathbb{N}$.

Definition 1.1.5 Let $\mathcal{F}$ be a field. A subset $\mathcal{K} \subseteq \mathcal{F}$ is called a **subfield** of $\mathcal{F}$ if $\mathcal{K}$ is itself a field under the same operations of addition and multiplication as $\mathcal{F}$.

Equivalently, $\mathcal{K}$ is a subfield of $\mathcal{F}$ if and only if the following conditions hold:

1. **Closure under Addition and Multiplication:** for all $x, y \in \mathcal{K}$,

$$x + y \in \mathcal{K}, \quad x \cdot y \in \mathcal{K}.$$

2. **Existence of Additive and Multiplicative Inverses:** for all $x \in \mathcal{K}$,

$$-x \in \mathcal{K},$$

and if $x \neq 0_{\mathcal{K}}$, then

$$x^{-1} \in \mathcal{K}.$$

3. **Containment of Identity Elements:**

$$0_{\mathcal{K}}, 1_{\mathcal{K}} \in \mathcal{K}.$$

Remark 1.1.6 This provides a simpler way to prove a set is a field. The third rule ensures that $\emptyset$ is not a subfield.

Proposition 1.1.7 Let $\mathcal{K}$ be a field and A, B are subfields of $\mathcal{K}$, then $A \cap B$ is also a subfield of $\mathcal{K}$.

Proposition 1.1.8 Let $\mathcal{K}$ be a field and A, B are subfields of $\mathcal{K}$, then $A \cup B$ is also a subfield of $\mathcal{K}$ if and only if $A \subseteq B$ or $B \subseteq A$.

Remark 1.1.9 For any elements in a field, we call them **numbers** or **scalars**.

Definition 1.1.10 A **vector space** V over a field $\mathcal{K}$ is a set equipped with two operations: vector addition and scalar multiplication, denoted by $(V, +, \cdot)$, satisfying the following axioms:

1. **Associative Laws:** for any $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$,

$$\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}.$$

2. **Existence of Identity Elements:** there exists a zero vector $\mathbf{0} \in V$ such that for any $\mathbf{v} \in V$,

$$\mathbf{v} + \mathbf{0} = \mathbf{v}.$$

3. **Existence of Additive Inverses:** for any $\mathbf{v} \in V$, there exists an element $-\mathbf{v} \in V$ such that

$$\mathbf{v} + (-\mathbf{v}) = \mathbf{0}.$$

4. **Commutative Law:** for any $\mathbf{u}, \mathbf{v} \in V$,

$$\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}.$$

5. **Distributive Laws:** for all $a, b \in \mathcal{K}$ and $\mathbf{u}, \mathbf{v} \in V$,

$$a \cdot (\mathbf{u} + \mathbf{v}) = a \cdot \mathbf{u} + a \cdot \mathbf{v}, \quad (a + b) \cdot \mathbf{v} = a \cdot \mathbf{v} + b \cdot \mathbf{v}.$$

6. **Compatibility of Scalar Multiplication:** for all $a, b \in \mathcal{K}$ and $\mathbf{v} \in V$,

$$(ab) \cdot \mathbf{v} = a \cdot (b \cdot \mathbf{v}).$$

7. **Identity of Scalar Multiplication:** for all $\mathbf{v} \in V$,

$$1_{\mathcal{K}} \cdot \mathbf{v} = \mathbf{v}.$$

Theorem 1.1.11 The zero vector $\mathbf{0}$ is unique.

Definition 1.1.12 Let V be a vector space over a field $\mathcal{K}$. A non-empty subset $W \subseteq V$ is called a **subspace** of V if W is itself a vector space over $\mathcal{K}$ under the same operations of addition and scalar multiplication as V . Equivalently, W is a subspace of V if and only if the following conditions hold for all $\mathbf{u}, \mathbf{v} \in W$ and $a \in \mathcal{K}$:

1. **Closure under Addition:**

$$\mathbf{u} + \mathbf{v} \in W.$$

2. **Closure under Scalar Multiplication:**

$$a \cdot \mathbf{u} \in W.$$

Notation 1.1.13 If W is a subspace of V , we write $W \leq V$.

Remark 1.1.14 We state W to be a non-empty set at first so we needn't to say $\mathbf{0} \in W$.

Remark 1.1.15 For any elements in a vector space, we call them **vectors**.

Example 1.1.16 If $\mathcal{F}$ is a field, then $\mathcal{F}^n$ is always a vector space over $\mathcal{F}$. Furthermore, $\mathcal{F}^n$ is a vector space over $\mathcal{K}$ whenever $\mathcal{K} \subseteq \mathcal{F}$ is a subfield.

Definition 1.1.17 Let V be a vector space over a field $\mathcal{K}$, and let vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$. Any vector of the form

$$\mathbf{v} = a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \dots + a_n \mathbf{v}_n,$$

where $a_1, a_2, \dots, a_n \in \mathcal{K}$, is called a **linear combination** of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$.

Definition 1.1.18 Let V be a vector space over a field $\mathcal{K}$, and let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\} \subseteq V$. The set of all linear combinations of vectors in S is called the **span** of S , denoted by

$$\text{span}(S) = \{a_1 \mathbf{v}_1 + a_2 \mathbf{v}_2 + \dots + a_n \mathbf{v}_n \mid a_1, a_2, \dots, a_n \in \mathcal{K}\}.$$

Theorem 1.1.19 The span of S is the smallest subspace of V containing S . Particularly, $\text{span}(\emptyset) = \{0\} = \text{span}\{0\}$.

Definition 1.1.20 Let V be a vector space over a field $\mathcal{K}$. A subset $S \subseteq V$ is called a **spanning set** of V or a set of **generators** if $\text{span}(S) = V$, that is, every vector in V can be expressed as a linear combination of finitely many vectors from S .

Definition 1.1.21 Let V be a vector space over a field $\mathcal{K}$, and let vectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n \in V$. The set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is said to be **linearly dependent** if there exist scalars $a_1, a_2, \dots, a_n \in \mathcal{K}$, not all zero, such that

$$a_1\mathbf{v}_1 + a_2\mathbf{v}_2 + \dots + a_n\mathbf{v}_n = \mathbf{0}.$$

Otherwise, if this equation holds only when $a_1 = a_2 = \dots = a_n = 0$, the set is said to be **linearly independent**.

Remark 1.1.22 $\{0\}$ is linearly dependent, but $\emptyset$ is linearly independent.

2 Bases

Definition 1.2.1 Let V be a vector space over a field $\mathcal{K}$. A subset $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\} \subseteq V$ is called a **basis** of V if it satisfies the following two conditions:

1. $\mathcal{B}$ is linearly independent.
2. $\mathcal{B}$ spans V , i.e. $\text{span}(\mathcal{B}) = V$.

Theorem 1.2.2 Every vector in V can be written uniquely as a linear combination of the vectors in $\mathcal{B}$.

Remark 1.2.3 Though we use set notation to denote a basis, the order *does* matter. For instance, two bases

$$\mathcal{B}_1 = \{(1, 0), (0, 1)\}, \quad \mathcal{B}_2 = \{(0, 1), (1, 0)\}$$

are considered to be different. You will know why when we discuss about matrix representation of linear maps.

Theorem 1.2.4 Let $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a set of elements of a vector space V over a field $\mathcal{K}$, and let r be a positive integer with $r \leq n$. We say that $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ is a **maximal subset of linearly independent elements** if $\mathbf{v}_1, \dots, \mathbf{v}_r$ are linearly independent, and moreover, for any $\mathbf{v}_i$ with $i > r$, the set $\{\mathbf{v}_1, \dots, \mathbf{v}_r, \mathbf{v}_i\}$ is linearly dependent.

If, in addition, $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a set of generators of V , then the set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$ is a **basis** of V .

3 Dimension

Theorem 1.3.1 Let V be a vector space over a field $\mathcal{K}$. Let $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$ be a basis of V over $\mathcal{K}$. Let $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n$ be elements of V , and assume that $n > m$. Then $\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n$ are linearly dependent.

Theorem 1.3.2 Let V be a vector space over a field $\mathcal{K}$, and suppose that one basis of V has n elements and another basis has m elements. Then $m = n$.

Definition 1.3.3 Let V be a vector space over a field $\mathcal{K}$. If V has a finite basis $\mathcal{B}$ consisting of n vectors, then V is said to be **finite-dimensional**, and the number n is called the **dimension** of V , denoted by $\dim(V) = n$. If no finite basis exists, V is said to be **infinite-dimensional**.

Remark 1.3.4 Since we prove that for a finite-dimensional vector space, the cardinality of every basis is the same, the dimension of a finite-dimensional vector space is well-defined.

Example 1.3.5 Let $\mathcal{K}$ be a field. Define

$$V = \mathcal{K}[x] = \{a_0 + a_1x + a_2x^2 + \dots + a_nx^n \mid n \in \mathbb{N}, a_i \in \mathcal{K}\},$$

the set of all polynomials with coefficients in $\mathcal{K}$.

Then V is a vector space over $\mathcal{K}$ with the usual polynomial addition and scalar multiplication.

The set

$$\{1, x, x^2, x^3, \dots\}$$

is linearly independent, and no finite subset of it spans V . Hence $\mathcal{K}[x]$ is an **infinite-dimensional** vector space over $\mathcal{K}$.

Definition 1.3.6 Let V be a vector space over a field $\mathcal{K}$. A subspace of V with dimension 1 is called a **line** in V . A subspace of V with dimension 2 is called a **plane** in V .

Definition 1.3.7 Let V be a finite-dimensional vector space over a field $\mathcal{K}$, and let $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ be an ordered basis of V . For any vector $\mathbf{v} \in V$, there exist unique scalars $a_1, a_2, \dots, a_n \in \mathcal{K}$ such that

$$\mathbf{v} = a_1\mathbf{e}_1 + a_2\mathbf{e}_2 + \dots + a_n\mathbf{e}_n.$$

The scalars $a_1, a_2, \dots, a_n$ are called the **components** or **coordinates** of $\mathbf{v}$ with respect to the basis $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$. And the n -tuple $A = (a_1, \dots, a_n)$ is called the **coordinate vector** of $\mathbf{v}$.

Definition 1.3.8 Let V be a vector space over a field $\mathcal{K}$, and let $S \subseteq V$. A subset $\mathcal{B} \subseteq S$ is called a **maximal linearly independent subset** of S if

1. $\mathcal{B}$ is linearly independent;
2. for any vector $\mathbf{v} \in S \setminus \mathcal{B}$, the set $\mathcal{B} \cup \{\mathbf{v}\}$ is linearly dependent.

Theorem 1.3.9 Let V be a vector space over a field $\mathcal{K}$, and let the set $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a maximal set of linearly independent elements of V . Then $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ is a basis of V .

Remark 1.3.10 You might think, “Oh! Why does the theorem above look similar to something previous. Is the author mentally unstable?” Actually, they are different because of their starting points.

Theorem 1.3.11 Let V be a vector space over a field $\mathcal{K}$ of dimension n , and let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ be linearly independent elements of V . Then $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ constitute a basis of V .

Proposition 1.3.12 Let V be a vector space over a field $\mathcal{K}$, and let W be a subspace of V . If $\dim W = \dim V$, then $V = W$.

Remark 1.3.13 This proposition gives us a way to prove that two sets (vector space) are the same.

Proposition 1.3.14 Let V be a vector space over a field $\mathcal{K}$ of dimension n . Let r be a positive integer with $r < n$, and let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r$ be linearly independent elements of V . Then there exist elements $\mathbf{v}_{r+1}, \mathbf{v}_{r+2}, \dots, \mathbf{v}_n$ in V such that

$$\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$$

is a basis of V .

Remark 1.3.15 All these theorems and propositions are telling one same truth: linearly independent sets might be too small to span a vector space, while spanning sets might be too large to be neat (linearly independent), and basis is exactly the boundary, i.e., the smallest spanning set or the largest linearly independent set. It's quite similar to supremum.

Theorem 1.3.16 Let V be a vector space over a field $\mathcal{K}$ having a basis consisting of n elements. Let W be a subspace of V that does not consist of the zero vector alone. Then W has a basis, and the dimension of W satisfies

$$\dim W \leq n.$$

Chapter 2 Matrices

These definitions are just for a quick review. Please refer to Artin's Algebra Chapter 1.1-1.3 for more detailed information.

1 Definitions

Definition 2.1.1 Let $\mathcal{K}$ be a field and let m, n be two positive integers. An array of elements in $\mathcal{K}$ of the form

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}$$

is called a **matrix** over $\mathcal{K}$ of size $m \times n$, or an m -by- n matrix.

Definition 2.1.2 For convenience, we may write $A = (a_{ij})$ for $1 \leq i \leq m$ and $1 \leq j \leq n$ to denote a matrix in $\mathcal{K}^{m \times n}$. The element a_{ij} is called the (i, j) -entry of A . Each **column** of A is a vector in $\mathcal{K}^m$, and each **row** of A is a vector in $\mathcal{K}^n$.

Definition 2.1.3 If $m = n$, the matrix A is called a **square matrix**. If $m \neq n$, A is called a **non-square matrix**. The **zero matrix** is the matrix all of whose entries are zero.

Definition 2.1.4 Let $A = (a_{ij})$ and $B = (b_{ij})$ be two matrices in $\mathbf{M}_{m \times n}(\mathcal{K})$, and let $\alpha \in \mathcal{K}$. **Matrix addition** and **scalar multiplication** are defined as:

$$(A + B) = (a_{ij} + b_{ij}), \quad (\alpha A) = (\alpha a_{ij}), \quad \text{for } 1 \leq i \leq m, 1 \leq j \leq n.$$

Remark 2.1.5 Let $\mathbf{M}_{m \times n}(\mathcal{K})$ (or $\text{Mat}_{m \times n}(\mathcal{K})$) denote the set of all $m \times n$ matrices over $\mathcal{K}$. Equipped with the matrix addition and scalar multiplication defined above, $\mathbf{M}_{m \times n}(\mathcal{K})$ forms a **vector space** over $\mathcal{K}$. Its dimension is

$$\dim \mathbf{M}_{m \times n}(\mathcal{K}) = mn.$$

Definition 2.1.6 The **standard basis** of the vector space $\mathbf{M}_{m \times n}(\mathcal{K})$ is the collection of mn matrices $\{e_{ij}\}$,

where each e_{ij} has entry

$$(e_{ij})_{pq} = \begin{cases} 1, & \text{if } p = i \text{ and } q = j, \\ 0, & \text{otherwise.} \end{cases}$$

Each e_{ij} has a single 1 in the (i, j) -position and zeros elsewhere. Every matrix $A = (a_{ij}) \in \mathbf{M}_{m \times n}(\mathcal{K})$ can be expressed uniquely as

$$A = \sum_{i=1}^m \sum_{j=1}^n a_{ij} e_{ij}.$$

Definition 2.1.7 Let $A = (a_{ij}) \in \mathbf{M}_{m \times n}(\mathcal{K})$ and $B = (b_{jk}) \in \mathbf{M}_{n \times p}(\mathcal{K})$. The **product** of A and B , denoted by AB , is defined as the $m \times p$ matrix

$$AB = (c_{ik}) \in \mathbf{M}_{m \times p}(\mathcal{K}),$$

where each entry c_{ik} is given by

$$c_{ik} = \sum_{j=1}^n a_{ij} b_{jk}, \quad \text{for } 1 \leq i \leq m, 1 \leq k \leq p.$$

Theorem 2.1.8 (Associative Law) Let $A \in \mathbf{M}_{m \times n}(\mathcal{K})$, $B \in \mathbf{M}_{n \times p}(\mathcal{K})$, and $C \in \mathbf{M}_{p \times q}(\mathcal{K})$. Then matrix multiplication is associative:

$$(AB)C = A(BC).$$

Theorem 2.1.9 (Distributive Laws) For matrices $A, A_1, A_2 \in \mathbf{M}_{m \times n}(\mathcal{K})$ and $B, B_1, B_2 \in \mathbf{M}_{n \times p}(\mathcal{K})$, matrix multiplication satisfies

$$A(B_1 + B_2) = AB_1 + AB_2, \quad (A_1 + A_2)B = A_1B + A_2B.$$

Definition 2.1.10 For each positive integer n , the **identity matrix** of order n , denoted by I_n , is the $n \times n$ matrix whose entries are defined by

$$(I_n)_{ij} = \begin{cases} 1, & \text{if } i = j, \\ 0, & \text{otherwise.} \end{cases}$$

The matrix I_n satisfies the property that for any $A \in \mathbf{M}_{m \times n}(\mathcal{K})$ and $B \in \mathbf{M}_{n \times p}(\mathcal{K})$,

$$AI_n = A, \quad I_n B = B.$$

Definition 2.1.11 Let $A \in \mathbf{M}_{n \times n}(\mathcal{K})$ be a square matrix. If there exists a matrix $B \in \mathbf{M}_{n \times n}(\mathcal{K})$ such that

$$AB = BA = I_n,$$

then A is called an **invertible matrix** (or **non-singular matrix**), and B is called the **inverse matrix** of A , denoted by A^{-1} .

If no such B exists, A is called a **singular matrix**.

Remark 2.1.12 Such inverse shown above is unique.

Definition 2.1.13 Let $A = (a_{ij}) \in \mathbf{M}_{m \times n}(\mathcal{K})$. The **transpose** of A , denoted by A^T , is the $n \times m$ matrix obtained by interchanging the rows and columns of A . Formally,

$$A^T = (a_{ji}) \in \mathbf{M}_{n \times m}(\mathcal{K}),$$

that is, the (i, j) -entry of A^T equals the (j, i) -entry of A .

If $A^T = A$, then A is called a **symmetric matrix**.

Theorem 2.1.14 (Properties of the Transpose) Let $A, B \in \mathbf{M}_{m \times n}(\mathcal{K})$ and $\alpha \in \mathcal{K}$. Then the transpose operation satisfies the following properties:

1. $(A^T)^T = A$;
2. $(A + B)^T = A^T + B^T$;
3. $(\alpha A)^T = \alpha A^T$;
4. If $A \in \mathbf{M}_{m \times n}(\mathcal{K})$ and $B \in \mathbf{M}_{n \times p}(\mathcal{K})$, then

$$(AB)^T = B^T A^T.$$

2 Linear Equations

Definition 2.2.1 A **linear system** over a field $\mathcal{K}$ is a collection of m linear equations in n variables or unknowns $x_1, x_2, \dots, x_n$ of the form

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1, \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2, \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = b_m, \end{cases}$$

where $a_{ij}, b_i \in \mathcal{K}$ for all i, j .

It can be written compactly in matrix form as

$$A\mathbf{x} = \mathbf{b},$$

where

$$A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix}, \quad \mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}, \quad \mathbf{b} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}.$$

Definition 2.2.2 A linear system $A\mathbf{x} = \mathbf{b}$ is called:

- **homogeneous** if $\mathbf{b} = \mathbf{0}$, i.e. the system is of the form $A\mathbf{x} = \mathbf{0}$;
- **inhomogeneous** (or nonhomogeneous) if $\mathbf{b} \neq \mathbf{0}$.

Definition 2.2.3 A vector $\mathbf{x} \in \mathcal{K}^n$ satisfying the equation $A\mathbf{x} = \mathbf{b}$ is called a **solution** of the system.

In the special case of a homogeneous system $A\mathbf{x} = \mathbf{0}$:

- the vector $\mathbf{0}$ is called the **trivial solution**;
- any nonzero vector $\mathbf{x} \neq \mathbf{0}$ satisfying $A\mathbf{x} = \mathbf{0}$ is called a **nontrivial solution**.

Theorem 2.2.4 Let

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = 0, \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = 0, \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n = 0, \end{cases}$$

be a homogeneous system of m linear equations in n unknowns, with coefficients in a field $\mathcal{K}$. Assume that $n > m$. Then the system has a **nontrivial solution** in $\mathcal{K}$.

Theorem 2.2.5 Assume that $m = n$ in the system above, and that the column vectors $A^1, A^2, \dots, A^n$ of the coefficient matrix A are linearly independent. Then the system has a **unique solution** in $\mathcal{K}$.

Chapter 3 Linear Mappings

1 Definitions

Definition 3.1.1 Let V and W be vector spaces over the same field $\mathcal{K}$. A mapping $T : V \rightarrow W$ is called a **linear mapping** (or **linear transformation**) if for all $\mathbf{u}, \mathbf{v} \in V$ and all scalars $\alpha, \beta \in \mathcal{K}$, we have

$$T(\alpha\mathbf{u} + \beta\mathbf{v}) = \alpha T(\mathbf{u}) + \beta T(\mathbf{v}).$$

Example 3.1.2 Let $A \in \mathbf{M}_{m \times n}(\mathcal{K})$. Define a mapping

$$T_A : \mathcal{K}^n \rightarrow \mathcal{K}^m, \quad T_A(\mathbf{x}) = A\mathbf{x},$$

for all $\mathbf{x} \in \mathcal{K}^n$. Then T_A is a **linear mapping**.

Indeed, for any $\mathbf{x}, \mathbf{y} \in \mathcal{K}^n$ and $\alpha, \beta \in \mathcal{K}$,

$$T_A(\alpha\mathbf{x} + \beta\mathbf{y}) = A(\alpha\mathbf{x} + \beta\mathbf{y}) = \alpha A\mathbf{x} + \beta A\mathbf{y} = \alpha T_A(\mathbf{x}) + \beta T_A(\mathbf{y}).$$

Hence T_A satisfies the linearity condition, and matrix left-multiplication defines a linear mapping from $\mathcal{K}^n$ to $\mathcal{K}^m$.

Example 3.1.3 Let $V = \mathcal{K}^n$ and $W = \mathcal{K}^m$. Denote by $\mathcal{L}(V, W)$ the set of all linear mappings from V to W over the field $\mathcal{K}$. That is,

$$\mathcal{L}(V, W) = \{ T : V \rightarrow W \mid T \text{ is linear} \}.$$

For $T, F \in \mathcal{L}(V, W)$ and $\alpha \in \mathcal{K}$, we define their **sum** and **scalar multiple** by

$$(T + F)(\mathbf{u}) = T(\mathbf{u}) + F(\mathbf{u}), \quad (\alpha T)(\mathbf{u}) = \alpha T(\mathbf{u}), \quad \forall \mathbf{u} \in V.$$

With these operations of addition and scalar multiplication, $\mathcal{L}(V, W)$ forms a vector space over $\mathcal{K}$.

Definition 3.1.4 Let V be a vector space over a field $\mathcal{K}$, and let $\mathbf{a} \in V$. The mapping $T_{\mathbf{a}} : V \rightarrow V$ defined by

$$T_{\mathbf{a}}(\mathbf{x}) = \mathbf{x} + \mathbf{a},$$

for all $\mathbf{x} \in V$, is called a **translation** by the vector $\mathbf{a}$.

A translation shifts every vector in V by the same displacement $\mathbf{a}$. Note that a translation is generally **not** a linear mapping, since

$$T_{\mathbf{a}}(\mathbf{0}) = \mathbf{a} \neq \mathbf{0}.$$

Theorem 3.1.5 Let V and W be vector spaces over a field $\mathcal{K}$. Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a basis of V , and let $\mathbf{w}_1, \dots, \mathbf{w}_n$ be arbitrary elements of W . Then there exists a unique linear mapping

$$T : V \rightarrow W$$

such that

$$T(\mathbf{v}_i) = \mathbf{w}_i, \quad i = 1, \dots, n.$$

Moreover, if $x_1, \dots, x_n \in \mathcal{K}$, then

$$T(x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + \dots + x_n\mathbf{v}_n) = x_1\mathbf{w}_1 + x_2\mathbf{w}_2 + \dots + x_n\mathbf{w}_n.$$

2 Kernel and Image

Definition 3.2.1 Let $T : V \rightarrow W$ be a linear mapping between vector spaces over a field $\mathcal{K}$. The **kernel** of T , denoted by $\ker(T)$, is the subset of V defined by

$$\ker(T) = \{ \mathbf{v} \in V \mid T(\mathbf{v}) = \mathbf{0} \}.$$

It consists of all vectors in V that are mapped to the zero vector in W .

Remark 3.2.2 The kernel $\ker(T)$ is a subspace of V .

Theorem 3.2.3 Let $F : V \rightarrow W$ be a linear map between vector spaces over a field $\mathcal{K}$. The following statements are equivalent:

1. $\ker(F) = \{\mathbf{0}\}$;
2. For all $\mathbf{v}, \mathbf{w} \in V$, if $F(\mathbf{v}) = F(\mathbf{w})$, then $\mathbf{v} = \mathbf{w}$.

In other words, F is injective if and only if its kernel is $\{\mathbf{0}\}$.

Remark 3.2.4 Note that F is a mapping if and only if F^{-1} is injective and surjective.

Theorem 3.2.5 Let $F : V \rightarrow W$ be a linear map whose kernel is $\{\mathbf{0}\}$. If $\mathbf{v}_1, \dots, \mathbf{v}_n$ are linearly independent elements of V , then $F(\mathbf{v}_1), \dots, F(\mathbf{v}_n)$ are linearly independent in W .

Definition 3.2.6 Let $T : V \rightarrow W$ be a linear map between vector spaces over a field $\mathcal{K}$. The **image** of T , denoted by $\text{im}(T)$, is the subset of W defined by

$$\text{im}(T) = \{ T(\mathbf{v}) \mid \mathbf{v} \in V \}.$$

It consists of all vectors in W that can be expressed as the image of some vector in V under T .

Remark 3.2.7 The image $\text{im}(T)$ is a subspace of W .

Theorem 3.2.8 (Rank–Nullity Theorem) Let $T : V \rightarrow W$ be a linear map between finite-dimensional vector spaces over $\mathcal{K}$. Then

$$\dim(\ker T) + \dim(\operatorname{im} T) = \dim V.$$

Here, $\dim(\ker T)$ is called the **nullity** of T , and $\dim(\operatorname{im} T)$ is called the **rank** of T .

Theorem 3.2.9 (Rank–Nullity Theorem for Matrices) Let $A \in \mathbf{M}_{m \times n}(\mathcal{K})$, and let its column vectors be

$$\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_n \in \mathcal{K}^m.$$

We call

$$\operatorname{Col}(A) = \operatorname{span}\{\mathbf{A}_1, \dots, \mathbf{A}_n\}$$

the **column space** of A , and

$$\operatorname{Null}(A) = \{\mathbf{x} \in \mathcal{K}^n \mid A\mathbf{x} = \mathbf{0}\}$$

the **null space** of A .

Then $\operatorname{Col}(A)$ is a subspace of $\mathcal{K}^m$, and $\operatorname{Null}(A)$ is a subspace of $\mathcal{K}^n$. Moreover,

$$\dim(\operatorname{Col}(A)) + \dim(\operatorname{Null}(A)) = n,$$

where n is the number of columns of A .

Theorem 3.2.10 Let $L : V \rightarrow W$ be a linear map between finite-dimensional vector spaces over $\mathcal{K}$. Assume that

$$\dim V = \dim W.$$

If $\ker L = \{\mathbf{0}\}$, or if $\operatorname{im} L = W$, then L is **bijective**.

Definition 3.2.11 If f is a mapping, and its inverse correspondence is also a mapping, then we say f is **invertible**.

Proposition 3.2.12 A mapping f is invertible if and only if it is bijective.

Theorem 3.2.13 Suppose V and W are vector spaces over a field $\mathcal{K}$ and $T : V \rightarrow W$ is a linear map that is invertible. Then its inverse $T^{-1} : W \rightarrow V$ is also a linear map.

Theorem 3.2.14 Let V and W be vector spaces over a field $\mathcal{K}$ with

$$\dim V = \dim W,$$

Suppose a linear map $T : V \rightarrow W$, if $\ker T = \{\mathbf{0}\}$, then T is invertible.

3 Isomorphism

Definition 3.3.1 Suppose V and W are vector spaces over a field $\mathcal{K}$. Any invertible linear map $T : V \rightarrow W$ is called an **isomorphism** between V and W .

Notation 3.3.2 We say that V and W are **isomorphic** if there exists an isomorphism $T : V \rightarrow W$. In this case we denote this fact by

$$V \cong W.$$

Theorem 3.3.3 Let V and W be finite-dimensional vector spaces over a field $\mathcal{K}$. Then $V \cong W$ if and only if $\dim V = \dim W$.

Remark 3.3.4 Suppose V and W are vector spaces over a field $\mathcal{K}$ with

$$\dim V = m, \quad \dim W = n.$$

Then $V \cong \mathcal{K}^m$ and $W \cong \mathcal{K}^n$.

Theorem 3.3.5 Let U, V, W be vector spaces over a field $\mathcal{K}$, with linear maps $F : U \rightarrow V$ and $G : V \rightarrow W$. Then the composition map

$$G \circ F : U \rightarrow W$$

is also a linear map.

Chapter 4 Linear Maps and Matrices

1 The Linear Map associated with a matrix

Theorem 4.1.1 Suppose $\mathcal{L} : \mathcal{K}^n \rightarrow \mathcal{K}^m$ is a linear map. Then there exists a unique matrix $A \in \mathbf{M}_{m \times n}(\mathcal{K})$ such that

$$\mathcal{L}(\mathbf{x}) = A\mathbf{x}, \quad \forall \mathbf{x} \in \mathcal{K}^n.$$

Remark 4.1.2 Suppose $A \in \mathbf{M}_{m \times n}(\mathcal{K})$, we call $\mathcal{L}_A : \mathcal{K}^n \rightarrow \mathcal{K}^m$ defined by

$$\mathcal{L}_A(\mathbf{x}) = A\mathbf{x}, \quad \forall \mathbf{x} \in \mathcal{K}^n$$

the **linear map associated with matrix A**. From the above theorem, we also know for any linear map from $\mathcal{K}^n$ to $\mathcal{K}^m$, it is uniquely associated to a matrix.

Proposition 4.1.3 Suppose $A \in \mathbf{M}_{m \times n}(\mathcal{K})$ and $B \in \mathbf{M}_{n \times p}(\mathcal{K})$. Let $\mathcal{L}_A : \mathcal{K}^n \rightarrow \mathcal{K}^m$ and $\mathcal{L}_B : \mathcal{K}^p \rightarrow \mathcal{K}^n$ be the linear maps associated with matrices A and B respectively. Then the composition map

$$\mathcal{L}_A \circ \mathcal{L}_B : \mathcal{K}^p \rightarrow \mathcal{K}^m$$

is the linear map associated with matrix AB . We often omit the symbol $\circ$ and just write

$$\mathcal{L}_A \mathcal{L}_B = \mathcal{L}_{AB},$$

since matrix multiplication is associative.

Proposition 4.1.4 Suppose $A \in \mathbf{M}_{m \times n}(\mathcal{K})$ is invertible, then

$$(\mathcal{L}_A)^{-1}(\mathbf{x}) = \mathcal{L}_{A^{-1}}(\mathbf{x}), \quad \forall \mathbf{x} \in \mathcal{K}^m.$$

2 The Matrix Representation of a linear map

Definition 4.2.1 Suppose V and W are vector spaces over a field $\mathcal{K}$ with

$$\dim V = m, \quad \dim W = n.$$

Let $\mathcal{B}_V = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$ be a basis of V , and $\mathcal{B}_W = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n\}$ be a basis of W . We consider the

following linear maps:

1. A linear map $T : V \rightarrow W$.
2. The linear map $\phi_V : V \rightarrow \mathcal{K}^m$ defined by

$$\phi_V(c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 + \cdots + c_m \mathbf{v}_m) = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_m \end{pmatrix} \in \mathcal{K}^m, \quad \forall c_1, c_2, \dots, c_m \in \mathcal{K}.$$

Note that ϕ_V is an isomorphism between V and $\mathcal{K}^m$.

3. The linear map $\phi_W : W \rightarrow \mathcal{K}^n$ defined by

$$\phi_W(d_1 \mathbf{w}_1 + d_2 \mathbf{w}_2 + \cdots + d_n \mathbf{w}_n) = \begin{pmatrix} d_1 \\ d_2 \\ \vdots \\ d_n \end{pmatrix} \in \mathcal{K}^n, \quad \forall d_1, d_2, \dots, d_n \in \mathcal{K}.$$

Note that ϕ_W is an isomorphism between W and $\mathcal{K}^n$.

Then the linear map

$$\mathcal{L} = \phi_W \circ T \circ \phi_V^{-1} : \mathcal{K}^m \rightarrow \mathcal{K}^n$$

has a unique associated matrix $A \in \mathbf{M}_{n \times m}(\mathcal{K})$. This matrix A is called the **matrix representation** of linear map T with respect to the bases $\mathcal{B}_V$ and $\mathcal{B}_W$, denoted by

$$A = [T]_{\mathcal{B}_V}^{\mathcal{B}_W} \quad \text{or} \quad A = \mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(T).$$

Remark 4.2.2 This is one of the ugliest notations in linear algebra.

Remark 4.2.3 Here's a graphical illustration for the matrix representation of linear maps:

$$\begin{array}{ccc} V & \xrightarrow{T} & W \\ \phi_V \downarrow & \uparrow \phi_V^{-1} & \uparrow \phi_W^{-1} \\ \mathcal{K}^m & \xrightarrow[\mathcal{L}: \mathcal{L}(x)=Ax]{} & \mathcal{K}^n \end{array}$$

If you're not familiar with matrix representations of linear maps, remember to draw this diagram when you get confused.

Proposition 4.2.4 Let $V = W$, $\dim V = n$ and $T = \text{Id}_V$ be the identity map on V with two bases $\mathcal{B}_1 = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ and $\mathcal{B}_2 = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n\}$. Then

$$\mathcal{M}_{\mathcal{B}_1}^{\mathcal{B}_2}(\text{Id}_V) = \mathbf{I}_n \quad \text{if and only if} \quad \mathcal{B}_1 = \mathcal{B}_2.$$

Example 4.2.5 Let $\mathbf{P}_n(\mathcal{K})$ be the vector space of all polynomials with coefficients in a field $\mathcal{K}$ and of degree at most n . That is,

$$\mathbf{P}_n(\mathcal{K}) = \{a_0 + a_1x + a_2x^2 + \cdots + a_nx^n \mid a_0, a_1, \dots, a_n \in \mathcal{K}\}.$$

Consider a linear map $T : \mathbf{P}_n(\mathcal{K}) \rightarrow \mathbf{P}_{n+1}(\mathcal{K})$ defined by

$$T(p) = \int_0^x p(t) dt \quad \forall p \in \mathbf{P}_n(\mathcal{K}).$$

Then the matrix representation of T with respect to the standard basis $\mathcal{B}_n = \{1, x, x^2, \dots, x^n\}$ of $\mathbf{P}_n(\mathcal{K})$ and the standard basis $\mathcal{B}_{n+1} = \{1, x, x^2, \dots, x^{n+1}\}$ of $\mathbf{P}_{n+1}(\mathcal{K})$ is

$$\mathcal{M}_{\mathcal{B}_{n+1}}^{\mathcal{B}_n}(T) = \begin{pmatrix} 0 & 0 & 0 & \cdots & 0 & 0 \\ 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & \frac{1}{2} & 0 & \cdots & 0 & 0 \\ 0 & 0 & \frac{1}{3} & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \frac{1}{n} & 0 \\ 0 & 0 & 0 & \cdots & 0 & \frac{1}{n+1} \end{pmatrix}.$$

Definition 4.2.6 Suppose V is a vector space over a field $\mathcal{K}$ with $\dim V = n$ and $T : V \rightarrow V$ is a linear map. If for a basis $\mathcal{B}$ of V , $\mathcal{M}_{\mathcal{B}}^{\mathcal{B}}(T)$ is a diagonal matrix, then we say the basis $\mathcal{B}$ **diagonalizes** the linear map T . If there exists such a basis $\mathcal{B}$, then we say the linear map T is **diagonalizable**. We call a matrix $A \in \mathbf{M}_{n \times n}(\mathcal{K})$ is **diagonalizable** if the linear map $\mathcal{L}_A : \mathcal{K}^n \rightarrow \mathcal{K}^n$ is diagonalizable.

Definition 4.2.7 Suppose $A, B \in \mathbf{M}_{n \times n}(\mathcal{K})$. We say A is **similar** to B if there exists an invertible matrix $P \in \mathbf{M}_{n \times n}(\mathcal{K})$ such that

$$A = PBP^{-1} \quad \text{or equivalently} \quad AP = PB.$$

Remark 4.2.8 Note that similarity relationship is a equivalence relationship, i.e., it is reflexive ($A \sim A$), symmetric ($A \sim B \implies B \sim A$), and transitive ($A \sim B \wedge B \sim C \implies A \sim C$). Later we will see similar matrices share lots in common.

Proposition 4.2.9 Note that $A \in \mathbf{M}_{n \times n}(\mathcal{K})$ is diagonalizable if and only if A is similar to a diagonal matrix. We write A in the form

$$AS = SD \text{ where } D = \begin{pmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n \end{pmatrix} \text{ and } S = \begin{pmatrix} | & | & \cdots & | \\ \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \\ | & | & \cdots & | \end{pmatrix}.$$

Hence, we have

$$A\mathbf{v}_i = \lambda_i\mathbf{v}_i, \quad i = 1, 2, \dots, n,$$

which motivates the following definition.

Definition 4.2.10 Suppose $A \in \mathbf{M}_{n \times n}(\mathcal{K})$ and $\lambda \in \mathcal{K}$ and $V \in \mathcal{K}^n$. We say λ is an **eigenvalue** of A with corresponding **eigenvector** $\mathbf{v} \neq \mathbf{0}$ if

$$A\mathbf{v} = \lambda\mathbf{v}.$$

If there exists a basis of $\mathcal{K}^n$ such that every basis vector is an eigenvector of A , we call such a basis an **eigenbasis** of A .

Theorem 4.2.11 Suppose V and W are vector spaces over a field $\mathcal{K}$ with

$$\dim V = m, \quad \dim W = n.$$

And $T : V \rightarrow W$ is a linear map. Let $\mathcal{B}_V = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$ be a basis of V , and $\mathcal{B}_W = \{\mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n\}$ be a basis of W . Let $\phi_V : V \rightarrow \mathcal{K}^m$ and $\phi_W : W \rightarrow \mathcal{K}^n$ be the isomorphisms defined as before. Then

$$\phi_W(T(\mathbf{v})) = \mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(T) \phi_V(\mathbf{v}), \quad \forall \mathbf{v} \in V.$$

Theorem 4.2.12 With the same set-up as above, suppose additionally $F : V \rightarrow W$ is another linear map. Then

$$\alpha T + \beta F : V \rightarrow W \quad \forall \alpha, \beta \in \mathcal{K}$$

is also a linear map and we have

$$\mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(\alpha T + \beta F) = \alpha \mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(T) + \beta \mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(F).$$

Thus, the mapping $\mathcal{L}(V, W) \rightarrow \mathbf{M}_{n \times m}(\mathcal{K})$ where $T \mapsto \mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(T)$ is a linear map.

Theorem 4.2.13 Still with the same set-up as above, additionally this time suppose $F : W \rightarrow U$ is a linear map where U is a vector space over a field $\mathcal{K}$ with $\dim U = l$ and $\mathcal{B}_U = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_l\}$ is a basis of U . We define $\phi_U : U \rightarrow \mathcal{K}^l$ accordingly. Then

$$\mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_U}(F \circ T) = \mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(F) \mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(T).$$

Remark 4.2.14 Here's a diagram illustration for the above theorem:

$$\begin{array}{ccccc} V & \xrightarrow{T} & W & \xrightarrow{F} & U \\ \phi_V \downarrow \uparrow \phi_V^{-1} & & \phi_W \downarrow \uparrow \phi_W^{-1} & & \phi_U \downarrow \uparrow \phi_U^{-1} \\ \mathcal{K}^m & \xrightarrow{\mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(T)} & \mathcal{K}^n & \xrightarrow{\mathcal{M}_{\mathcal{B}_W}^{\mathcal{B}_U}(F)} & \mathcal{K}^l \end{array}$$

Proposition 4.2.15 Suppose V and W are isomorphic vector spaces over a field $\mathcal{K}$ with

$$\dim V = \dim W = n$$

and $T : V \rightarrow W$ is an isomorphism. Let $\mathcal{B}_V$ be a basis of V , and $\mathcal{B}_W$ be a basis of W . Then we have

$$\left(\mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_W}(T)\right)^{-1} = \mathcal{M}_{\mathcal{B}_W}^{\mathcal{B}_V}(T^{-1}).$$

Proposition 4.2.16 Assume $\mathcal{B}_1$ and $\mathcal{B}_2$ are two bases of a vector space V over a field $\mathcal{K}$ with

$$\dim V = n.$$

1. We know

$$\mathcal{M}_{\mathcal{B}_2}^{\mathcal{B}_1}(\text{Id}_V) \mathcal{M}_{\mathcal{B}_1}^{\mathcal{B}_2}(\text{Id}_V) = I_n = \mathcal{M}_{\mathcal{B}_1}^{\mathcal{B}_2}(\text{Id}_V) \mathcal{M}_{\mathcal{B}_2}^{\mathcal{B}_1}(\text{Id}_V).$$

Here these two matrices are called **change-of-basis matrices**.

2. For any linear map $T : V \rightarrow V$, we often simplify the notation $\mathcal{M}_{\mathcal{B}_V}^{\mathcal{B}_V}(T)$ to $\mathcal{M}_{\mathcal{B}_V}(T)$.

3. If $T : V \rightarrow V$ and we have two bases $\mathcal{B}_1$ and $\mathcal{B}_2$ of V , then $\mathcal{M}_{\mathcal{B}_2}(T)$ and $\mathcal{M}_{\mathcal{B}_1}(T)$ are similar matrices. In fact, we have

$$\mathcal{M}_{\mathcal{B}_2}(T) = \mathcal{M}_{\mathcal{B}_1}^{\mathcal{B}_2}(\text{Id}_V) \mathcal{M}_{\mathcal{B}_1}(T) \mathcal{M}_{\mathcal{B}_2}^{\mathcal{B}_1}(\text{Id}_V).$$

Using the following diagram might be more intuitive:

$$\begin{array}{ccc} \mathcal{K}^n & \xrightarrow{\quad \mathcal{M}_{\mathcal{B}_2}(T) \quad} & \mathcal{K}^n \\ \phi_2^{-1} \uparrow & & \downarrow \phi_2 \\ & & \\ V & \xrightarrow{\quad T \quad} & V \\ \phi_1 \uparrow & & \downarrow \phi_1^{-1} \\ \mathcal{K}^n & \xrightarrow{\quad \mathcal{M}_{\mathcal{B}_1}(T) \quad} & \mathcal{K}^n \end{array}$$

Chapter 5 Scalar Products and Orthogonality

1 Scalar Products

Definition 5.1.1 Suppose V is a vector space over a field $\mathcal{K}(= \mathbb{R})$. A **scalar product** on V is a mapping

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathcal{K}$$

such that for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$ and all $\alpha, \beta \in \mathcal{K}$, the following properties hold:

1. (Symmetry) $\langle \mathbf{u}, \mathbf{v} \rangle = \langle \mathbf{v}, \mathbf{u} \rangle$.
2. (Linearity in the first argument) $\langle \alpha \mathbf{u} + \beta \mathbf{v}, \mathbf{w} \rangle = \alpha \langle \mathbf{u}, \mathbf{w} \rangle + \beta \langle \mathbf{v}, \mathbf{w} \rangle$.

Additionally, we say the scalar product is **non-degenerate** if

$$\langle \mathbf{v}, \mathbf{u} \rangle = 0, \quad \forall \mathbf{u} \in V \implies \mathbf{v} = \mathbf{0}.$$

Remark 5.1.2 The linearity in the second argument also holds due to the conjugate symmetry.

Definition 5.1.3 Suppose V is a vector space over a field $\mathcal{K}$ with a scalar product $\langle \cdot, \cdot \rangle$. We say two vectors $\mathbf{u}, \mathbf{v} \in V$ are **orthogonal** or **perpendicular** if

$$\langle \mathbf{u}, \mathbf{v} \rangle = 0.$$

Moreover, suppose $S \subseteq V$ is nonempty. We define the **orthogonal set** of S , as

$$S^\perp = \{ \mathbf{v} \in V \mid \langle \mathbf{v}, \mathbf{s} \rangle = 0, \quad \forall \mathbf{s} \in S \}.$$

And we say $\mathbf{v} \in S^\perp$ is **orthogonal** or **perpendicular** to S and write $\mathbf{v} \perp S$.

Proposition 5.1.4 Suppose V is a vector space over a field $\mathcal{K}$ with a scalar product $\langle \cdot, \cdot \rangle$. If a set $S \leq V$, then $S^\perp$ is a subspace of V .

Example 5.1.5 Suppose $A \in \mathbf{M}_{m \times n}(\mathcal{K})$, then $\mathbf{x}$ is a solution of the homogeneous linear system

$$A\mathbf{x} = \mathbf{0}$$

if and only if $\mathbf{x} \in (\text{Row}(A))^\perp$, where $\text{Row}(A)$ is the row space of matrix A . That is,

$$\text{Row}(A) \perp \text{Null}(A).$$

Definition 5.1.6 Suppose V is a finite-dimensional vector space over a field $\mathcal{K}$ with a scalar product $\langle \cdot, \cdot \rangle$. Then V has a basis and any basis $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ of V is called an **orthogonal basis** if

$$\langle \mathbf{v}_i, \mathbf{v}_j \rangle = 0, \quad \forall i \neq j \in \{1, 2, \dots, n\}.$$

Definition 5.1.7 Suppose V is a vector space over a field $\mathcal{K} (= \mathbb{R})$ with a scalar product $\langle \cdot, \cdot \rangle$. We call this scalar product **positive-definite** if

$$\langle \mathbf{v}, \mathbf{v} \rangle \begin{cases} > 0, \mathbf{v} \neq \mathbf{0}, \\ = 0, \mathbf{v} = \mathbf{0}. \end{cases}$$

If the scalar product is positive-definite, we define the **norm** of a vector $\mathbf{v} \in V$ as

$$\|\mathbf{v}\| = \sqrt{\langle \mathbf{v}, \mathbf{v} \rangle}.$$

Then we can define the **distance** between two vectors $\mathbf{u}, \mathbf{v} \in V$ as

$$\text{dist}(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|.$$

Also, if $\mathbf{v} \in V$ satisfies $\|\mathbf{v}\| = 1$, we say $\mathbf{v}$ is a **unit vector**. Note that for any nonzero vector $\mathbf{v} \in V$, the vector

$$\frac{\mathbf{v}}{\|\mathbf{v}\|}$$

is a unit vector called the **normalization** of $\mathbf{v}$.

Remark 5.1.8 Here we write $\mathcal{K} (= \mathbb{R})$ because we need “ > 0 ” to define positive-definiteness. Actually $\mathcal{K}$ can be any field homomorphic to $\mathbb{R}$.

Example 5.1.9 Let f, g be real-valued functions integrable on the interval I containing $[0, 1]$. We define

$$\langle f, g \rangle = \int_0^1 f(x)g(x) dx, \quad \|f\| = \sqrt{\langle f, f \rangle}.$$

Then $\langle \cdot, \cdot \rangle$ is a positive-definite scalar product on the vector space of all real-valued functions integrable on the interval I .

Proposition 5.1.10 Suppose V is a vector space over a field $\mathcal{K} (= \mathbb{R})$ with a positive-definite scalar product $\langle \cdot, \cdot \rangle$.

1. For $c \in \mathcal{K}$ and $\mathbf{v} \in V$, we have $\|c\mathbf{v}\| = |c| \|\mathbf{v}\|$.
2. For any $\mathbf{u}, \mathbf{v} \in V$,

$$\|\mathbf{u} - \mathbf{v}\| = \|\mathbf{u} + \mathbf{v}\| \quad \text{if and only if} \quad \mathbf{u} \perp \mathbf{v}.$$

3. (Pythagorean theorem) For any $\mathbf{u}, \mathbf{v} \in V$ with $\mathbf{u} \perp \mathbf{v}$,

$$\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2.$$

4. (Parallelogram law) For any $\mathbf{u}, \mathbf{v} \in V$,

$$\|\mathbf{u} + \mathbf{v}\|^2 + \|\mathbf{u} - \mathbf{v}\|^2 = 2(\|\mathbf{u}\|^2 + \|\mathbf{v}\|^2).$$

5. (Cauchy-Schwarz inequality) For any $\mathbf{u}, \mathbf{v} \in V$,

$$|\langle \mathbf{u}, \mathbf{v} \rangle| \leq \|\mathbf{u}\| \|\mathbf{v}\|.$$

6. (Triangle inequality) For any $\mathbf{u}, \mathbf{v} \in V$,

$$\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|.$$

7. Suppose $\mathbf{u}, \mathbf{v} \in V$ and $\mathbf{u} \neq \mathbf{0}$. Then

$$c = \frac{\langle \mathbf{v}, \mathbf{u} \rangle}{\langle \mathbf{u}, \mathbf{u} \rangle}$$

is called the **component** of $\mathbf{v}$ in the direction of $\mathbf{u}$ and we call $c\mathbf{u}$ the **projection** of $\mathbf{v}$ onto $\mathbf{u}$. Note that

$$(\mathbf{v} - c\mathbf{u}) \perp \mathbf{u}.$$

Furthermore,

$$\|\mathbf{v} - c\mathbf{u}\| = \min_{\alpha \in \mathcal{K}} \|\mathbf{v} - \alpha\mathbf{u}\|.$$

8. More generally, suppose $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in V$ are non-zero vectors that are pairwise orthogonal. For any $\mathbf{v} \in V$, we define the **component** of $\mathbf{v}$ along $\mathbf{u}_i$ to be

$$c_i = \frac{\langle \mathbf{v}, \mathbf{u}_i \rangle}{\langle \mathbf{u}_i, \mathbf{u}_i \rangle}, \quad i = 1, 2, \dots, k.$$

We call $c_i\mathbf{u}_i$ the **projection** of $\mathbf{v}$ onto $\mathbf{u}_i$. Note

$$\langle \mathbf{v} - \sum_{i=1}^k c_i\mathbf{u}_i, \mathbf{u}_j \rangle = \langle \mathbf{v}, \mathbf{u}_j \rangle - \sum_{i=1}^k c_i \langle \mathbf{u}_i, \mathbf{u}_j \rangle = 0, \quad j = 1, 2, \dots, k.$$

and we have

$$\left(\mathbf{v} - \sum_{i=1}^k c_i\mathbf{u}_i \right) \perp \mathbf{u}, \quad \forall \mathbf{u} \in \text{span} \{ \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \}.$$

We also call $\sum_{i=1}^k c_i\mathbf{u}_i$ the **projection** of $\mathbf{v}$ onto $\text{span} \{ \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \}$. Furthermore,

$$\left\| \mathbf{v} - \sum_{i=1}^k c_i\mathbf{u}_i \right\| = \min_{\alpha_1, \alpha_2, \dots, \alpha_k \in \mathcal{K}} \left\| \mathbf{v} - \sum_{i=1}^k \alpha_i\mathbf{u}_i \right\|.$$

Definition 5.1.11 Suppose V is a finite-dimensional vector space over a field $\mathcal{K} (= \mathbb{R})$ with a scalar product $\langle \cdot, \cdot \rangle$. A basis $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ of V is called an **orthonormal basis** if

$$\langle \mathbf{v}_i, \mathbf{v}_j \rangle = \begin{cases} 1, & i = j, \\ 0, & i \neq j, \end{cases} \quad \forall i, j \in \{1, 2, \dots, n\}.$$

Theorem 5.1.12 (Bessel's Inequality) Suppose V is a finite-dimensional vector space over a field $\mathcal{K} (= \mathbb{R})$ with a positive-definite scalar product $\langle \cdot, \cdot \rangle$. Let $\{w_1, w_2, \dots, w_k\}$ be an orthogonal set of vectors in V . Then for any $\mathbf{v} \in V$, we have

$$\langle \mathbf{v}, \mathbf{v} \rangle \geq \sum_{i=1}^k \frac{|\langle \mathbf{v}, w_i \rangle|^2}{\langle w_i, w_i \rangle}.$$

with equality if and only if $\mathbf{v} \in \text{span}\{w_1, w_2, \dots, w_k\}$.

Lemma 5.1.13 Suppose V is a finite-dimensional vector space over $\mathcal{K} (= \mathbb{R})$ with a positive-definite scalar product $\langle \cdot, \cdot \rangle$. Let W be a proper subspace of V . If $\{w_1, w_2, \dots, w_k\}$ is an orthonormal basis of W , then there exist vectors $w_{k+1}, w_{k+2}, \dots, w_n \in V$ such that $\{w_1, w_2, \dots, w_k, w_{k+1}, w_{k+2}, \dots, w_n\}$ is an orthonormal basis of V .

Proof. This is a constructive proof via the Gram-Schmidt process.

Note that we do know there exists $x_{k+1}, x_{k+2}, \dots, x_n \in V$ such that $\{w_1, w_2, \dots, w_k, x_{k+1}, x_{k+2}, \dots, x_n\}$ is a basis of V . Next we will convert this basis to an orthonormal basis via projection.

Step 1: Project x_{k+1} onto W and denote the projection as

$$w_{k+1} = x_{k+1} - \sum_{i=1}^k \frac{\langle x_{k+1}, w_i \rangle}{\langle w_i, w_i \rangle} w_i.$$

Note that since

$$\langle w_{k+1}, w_j \rangle = \langle x_{k+1}, w_j \rangle - \sum_{i=1}^k \frac{\langle x_{k+1}, w_i \rangle}{\langle w_i, w_i \rangle} \langle w_i, w_j \rangle = 0, \quad j = 1, 2, \dots, k,$$

we have $w_{k+1} \perp W$. Also, $w_{k+1} \neq \mathbf{0}$ since otherwise $x_{k+1} \in W$ which contradicts the fact that $\{w_1, w_2, \dots, w_k, x_{k+1}, x_{k+2}, \dots, x_n\}$ is a basis of V . Then we normalize w_{k+1} .

Inductive Step: We proceed by induction to construct the following vectors $w_{k+2}, w_{k+3}, \dots, w_n$. □

Theorem 5.1.14 Suppose V is a finite-dimensional nontrivial vector space over a field $\mathcal{K} (= \mathbb{R})$ with a positive-definite scalar product $\langle \cdot, \cdot \rangle$. Then V has an orthonormal basis.

Theorem 5.1.15 Suppose V is a vector space over a field $\mathcal{K} (= \mathbb{R})$ with a positive-definite scalar product $\langle \cdot, \cdot \rangle$ and $\dim V = n$. Define W be a nontrivial subspace of V with $\dim W = r$ ($1 \leq r \leq n - 1$). Define the **orthogonal complement** of W in V , which is a subspace of V , as

$$W^\perp = \{\mathbf{v} \in V \mid \langle \mathbf{v}, \mathbf{w} \rangle = 0, \quad \forall \mathbf{w} \in W\}.$$

Then we have

$$\dim W + \dim W^\perp = \dim V, \quad W \oplus W^\perp = V.$$

2 Hermitian Products

For vector space V over $\mathbb{C}$, if we want to have a “scalar product” that is “positive-definite” so that

$$\langle \mathbf{v}, \mathbf{v} \rangle \in \mathbb{R} \text{ and } \langle \mathbf{v}, \mathbf{v} \rangle \geq 0,$$

we need to modify our definition of “scalar product”.

Definition 5.2.1 Suppose V is a vector space over $\mathbb{C}$. A **Hermitian scalar product** on V is a mapping

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C}$$

satisfying the following properties:

1. (Conjugate symmetry) $\langle \mathbf{u}, \mathbf{v} \rangle = \overline{\langle \mathbf{v}, \mathbf{u} \rangle}$.
2. (Linearity in the first argument) $\langle \alpha \mathbf{u} + \beta \mathbf{v}, \mathbf{w} \rangle = \alpha \langle \mathbf{u}, \mathbf{w} \rangle + \beta \langle \mathbf{v}, \mathbf{w} \rangle$.

Additionally, we say the Hermitian scalar product is **positive-definite** if

$$\langle \mathbf{v}, \mathbf{v} \rangle \begin{cases} > 0, \mathbf{v} \neq \mathbf{0}, \\ = 0, \mathbf{v} = \mathbf{0}. \end{cases}$$

Note that

$$\langle \mathbf{v}, \mathbf{v} \rangle = \overline{\langle \mathbf{v}, \mathbf{v} \rangle} \implies \langle \mathbf{v}, \mathbf{v} \rangle \in \mathbb{R}.$$

the positive-definiteness is well-defined.

Remark 5.2.2 (Conjugate Linearity) Note that the linearity in the second argument does not hold.

Remark 5.2.3 For vector space V over $\mathbb{C}$, given a Hermitian product, we say x, y are orthogonal if $\langle x, y \rangle = 0$ ($= \langle y, x \rangle$). Concepts like orthogonal complement can be defined accordingly. If additionally the Hermitian product is positive-definite, concepts like norm, distance, orthogonal basis, orthonormal basis, etc. can be defined accordingly.

Definition 5.2.4 Given $A \in \mathbf{M}_{m \times n}(\mathbb{R})$, we call $\text{Im}(A)$ the **column space** of A and $\text{Im}(A^T)$ the **row space** of A . We also call $\dim \text{Im}(A)$ the **column rank** of A and $\dim \text{Im}(A^T)$ the **row rank** of A . Since the column rank and the row rank are equal, we simply call it the **rank** of A and denote it as $\text{rank}(A)$.

Remark 5.2.5 Let $A \in \mathbf{M}_{m \times n}(\mathbb{R})$ and $R = \text{rref}(A)$. Then we know $\ker(A) = \ker(R)$ and thus $\text{rank}(A) = \text{rank}(R) = \text{pivots of } R$.

3 Bilinearity and Quadratic Form

Definition 5.3.1 Suppose U, V, W are vector spaces over a field $\mathcal{K}$ and $g : U \times V \rightarrow W$ be a map. We say g is **bilinear** if for each fixed $u \in U$, the map $v \mapsto g(u, v)$ is linear from V to W , and for each fixed $v \in V$, the map $u \mapsto g(u, v)$ is linear from U to W .

Example 5.3.2 Let $A \in \mathbf{M}_{m \times n}(\mathcal{K})$. Define $g_A : \mathcal{K}^m \times \mathcal{K}^n \rightarrow \mathcal{K}$ as $g_A(x, y) = x^T A y$. Then g_A is a bilinear map.

Theorem 5.3.3 Given any bilinear map $g : \mathcal{K}^m \times \mathcal{K}^n \rightarrow \mathcal{K}$, there exists a unique matrix $A \in \mathbf{M}_{m \times n}(\mathcal{K})$ such that $g(x, y) = x^T A y (= g_A)$ for all $x \in \mathcal{K}^m$ and $y \in \mathcal{K}^n$.

Theorem 5.3.4 The set of all bilinear maps from $\mathcal{K}^m \times \mathcal{K}^n$ to $\mathcal{K}$, denoted by $\text{Bil}(\mathcal{K}^m \times \mathcal{K}^n, \mathcal{K})$, is a vector space over $\mathcal{K}$. The association $A \mapsto g_A$ is a vector space isomorphism from $\mathbf{M}_{m \times n}(\mathcal{K})$ to $\text{Bil}(\mathcal{K}^m \times \mathcal{K}^n, \mathcal{K})$. That is, we have

$$\dim \text{Bil}(\mathcal{K}^m \times \mathcal{K}^n, \mathcal{K}) = \dim \mathbf{M}_{m \times n}(\mathcal{K}) = mn.$$

Remark 5.3.5 Note that a scalar product define on a vector space V is a bilinear map from $V \times V$ to $\mathcal{K}$. Particularly, if $V = \mathcal{K}^n$, then the unique matrix $A \in \mathbf{M}_{n \times n}(\mathcal{K})$ corresponding to the scalar product is symmetric, i.e., $A^T = A$.

Remark 5.3.6 In some textbook, a scalar product is also called a symmetric bilinear form.

Definition 5.3.7 Suppose V is a vector space over a field $\mathcal{K}$ and $\langle \cdot, \cdot \rangle$ is a scalar product on V . We call the map $f : V \rightarrow \mathcal{K}$ defined by $f(\mathbf{v}) = \langle \mathbf{v}, \mathbf{v} \rangle$ a **quadratic form** on V .

Example 5.3.8 For the case $V = \mathcal{K}^n$, a quadratic form on V can be uniquely associated with a symmetric matrix $A \in \mathbf{M}_{n \times n}(\mathcal{K})$ such that

$$f(x) = x^T A x = \sum_{i=1}^n \sum_{j=1}^n x_i a_{ij} x_j \quad \forall x \in \mathcal{K}^n.$$

Particularly, if A is a diagonal matrix, then we have

$$f(x) = \sum_{i=1}^n a_{ii} x_i^2.$$

4 Determinants

Recall in last semester we defined the determinant of a 2×2 matrix $A = \begin{bmatrix} ab \\ cd \end{bmatrix}$ as $\det(A) = \begin{vmatrix} ab \\ cd \end{vmatrix} = ad - bc$. And we conclude that A is invertible if and only if $\det(A) \neq 0$. Furthermore, we know $A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} d - b \\ -ca \end{bmatrix}$ when A is invertible.

We recall more properties of determinants of 2×2 matrices:

1. $\det(I_2) = 1$.
2. Bilinearity: For any $a, b, c, d \in \mathcal{K}$, we have

$$\begin{vmatrix} a+bc & \\ & de \end{vmatrix} = \begin{vmatrix} ac & \\ & de \end{vmatrix} + \begin{vmatrix} bc & \\ & de \end{vmatrix}, \quad \begin{vmatrix} ab+c & \\ & de \end{vmatrix} = \begin{vmatrix} ab & \\ & de \end{vmatrix} + \begin{vmatrix} c & \\ & de \end{vmatrix}.$$

3. Alternating property: For any $a, b, c, d \in \mathcal{K}$, we have $\begin{vmatrix} ab & \\ & cd \end{vmatrix} = -\begin{vmatrix} cd & \\ & ab \end{vmatrix}$.
4. $\det(A) = \det(A^T)$.

Now we try to expand the definition of determinant to more general cases.

Definition 5.4.1 Based on the definition of 2×2 determinant, we define 3×3 **determinant** as follows: suppose

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \in \mathbf{M}_{3 \times 3}(\mathcal{K}),$$

then we define

$$\det(A) = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}.$$

Remark 5.4.2 We call this formula the **Laplace expansion** or the **cofactor formula** by the first row.

Theorem 5.4.3 The 3×3 determinant satisfies the following properties:

1. $\det(I_3) = 1$.
2. Trilinearity: as a function of the columns in $A \in \mathbf{M}_{m \times n \times 3}(\mathcal{K})$, the determinant is linear in each column when the other two columns are fixed. For example, $\det(A_1, \alpha x + \beta y, A_3) = \alpha \det(A_1, x, A_3) + \beta \det(A_1, y, A_3)$ for any $A_1, A_3 \in \mathcal{K}^3$, $x, y \in \mathcal{K}^3$ and $\alpha, \beta \in \mathcal{K}$.
3. If two adjacent columns of A are equal, then $\det(A) = 0$.

Definition 5.4.4 Suppose function $D : \mathbf{M}_{n \times n}(\mathcal{K}) \rightarrow \mathcal{K}$ or alternatively $D : \mathcal{K}^n \times \mathcal{K}^n \times \cdots \times \mathcal{K}^n \rightarrow \mathcal{K}$ (with n copies of $\mathcal{K}^n$) with notation $D(A)$ or $D(A_1, A_2, \dots, A_n)$ where A_i is the i -th column of A . We say:

1. D is **multilinear**: as a function of the columns in $A \in \mathbf{M}_{m \times n \times n}(\mathcal{K})$, D is linear in each column when the other $n - 1$ columns are fixed.
2. D is **alternating**: if two adjacent columns of A are equal, then $D(A) = 0$.

Theorem 5.4.5 (Determinant Function) There exists a unique multilinear and alternating function $D : \mathbf{M}_{n \times n}(\mathcal{K}) \rightarrow \mathcal{K}$ such that $D(I_n) = 1$. We call this unique D the $n \times n$ **determinant** function and denote it as $\det(\cdot)$.

Proof. We prove in a constructive and inductive way.

Existence:

Initial Cases: For $n = 1$, we define $\det([a]) = a$ for any $a \in \mathcal{K}$. For $n = 2$ and $n = 3$, we have already defined the determinant.

Inductive assumption: suppose such D exists as a map from $\mathbf{M}_{k \times k}(\mathcal{K})$ to $\mathcal{K}$ such that the three properties hold for any $k \leq n - 1$.

Inductive step: For

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1j} & \cdots & a_{1n} \\ \vdots & & \vdots & & \vdots \\ a_{i1} & \cdots & a_{ij} & \cdots & a_{in} \\ \vdots & & \vdots & & \vdots \\ a_{n1} & \cdots & a_{nj} & \cdots & a_{nn} \end{bmatrix} \in \mathbf{M}_{n \times n}(\mathcal{K}),$$

we define

$$D(A) = \sum_{j=1}^n (-1)^{i+j} a_{ij} D(\tilde{A}_{ij}),$$

where $\tilde{A}_{ij}$ is the $(n-1) \times (n-1)$ matrix obtained by deleting the i -th row and the j -th column of A . By the inductive assumption, $D(\tilde{A}_{ij})$ is well-defined and satisfies the three properties. We will show that such D satisfies the three properties as well.

For multilinearity, we consider the individual term $(-1)^{i+j} a_{ij} D(\tilde{A}_{ij})$ in the sum and a particular $k \in \{1, 2, \dots, n\}$.

- If $j \neq k$, then $(-1)^{i+j} a_{ij}$ is independent of A_k and $D(\tilde{A}_{ij})$ is linear in A_k by the inductive assumption. Thus $(-1)^{i+j} a_{ij} D(\tilde{A}_{ij})$ is linear in A_k .
- If $j = k$, then $D(\tilde{A}_{ij})$ is independent of A_k , but $(-1)^{i+j} a_{ij}$ is linear in A_k . Thus $(-1)^{i+j} a_{ij} D(\tilde{A}_{ij})$ is linear in A_k .

Hence $D(A)$ is linear in A_k .

For alternating property, suppose $A_k = A_{k+1}$ for some $k \in \{1, 2, \dots, n-1\}$. Consider:

- If $j \neq k$ and $j \neq k+1$, then $D(\tilde{A}_{ij})$ has two adjacent columns equal and thus $D(\tilde{A}_{ij}) = 0$ by the inductive assumption. Thus $(-1)^{i+j} a_{ij} D(\tilde{A}_{ij}) = 0$.
- In the other cases, we only have

$$D(A) = (-1)^{i+k} a_{ik} D(\tilde{A}_{ik}) + (-1)^{i+k+1} a_{i,k+1} D(\tilde{A}_{i,k+1}) = 0.$$

For identity, we have

$$D(I_n) = (-1)^{i+i} D(I_{n-1}) = 1.$$

Uniqueness: This part is finished in Homework. The core idea is to utilize the permutation of columns and the alternating property to show that D is determined by its value on I_n . $\square$

Theorem 5.4.6 Here are some additional properties of the determinant function:

1. If the i -th and j -th ($i \neq j$) columns of A are swapped, then $\det(A)$ changes its sign.
2. If A has two identical columns or rows, then $\det(A) = 0$.
3. If one adds a scalar multiple of one column to another column, then $\det(A)$ does not change.
4. $\det(A) = \det(A^T)$.

5. $\det(AB) = \det(A) \det(B)$ for any $A, B \in \mathbf{M}_{n \times n}(\mathcal{K})$.

Theorem 5.4.7 (Cramer's Rule) Suppose $A \in \mathbf{M}_{m \times n n \times n}(\mathcal{K})$ satisfies $\det(A) \neq 0$ and $b \in \mathcal{K}^n$. If $x \in \mathcal{K}^n$ is a solution to the linear system $Ax = b$, then x satisfies

$$x_i = \frac{\det(A_{ib})}{\det(A)}, \quad i = 1, 2, \dots, n,$$

where A_{ib} is the matrix obtained by replacing the i -th column of A with b .

Proof. Note $b = x_1 A_1 + x_2 A_2 + \dots + x_n A_n$, where A_i is the i -th column of A . Then we have

$$\det(A_{ib}) = \det\left(A_1, \dots, A_{i-1}, \sum_{j=1}^n x_j A_j, A_{i+1}, \dots, A_n\right) = x_i \det(A).$$

□

Theorem 5.4.8 Suppose $A \in \mathbf{M}_{n \times n}(\mathcal{K})$. If $\det(A) \neq 0$, then the columns of A are linearly independent and thus A is invertible.

Proof. It's equivalent to show that if the columns of A are linearly dependent, then $\det(A) = 0$. Suppose there exists $k \in \{1, 2, \dots, n\}$ such that

$$A_k = \sum_{j=1, j \neq k}^n \alpha_j A_j,$$

where $\alpha_j \in \mathcal{K}$ for $j = 1, 2, \dots, n$ and α_j are not all zero. Then we have

$$\det(A) = \det\left(A_1, \dots, A_{k-1}, \sum_{j=1, j \neq k}^n \alpha_j A_j, A_{k+1}, \dots, A_n\right) = 0.$$

□

Definition 5.4.9 Consider the following two types of column operations:

1. Add a scalar multiple of one column to another column.
2. Interchange two columns.

We call two matrices $A, B \in \mathbf{M}_{m \times n n \times n}(\mathcal{K})$ are **column equivalent** if B can be obtained from A by a succession of column operations of the above two types.

Theorem 5.4.10 Let $A, B \in \mathbf{M}_{m \times n n \times n}(\mathcal{K})$ be column equivalent. Then:

1. $\text{rank}(A) = \text{rank}(B)$.
2. A is invertible if and only if B is invertible.
3. $\det A = 0$ if and only if $\det B = 0$.

Proof. For (1), we only need to show that the column operations of the above two types do not change the rank of a matrix. This is straightforward since these two types of column operations do not change the linear dependence or independence of columns.

For (2), since A is invertible, we have $\text{rank}(A) = n$. By (1), we have $\text{rank}(B) = n$ and thus B is invertible.

For (3), since the column operations only changes the sign of the determinant, we have $\det A = 0$ if and only if $\det B = 0$. $\square$

Theorem 5.4.11 Suppose $A \in \mathbf{M}_{m \times n \times n}(\mathcal{K})$ is column equivalent to a lower-triangular matrix L .

Theorem 5.4.12 Suppose $A \in \mathbf{M}_{m \times n \times n}(\mathcal{K})$. The following conditions are equivalent:

1. A is invertible.
2. The columns of A are linearly independent.
3. $\det A \neq 0$.

Remark 5.4.13 To calculate $\det A$, we can:

1. Laplace expansion by a row or a column.
2. Permutation definition.
3. Convert A to a lower-triangular or upper-triangular matrix L by column operations of the above two types and then calculate $\det L$ as the product of diagonal entries of L .

5 Symmetric, Hermitian and Unitary Operators

Definition 5.5.1 Suppose V is a vector space over $\mathcal{K}$. A linear map $T : V \rightarrow V$ is called a **linear operator** on V .

Definition 5.5.2 Let V be a finite dimensional vector space over field $\mathcal{K}$. The **dual space** of V is denoted by V^* and is defined as

$$V^* = \mathcal{L}(V, \mathcal{K}),$$

i.e., the vector space of all linear maps from V to $\mathcal{K}$. An element of V^* is called a **linear functional** on V .

Example 5.5.3

1. Let V be a finite dimensional vector space over field $\mathcal{K}$ with a scalar product. Suppose $v_0 \in V$. Define

$$f_{v_0} : V \rightarrow \mathcal{K}, \quad f_{v_0}(v) = \langle v, v_0 \rangle \quad \text{for any } v \in V.$$

Then f_{v_0} is a linear functional on V .

2. Let V be a finite dimensional vector space over $\mathbb{C}$ with a hermitian product. Suppose $v_0 \in V$. Define

$$f_{v_0} : V \rightarrow \mathbb{C}, \quad f_{v_0}(v) = \langle v, v_0 \rangle \quad \text{for any } v \in V.$$

Then f_{v_0} is a linear functional on V .

However, the map

$$g_{v_0} : V \rightarrow \mathbb{C}, \quad g_{v_0}(v) = \langle v_0, v \rangle \quad \text{for any } v \in V$$

is, in general, not a linear functional on V by the conjugate linearity in the second argument of the hermitian product.

Lemma 5.5.4 Let V be a finite dimensional vector space over field $\mathcal{K}$.

1. $\dim(V) = n$ if and only if $\dim(V^*) = n$. Thus V , V^* , and $\mathcal{K}^n$ are all isomorphic to each other as vector spaces over $\mathcal{K}$.
2. Suppose $\{v_1, \dots, v_n\}$ is a basis of V . Then there exists a unique basis $\{f_1, \dots, f_n\}$ of V^* such that

$$f_i(v_j) = \delta_{ij} \quad \text{for any } i, j \in \{1, \dots, n\},$$

where δ_{ij} is the Kronecker delta function.

3. Suppose in addition that V is equipped with a non-degenerate scalar product. Then the map

$$\Phi : V \rightarrow V^*, \quad \Phi(v) = f_v,$$

where

$$f_v(w) = \langle w, v \rangle \quad \text{for any } w \in V,$$

is an isomorphism of vector spaces over $\mathcal{K}$.

Proof.

1. Assume $\dim(V) = n$, and let $\{v_1, \dots, v_n\}$ be a basis of V . For each $i \in \{1, \dots, n\}$, define

$$f_i \left(\sum_{j=1}^n a_j v_j \right) = a_i.$$

This is well-defined because every vector in V has a unique expression in the basis $\{v_1, \dots, v_n\}$, and each f_i is linear.

We claim that $\{f_1, \dots, f_n\}$ is a basis of V^* . First, it is linearly independent: if

$$c_1 f_1 + \dots + c_n f_n = 0,$$

then evaluating at v_j gives

$$0 = (c_1 f_1 + \dots + c_n f_n)(v_j) = c_j,$$

so all $c_j = 0$.

Next, it spans V^* . Let $f \in V^*$. Set

$$a_i = f(v_i) \quad (i = 1, \dots, n).$$

Then for any $v = \sum_{j=1}^n x_j v_j$,

$$f(v) = f\left(\sum_{j=1}^n x_j v_j\right) = \sum_{j=1}^n x_j f(v_j) = \sum_{j=1}^n x_j a_j = \sum_{j=1}^n a_j f_j(v).$$

Hence

$$f = \sum_{j=1}^n a_j f_j.$$

Therefore $\{f_1, \dots, f_n\}$ is a basis of V^* , so $\dim(V^*) = n$.

Conversely, if $\dim(V^*) = n$, then applying the same result to the vector space V^* gives

$$\dim((V^*)^*) = n.$$

Since V is finite dimensional, the canonical map from V to $(V^*)^*$ is an isomorphism, hence

$$\dim(V) = \dim((V^*)^*) = n.$$

So $\dim(V) = n$ if and only if $\dim(V^*) = n$.

Since any n -dimensional vector space over $\mathcal{K}$ is isomorphic to $\mathcal{K}^n$, it follows that $V \cong V^* \cong \mathcal{K}^n$.

2. Existence has already been established in part (1): the functionals f_i defined by

$$f_i\left(\sum_{j=1}^n a_j v_j\right) = a_i$$

satisfy

$$f_i(v_j) = \delta_{ij}.$$

We now show uniqueness. Suppose $\{g_1, \dots, g_n\}$ is another basis of V^* such that

$$g_i(v_j) = \delta_{ij} \quad \text{for all } i, j.$$

Let $v = \sum_{j=1}^n a_j v_j$. Then

$$g_i(v) = g_i\left(\sum_{j=1}^n a_j v_j\right) = \sum_{j=1}^n a_j g_i(v_j) = \sum_{j=1}^n a_j \delta_{ij} = a_i = f_i(v).$$

Since this holds for every $v \in V$, we have $g_i = f_i$ for all i . Hence the basis is unique.

3. First, Φ is linear. For any $u, v, w \in V$ and $a, b \in \mathcal{K}$,

$$\Phi(au + bv) = f_{au+bv},$$

and for every $w \in V$,

$$f_{au+bv}(w) = \langle w, au + bv \rangle = a\langle w, u \rangle + b\langle w, v \rangle = af_u(w) + bf_v(w).$$

Hence

$$\Phi(au + bv) = a\Phi(u) + b\Phi(v).$$

Next, Φ is injective. If $\Phi(v) = 0$, then $f_v = 0$, i.e.

$$\langle w, v \rangle = 0 \quad \text{for all } w \in V.$$

By non-degeneracy of the scalar product, this implies $v = 0$.

Finally, since $\dim(V) = \dim(V^*)$ by part (1), a linear injective map

$$\Phi : V \rightarrow V^*$$

between finite dimensional vector spaces of the same dimension must be surjective. Therefore Φ is an isomorphism. □

Theorem 5.5.5 Suppose V is a finite-dimensional vector space over $\mathcal{K}$ with a non-degenerate scalar product $\langle \cdot, \cdot \rangle$ and $T : V \rightarrow V$ is a linear operator. Then there exists a unique linear operator $T^T : V \rightarrow V$ such that

$$\langle T(\mathbf{v}), \mathbf{w} \rangle = \langle \mathbf{v}, T^T(\mathbf{w}) \rangle, \quad \forall \mathbf{v}, \mathbf{w} \in V.$$

Proof. Let $L : V \rightarrow \mathcal{K}$ be $L(\mathbf{v}) = \langle T(\mathbf{v}), \mathbf{w} \rangle$ for any $\mathbf{v} \in V$. Note L is a linear map from V to $\mathcal{K}$ and thus a linear functional. Note by the above lemma, we know there exists a unique $\tilde{w} \in V$ such that $L(x) = f_{\tilde{w}}(x) = \langle x, \tilde{w} \rangle$. Let $T^T(\mathbf{w}) = \tilde{w}$.

Suppose $L_1(\mathbf{v}) = \langle T(\mathbf{v}), \mathbf{w}_1 \rangle$ and $\tilde{w}_1 \in V$ is the unique vector in V such that $L_1 = f_{\tilde{w}_1}$. Suppose $L_2(\mathbf{v}) = \langle T(\mathbf{v}), \mathbf{w}_2 \rangle$ and $\tilde{w}_2 \in V$ is the unique vector in V such that $L_2 = f_{\tilde{w}_2}$.

Note $L' := \alpha L_1 + \beta L_2$ satisfies

$$L'(\mathbf{v}) = \alpha \langle T(\mathbf{v}), \mathbf{w}_1 \rangle + \beta \langle T(\mathbf{v}), \mathbf{w}_2 \rangle = \langle T(\mathbf{v}), \alpha \mathbf{w}_1 + \beta \mathbf{w}_2 \rangle.$$

Thus $T^T(\alpha \mathbf{w}_1 + \beta \mathbf{w}_2) = \alpha T^T(\mathbf{w}_1) + \beta T^T(\mathbf{w}_2)$ and thus T^T is linear. We still need to verify $\langle T(\mathbf{v}), \mathbf{w} \rangle = \langle \mathbf{v}, T^T(\mathbf{w}) \rangle$ for all $\mathbf{v}, \mathbf{w} \in V$. This is straightforward since $L(\mathbf{v}) = \langle T(\mathbf{v}), \mathbf{w} \rangle = f_{\tilde{w}}(\mathbf{v}) = \langle \tilde{w}, \mathbf{v} \rangle = \langle T^T(\mathbf{w}), \mathbf{v} \rangle$. □

Remark 5.5.6 We often just write $\langle T(\mathbf{v}), \mathbf{w} \rangle = \langle \mathbf{v}, T^T(\mathbf{w}) \rangle$, $\langle T\mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{v}, T^T\mathbf{w} \rangle$ or $\langle A\mathbf{v}, \mathbf{w} \rangle = \langle \mathbf{v}, A^T\mathbf{w} \rangle$ where A is a general operator (instead of a matrix) on V .

Definition 5.5.7 Suppose V is a finite-dimensional vector space over field $\mathcal{K}$ and T is an operator on V . Suppose $\mathcal{B}$ is a basis for V and let $\mathcal{M}$ be the matrix representation of T with respect to basis $\mathcal{B}$, then we define the **determinant of operator T** to be the determinant of matrix $\mathcal{M}$, i.e., $\det(T) = \det(\mathcal{M})$.

Remark 5.5.8 Note that the determinant of an operator is well-defined since the value of the determinant of the matrix representation of T does not depend on the choice of basis $\mathcal{B}$: suppose that if a different basis $\mathcal{B}'$ is chosen and $\mathcal{M}'$ is the matrix representation of T with respect to $\mathcal{B}'$, then we have $\mathcal{M}' = P^{-1}\mathcal{M}P$ for some invertible

matrix P and thus

$$\det(\mathcal{M}') = \det(P^{-1}\mathcal{M}P) = \det(P^{-1}) \det(\mathcal{M}) \det(P) = \det(\mathcal{M}).$$

Theorem 5.5.9 Suppose $V, \mathcal{K}, \langle \cdot, \cdot \rangle$ satisfy the condition in Theorem 5.5.5. Let A, B be operators on V and $c \in \mathcal{K}$. Then

1. $(A + B)^T = A^T + B^T$.
2. $(cA)^T = cA^T$.
3. $(AB)^T = B^T A^T$, where AB means the composite operator: $V \rightarrow V : AB(x) = (A \circ B)(x) = A(B(x))$.
4. $(A^T)^T = A$.

Definition 5.5.10 Suppose $V, \mathcal{K}, \langle \cdot, \cdot \rangle$ satisfy the condition in Theorem 5.5.5 and A is an operator on V . We say A is **symmetric** if $A^T = A$.

Lemma 5.5.11 Suppose V is a finite-dimensional vector space over $\mathbb{C}$ with a positive definite hermitian product $\langle \cdot, \cdot \rangle$. Given a functional L on V , there exists a unique vector $\tilde{w} \in V$ such that $L(x) = \langle x, \tilde{w} \rangle$ for all $x \in V$.

Theorem 5.5.12 Suppose $V, \mathbb{C}, \langle \cdot, \cdot \rangle$ satisfy the condition in Lemma 5.5.11 and A is an operator on V . Then there exists a unique operator A^* on V such that $\langle Ax, y \rangle = \langle x, A^*y \rangle$ for all $x, y \in V$.

Example 5.5.13 Let $V = \mathbb{C}^n$ with the standard hermitian product $\langle x, y \rangle = \sum_{i=1}^n x_i \overline{y_i}$. For any $A \in \mathbf{M}_{n \times n}(\mathbb{C})$, we have

$$\langle Ax, y \rangle = x^T A^T \overline{y} = x^T \overline{\overline{A^T y}} = \langle x, \overline{A^T y} \rangle.$$

Thus $A^* = \overline{A^T}$.

Definition 5.5.14 Suppose $V, \mathbb{C}, \langle \cdot, \cdot \rangle$ satisfy the condition in Lemma 5.5.11. An operator A on V is called **self-adjoint** or **hermitian** if $A^* = A$. Particularly, if A is hermitian, then $\langle Ax, y \rangle = \langle x, Ay \rangle$ for all $x, y \in V$.

Theorem 5.5.15 Suppose $V, \mathbb{C}, \langle \cdot, \cdot \rangle$ satisfy the condition in Lemma 5.5.11. Let A and B be operators on V and $\alpha \in \mathbb{C}$. Then

1. $(A + B)^* = A^* + B^*$.
2. $(\alpha A)^* = \overline{\alpha} A^*$.
3. $(AB)^* = B^* A^*$, where AB means the composite operator: $V \rightarrow V : AB(x) = (A \circ B)(x) = A(B(x))$.
4. $(A^*)^* = A$.

Definition 5.5.16 Suppose V is a finite-dimensional vector space over $\mathbb{R}$ with a positive definite scalar product $\langle \cdot, \cdot \rangle$. An operator A on V is called (real-) **unitary** or **orthogonal** if $\langle Ax, Ay \rangle = \langle x, y \rangle$ for all $x, y \in V$.

Theorem 5.5.17 Suppose $V, \mathbb{R}, \langle \cdot, \cdot \rangle$ satisfy the condition in Definition 5.5.16. Let A be an operator on V . Then the following conditions on A are equivalent:

- (a) A is real-unitary, i.e., $\langle Ax, Ay \rangle = \langle x, y \rangle$ for all $x, y \in V$.
- (b) A preserves norm of vectors, i.e., $\|Ax\| = \|x\|$ for all $x \in V$.
- (c) If $v \in V$ is a unit vector, then Av is also a unit vector.
- (d) $A^T A = I$, where I is the identity operator on V .

Proof. (a) $\Rightarrow$ (b): by definition $\|Av\| = \sqrt{\langle Av, Av \rangle} = \sqrt{\langle v, v \rangle} = \|v\|$ for any $v \in V$.

(b) $\Rightarrow$ (c): trivial.

(d) $\Rightarrow$ (a): for any $x, y \in V$, we have $\langle Ax, Ay \rangle = \langle x, A^T Ay \rangle = \langle x, y \rangle$.

(b) $\Rightarrow$ (a): for any $x, y \in V$, we have

$$\begin{aligned} \langle Ax, Ay \rangle &= \frac{1}{4} (\|Ax + Ay\|^2 - \|Ax - Ay\|^2) \\ &= \frac{1}{4} (\|x + y\|^2 - \|x - y\|^2) = \langle x, y \rangle. \end{aligned}$$

(c) $\Rightarrow$ (b):

Case 1: if $x = 0$, then $\|Ax\| = \|0\| = \|x\|$.

Case 2: if $x \neq 0$, then $\frac{x}{\|x\|}$ is a unit vector and thus

$$\left\| A \frac{x}{\|x\|} \right\|^2 = 1 = \left\langle \frac{x}{\|x\|}, \frac{x}{\|x\|} \right\rangle = \frac{1}{\|x\|^2} \langle x, x \rangle = \frac{\|x\|^2}{\|x\|^2} = 1.$$

Thus $\|Ax\| = \|x\|$.

(a) $\Rightarrow$ (d): for any $x, y \in V$, we have

$$\langle Ax, Ay \rangle = \langle x, y \rangle = \langle x, A^T Ay \rangle \implies \langle x, (A^T A - I)y \rangle = 0.$$

By the non-degeneracy of the scalar product, we have $(A^T A - I)y = 0$ for all $y \in V$ and thus $A^T A = I$. $\square$

Example 5.5.18 Let

$$R(\theta) = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}.$$

For a vector $x \in \mathbb{R}^2$, Rx rotated counterclockwise by θ . We also call such a matrix unitary if the associated operator is unitary.

Definition 5.5.19 Suppose V is a finite-dimensional vector space over $\mathbb{C}$ with a positive definite hermitian product $\langle \cdot, \cdot \rangle$. An operator A on V is called (complex-)unitary if $\langle Ax, Ay \rangle = \langle x, y \rangle$ for all $x, y \in V$.

Theorem 5.5.20 Suppose $V, \mathbb{C}, \langle \cdot, \cdot \rangle$ satisfy the condition in Definition 5.5.19. Then the following conditions on an operator A on V are equivalent:

- (a) A is complex-unitary, i.e., $\langle Ax, Ay \rangle = \langle x, y \rangle$ for all $x, y \in V$.
- (b) A preserves norm of vectors, i.e., $\|Ax\| = \|x\|$ for all $x \in V$.

(c) If $v \in V$ is a unit vector, then Av is also a unit vector.

(d) $A^*A = I$, where I is the identity operator on V .

Proposition 5.5.21 Let V be a finite-dimensional vector space over $\mathbb{R}$ with a positive definite scalar product $\langle \cdot, \cdot \rangle$, or over $\mathbb{C}$ with a positive definite hermitian product $\langle \cdot, \cdot \rangle$. Suppose A is an operator on V . Let $\{v_1, v_2, \dots, v_n\}$ be an orthonormal basis for V .

(a) If A is real-/complex-unitary, then $\{Av_1, Av_2, \dots, Av_n\}$ is also an orthonormal basis for V .

(b) If $\{Av_1, Av_2, \dots, Av_n\}$ is also an orthonormal basis for V , then A is real-/complex-unitary.

Proof. For (2), we can prove that A is unitary by showing that $\|Av\| = \|v\|$ for any $v \in V$. Since $\{v_1, v_2, \dots, v_n\}$ is an orthonormal basis for V , we can write $v = \sum_{i=1}^n \alpha_i v_i$ for some unique α_i for $i = 1, 2, \dots, n$. Then we have

$$\|v\|^2 = \langle v, v \rangle = \sum_{i=1}^n |\alpha_i|^2, \quad \|Av\|^2 = \langle Av, Av \rangle = \sum_{i=1}^n |\alpha_i|^2.$$

Thus $\|Av\| = \|v\|$ and A is unitary. □

Proposition 5.5.22 Suppose $V, \mathbb{C}, \langle \cdot, \cdot \rangle$ satisfy the condition in Definition 5.5.19. Then the following conditions on an operator A on V are equivalent:

1. $AA^* = A^*A$.
2. $\|Av\| = \|A^*v\|$ for all $v \in V$.
3. We can write $A = S + iT$ where S and T are self-adjoint operators on V such that $ST = TS$. Moreover,

$$S = \frac{A + A^*}{2}, \quad T = \frac{A - A^*}{2i}.$$

6 Eigenvectors and eigenvalues

Recall: in Definition 4.2.10 we defined the concept of eigenvectors and eigenvalues. After introducing operators, we can discuss more properties of eigenvectors and eigenvalues in this subsection.

Definition 5.6.1 Suppose V is a vector space over $\mathcal{K}$ and A is an operator on V . We say $\lambda \in \mathcal{K}$ is an **eigenvalue** of A if there exists a nonzero vector $v \in V$ such that $Av = \lambda v$. In this case, we call v an **eigenvector** of A corresponding to eigenvalue λ .

Remark 5.6.2 Note that for a given eigenvector v of A , the corresponding eigenvalue λ is unique since $v \neq 0$. However, for a given eigenvalue λ , the corresponding eigenvectors are not unique.

Definition 5.6.3 E_λ or V_λ is defined to be a subspace of V : $E_\lambda = \{v \in V : Av = \lambda v\}$ when λ is an eigenvalue of A . We call E_λ the **eigenspace** of A corresponding to λ .

Example 5.6.4 1. Suppose V is the vector space over $\mathbb{R}$ consisting of all infinitely differentiable functions defined on $\mathbb{R}$. Let $D : V \rightarrow V$ be the differentiation operator, i.e., $D(f) = f'$. Then $e^{\lambda x}$ is an eigenvec-
tor/eigenfunction of D corresponding to eigenvalue λ for any $\lambda \in \mathbb{R}$.

2. Let

$$R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

be an operator on $\mathbb{R}^2$. Then

- if $\theta = k\pi$ for $k \in \mathbb{Z}$, then $R(\theta)$ has two eigenvalues 1 and -1 .
- if $\theta \neq k\pi$ for $k \in \mathbb{Z}$, then $R(\theta)$ has no eigenvalue.

Theorem 5.6.5 Let V be a vector space over $\mathcal{K}$ and A be an operator on V . Suppose $\{v_1, \dots, v_m\}$ are eigen-
vectors of V corresponding to eigenvalues $\lambda_1, \dots, \lambda_m$ respectively. If $\lambda_i \neq \lambda_j$ for any $i \neq j$, then $\{v_1, \dots, v_m\}$
is linearly independent.

Proof. We can prove the theorem by induction on m . The case $m = 1$ is trivial.

Inductive step: suppose the statement holds for case m and we will prove for the case $m + 1$. Suppose $\alpha_1, \alpha_2, \dots, \alpha_{m+1} \in \mathcal{K}$ such that

$$\sum_{i=1}^{m+1} \alpha_i v_i = 0.$$

Applying λ_{m+1} to both sides of the above equation, we have

$$\sum_{i=1}^{m+1} \alpha_i \lambda_{m+1} v_i = 0.$$

Applying A to both sides of the first equation, we have

$$\sum_{i=1}^{m+1} \alpha_i A v_i = \sum_{i=1}^{m+1} \alpha_i \lambda_i v_i = 0.$$

Thus

$$\sum_{i=1}^m \alpha_i (\lambda_{m+1} - \lambda_i) v_i = 0.$$

By the inductive hypothesis, we have $\alpha_i (\lambda_{m+1} - \lambda_i) = 0$ for $i = 1, 2, \dots, m$. Since $\lambda_i \neq \lambda_{m+1}$ for $i = 1, 2, \dots, m$, we have $\alpha_i = 0$ for $i = 1, 2, \dots, m$. Thus $\alpha_{m+1} v_{m+1} = 0$ and thus $\alpha_{m+1} = 0$ since v_{m+1} is nonzero. We have shown that $\alpha_1 = \alpha_2 = \dots = \alpha_{m+1} = 0$ and thus $\{v_1, v_2, \dots, v_{m+1}\}$ is linearly independent. $\square$

Proposition 5.6.6 Suppose V with $\dim V = n$ and A is an operator on V . If A has n distinct eigenvalues, then the corresponding eigenvectors form a basis $\mathcal{B}$ for V . Moreover, the matrix representation of A with respect to

basis $\mathcal{B}$ is a diagonal matrix with diagonal entries being the eigenvalues of A , i.e.,

$$\mathcal{M}_{\mathcal{B}}(A) = \begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ 0 & \lambda_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & \lambda_n \end{bmatrix}.$$

Remark 5.6.7 Note

A has n distinct eigenvalues $\implies A$ has an eigenbasis.

However,

A has an eigenbasis $\not\Rightarrow A$ has n distinct eigenvalues.

For example, let $A = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ be an operator on $\mathbb{R}^2$. Then A has only one eigenvalue 0 but the corresponding eigenspace is $\mathbb{R}^2$ and thus any basis of $\mathbb{R}^2$ is an eigenbasis for A .

Theorem 5.6.8 Let V be a finite-dimensional vector space over $\mathcal{K}$ and A be an operator on V . Suppose $\lambda \in \mathcal{K}$. Then λ is an eigenvalue of A if and only if $A - \lambda I$ is not invertible, where I is the identity operator on V .

Definition 5.6.9 Let V be a vector space with $\dim V = n$ and A be an operator on V . Then

$$P_A(t) = \det(A - tI)$$

is called the **characteristic polynomial** of A . Note that P_A is a polynomial of degree n . Suppose

$$P_A(t) = (t - \lambda_1)^{m_1} (t - \lambda_2)^{m_2} \cdots (t - \lambda_k)^{m_k} \quad \text{with} \quad \sum_{i=1}^k m_i = n$$

where $\lambda_1, \lambda_2, \dots, \lambda_k$ are distinct eigenvalues of A and m_i is the multiplicity of λ_i for $i = 1, 2, \dots, k$. Then we call m_i the **algebraic multiplicity** of eigenvalue λ_i . We also define the **geometric multiplicity** of eigenvalue λ_i to be $E_{\lambda} = \ker(A - \lambda I)$, i.e., how many linearly independent eigenvectors of A corresponding to λ_i .

Remark 5.6.10 There're two cases that a matrix A cannot be diagonalized, i.e., A does not have an eigenbasis:

- A has no roots in $\mathbb{R}$, i.e., A has no eigenvalue. We can expand the field $\mathbb{R}$ to $\mathbb{C}$ and then A has eigenvalues in $\mathbb{C}$ since $\mathbb{C}$ is algebraically closed.
- A has eigenvalues but does not have enough linearly independent eigenvectors. For example, let $A = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$. Since the characteristic polynomial

$$\det(tI - A) = \det \begin{pmatrix} t - 2 & -1 \\ 0 & t - 2 \end{pmatrix} = (t - 2)^2.$$

Thus A has only one eigenvalue 2 with algebraic multiplicity 2. However, the eigenspace corresponding to eigenvalue 2 is $\{(x, 0) : x \in \mathbb{R}\}$ and thus the geometric multiplicity of eigenvalue 2 is 1. Thus A does not have an eigenbasis and thus cannot be diagonalized.

Theorem 5.6.11 If $A \in \mathbf{M}_{m \times n n \times n}(\mathbb{R})$ is symmetric, i.e., $A^T = A$, and $\lambda \in \mathbb{C}$ is an eigenvalue of A , then $\lambda \in \mathbb{R}$.

Proof. Suppose $\lambda \in \mathbb{C}$ and $x \in \mathbb{C}^n$ such that $Ax = \lambda x$. With the standard Hermitian product, we have

$$\begin{aligned}\langle Ax, x \rangle &= \langle \lambda x, x \rangle = \lambda \langle x, x \rangle, \\ \langle Ax, x \rangle &= \langle x, A^* x \rangle = \langle x, \overline{A^T} x \rangle = \langle x, \overline{Ax} \rangle = \langle x, \overline{\lambda x} \rangle = \overline{\lambda} \langle x, x \rangle.\end{aligned}$$

Since $\langle x, x \rangle > 0$, we have $\lambda = \overline{\lambda}$ and thus $\lambda \in \mathbb{R}$. □

Remark 5.6.12 Note

- the eigenvectors for a symmetric real matrix are not necessarily real.
- Suppose $Ax = \lambda x$ for some $x = u + iv$ where $u, v \in \mathbb{R}^n$. Every eigenvalue λ of A must have an eigenvector in $\mathbb{R}^n$.

Definition 5.6.13 Let V be a finite-dimensional vector space over $\mathbb{R}$ with $\dim V \geq 1$ and a positive definite scalar product $\langle \cdot, \cdot \rangle$. Let A be a symmetric operator on V . We say a subspace W of V is **invariant** under A if $Aw \in W$ for all $w \in W$. In this case, we also say A preserves W .

Lemma 5.6.14 Let V and A be defined in Definition 5.6.13. If W is a subspace of V invariant under A , then the orthogonal complement $W^\perp$ of W is also invariant under A .

Proof. Suppose W is invariant under A . By definition, for any $w \in W$, $Aw \in W$. Let $v \in W^\perp$ and we will show that $Av \in W^\perp$. Since A is symmetric, we have $\langle Av, w \rangle = \langle v, Aw \rangle = 0$ for any $w \in W$. Thus $Av \in W^\perp$ and thus $W^\perp$ is invariant under A . □

Theorem 5.6.15 Let V be a finite-dimensional vector space over $\mathbb{R}$ with $\dim V \geq 1$ and a positive definite scalar product $\langle \cdot, \cdot \rangle$. If A is a symmetric operator on V , then A has a orthonormal eigenbasis.

Proof. We prove by induction on $\dim V$.

Initial case: By the real version of Theorem 5.6.11, A has a real eigenvalue λ and we pick an eigenvector with length 1 to form an orthonormal basis.

Inductive step: suppose the statement holds for V with $\dim V = n$. We need to prove for the case $\dim V = n+1$. Note that we can pick an eigenvector v_1 of A with $\|v_1\| = 1$ due to the same reasoning in initial case. Let $V_1 = \text{span}\{v_1\}$ and V_1 is invariant under A . By the previous lemma, the orthogonal complement $V_1^\perp$ of V_1 is also invariant under A . Since $V_1 \oplus V_1^\perp = V$ and $\dim V_1^\perp = n$, by the inductive hypothesis, the restricted $A : V_1^\perp \rightarrow V_1^\perp$ has an orthonormal eigenbasis $\{v_2, v_3, \dots, v_{n+1}\}$ for $V_1^\perp$. Thus $\{v_1, v_2, \dots, v_{n+1}\}$ is an orthonormal eigenbasis for V . □

Remark 5.6.16 A real symmetric matrix/operator is always diagonalizable with an orthonormal eigenbasis. Particularly, suppose $A \in \mathbf{M}_{m \times n n \times n}(\mathbb{R})$ is symmetric, let $\{x_1, x_2, \dots, x_n\}$ be an orthonormal eigenbasis of A

and let $\lambda_1, \lambda_2, \dots, \lambda_n$ be the corresponding eigenvalues. Then we have

$$AX = X\Lambda, \quad \text{where } X = \begin{pmatrix} ||| \\ x_1 x_2 \cdots x_n \\ ||| \end{pmatrix}, \quad \Lambda = \begin{pmatrix} \lambda_1 0 \cdots 0 \\ 0 \lambda_2 \cdots 0 \\ \vdots \vdots \vdots \\ 0 0 \cdots \lambda_n \end{pmatrix}.$$

Note that

$$X^T X = \begin{pmatrix} \langle x_1, x_1 \rangle \langle x_1, x_2 \rangle \cdots \langle x_1, x_n \rangle \\ \langle x_2, x_1 \rangle \langle x_2, x_2 \rangle \cdots \langle x_2, x_n \rangle \\ \vdots \vdots \vdots \\ \langle x_n, x_1 \rangle \langle x_n, x_2 \rangle \cdots \langle x_n, x_n \rangle \end{pmatrix} = \begin{pmatrix} 1 0 \cdots 0 \\ 0 1 \cdots 0 \\ \vdots \vdots \vdots \\ 0 0 \cdots 1 \end{pmatrix} = I.$$

Thus X is unitary and we call A **unitarily/orthogonally diagonalizable**. Since Λ is diagonal, we say A is **unitarily similar to a diagonal matrix** and $A = X\Lambda X^T = X\Lambda X^{-1}$ is a **spectral decomposition** of A with $\lambda_1, \lambda_2, \dots, \lambda_n$ being the **spectrum** of A .

The converse of Theorem 5.6.15 is also true: if $A \in \mathbf{M}_{m \times n n \times n}(\mathbb{R})$ has an orthonormal eigenbasis, i.e., $A = X\Lambda X^T$ for some unitary matrix X and diagonal matrix Λ , then A is symmetric since $A^T = (X\Lambda X^T)^T = X\Lambda X^T = A$.

Theorem 5.6.17 Let V be a finite-dimensional vector space over $\mathbb{C}$ with $\dim V \geq 1$ and a positive definite hermitian product $\langle \cdot, \cdot \rangle$. If A is a hermitian operator on V and $\lambda \in \mathbb{C}$ is an eigenvalue of A , then $\lambda \in \mathbb{R}$.

Definition 5.6.18 Let V be as in Theorem 5.6.17 and A be a hermitian operator on V . We say a subspace W of V is **invariant** under A if $Aw \in W$ for all $w \in W$. In this case, we also say A preserves W .

Lemma 5.6.19 Let V and A be defined in the above definition. If W is a subspace of V invariant under A , then the orthogonal complement $W^\perp$ of W is also invariant under A .

Theorem 5.6.20 Let V be a finite-dimensional vector space over $\mathbb{C}$ with $\dim V \geq 1$ and a positive definite hermitian product $\langle \cdot, \cdot \rangle$. If A is a hermitian operator on V , then A has an orthonormal eigenbasis.

Remark 5.6.21 If $A \in \mathbf{M}_{m \times n n \times n}(\mathbb{C})$ is hermitian, then A is unitarily diagonalizable, i.e., there exists a unitary matrix X such that X^*AX is diagonal. Moreover, the converse also holds: if $A \in \mathbf{M}_{m \times n n \times n}(\mathbb{C})$ is unitarily diagonalizable, then A is hermitian.

Lemma 5.6.22 Let V be a finite-dimensional vector space over $\mathbb{C}$ with $\dim V \geq 1$ and a positive definite hermitian product $\langle \cdot, \cdot \rangle$. If A is a unitary operator on V and W is invariant under A , then the orthogonal complement $W^\perp$ of W is also invariant under A .

Proof. Suppose $Aw \in W$ for any $w \in W$ and A is unitary. Let $v \in W^\perp$, then $\langle v, w \rangle = 0$ for any $w \in W$. We want to show $\langle Av, w \rangle = \langle v, A^*w \rangle = \langle v, A^{-1}w \rangle = 0$ for all $w \in W$, i.e., $Av \in W^\perp$. Note that if we restrict A to be on W , i.e., $A_W : W \rightarrow W$, then $0 \in \ker A_W \leq \ker A = \{0\}$, then A_W is an isomorphism on W . Hence, $A_W^{-1} : W \rightarrow W$ is also an isomorphism and $A^{-1}w \in W$ for any $w \in W$. Thus $\langle v, A^{-1}w \rangle = 0$ for any $w \in W$

and thus $Av \in W^\perp$. □

Theorem 5.6.23 For the complex case, if A is a complex-unitary operator on V , then A has an orthonormal eigenbasis. Particularly, if $A \in \mathbf{M}_{m \times nn \times n}(\mathbb{C})$ is unitary, then $A = U\Lambda U^{-1} = U\Lambda U^*$ for some complex-unitary matrix U and diagonal matrix Λ .

Remark 5.6.24 The eigenvalues of a complex-unitary operator must satisfy $|\lambda| = 1$:

$$\langle Av, Av \rangle = \langle \lambda v, \lambda v \rangle = \lambda \bar{\lambda} \langle v, v \rangle = |\lambda|^2 \langle v, v \rangle.$$

Since A is unitary, we have $\langle Av, Av \rangle = \langle v, v \rangle$ and thus $|\lambda| = 1$.

Remark 5.6.25 A complex-unitary matrix/operator is not necessarily hermitian:

$$A = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad A^* = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \neq A.$$

A hermitian matrix is not necessarily complex-unitary:

$$A = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}, \quad A^* = A, \quad \text{but } A^*A = \begin{pmatrix} 1 & 0 \\ 0 & 4 \end{pmatrix} \neq I.$$

Theorem 5.6.26 For the real case, if A is a real-unitary operator on V , then V can be expressed as $V = V_1 \oplus V_2 \oplus \cdots \oplus V_r$ where

1. each V_i is a subspace of V invariant under A
2. each V_i is mutually orthogonal to each other, i.e., $\langle v_i, v_j \rangle = 0$ for any $v_i \in V_i$ and $v_j \in V_j$ with $i \neq j$
3. $\dim V_i = 1$ or 2 for $i = 1, 2, \dots, r$.

Proof. We shall only prove the case $V = \mathbb{R}^n$ with the standard scalar product and A is a real unitary operator. The general case can be reduced to this case by choosing an orthonormal basis for V and consider the matrix representation of A with respect to this basis.

We view A as a complex-unitary matrix, i.e., $A \in \mathbf{M}_{m \times nn \times n}(\mathbb{C})$ and $A^*A = I$. Suppose $z \neq 0$, $z \in \mathbb{C}^n$ is an eigenvector of A corresponding to eigenvalue $\lambda \in \mathbb{C}$. Namely, $Az = \lambda z$ and $|\lambda| = 1$. Let $z = x + iy$ for $x, y \in \mathbb{R}^n$ and $\lambda = \cos \theta + i \sin \theta$ for some $\theta \in \mathbb{R}$. We discuss by case.

Case 1: if λ is real, then $\lambda = \pm 1$. Then $Ax + iAy = A(x + iy) = \lambda(x + iy) = \lambda x + i\lambda y$. Since $Ax, Ay, \lambda x, \lambda y$ are all real vectors, we have $Ax = \lambda x$ and $Ay = \lambda y$. We can let v to be a non-zero vector in $\{x, y\}$ and $v_1 = \frac{v}{\|v\|}$.

We let $V_1 = \text{span}\{v_1\}$, then V_1 is a one-dimensional subspace of $\mathbb{R}^n$ invariant under A . So $V_1^\perp$ is also invariant under A and $V = V_1 \oplus V_1^\perp$.

Case 2: if λ is not real, then $Az = \lambda z$ and $A\bar{z} = \bar{\lambda}\bar{z}$ since A is a real matrix. Thus $\bar{z}$ is also an eigenvector of A corresponding to eigenvalue $\bar{\lambda}$. Since $\{z, \bar{z}\}$ are linearly independent, we know $\{x, y\}$ are linearly independent. Note

$$\begin{cases} Ax = \cos \theta x - \sin \theta y \in \text{span}\{x, y\}, \\ Ay = \sin \theta x + \cos \theta y \in \text{span}\{x, y\}. \end{cases}$$

Let $V_1 = \text{span}\{x, y\}$, then V_1 is A -invariant and $\dim V_1 = 2$. So $V_1^\perp$ is also invariant under A and $V = V_1 \oplus V_1^\perp$. By induction on n , we know V can be reduced to the direct sum of A -invariant subspaces with dimension 1 or 2. $\square$

Corollary 5.6.27 For the real case, if A is a real-unitary operator on V , then there exists a basis $\mathcal{B}$ for V such that

$$\mathcal{M}_{\mathcal{B}}(A) = \begin{pmatrix} M_1 & 0 & \cdots & 0 \\ 0 & M_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & M_r \end{pmatrix}$$

where each M_i for $i = 1, 2, \dots, r$ satisfies one of the following conditions:

$$M_i = 1, \quad M_i = -1, \quad M_i = \begin{pmatrix} \cos \theta_i & -\sin \theta_i \\ \sin \theta_i & \cos \theta_i \end{pmatrix} \text{ with } \theta_i \neq k\pi \text{ for } k \in \mathbb{Z}.$$

Remark 5.6.28 Operators that have orthonormal eigenbasis $\mathcal{B}$, i.e., symmetric operators, hermitian operators and complex-unitary operators, are easy to handle. However, some operators do not have orthonormal eigenbasis (or even just an eigenbasis), but it's still possible to pick a basis $\mathcal{B}$ such that $\mathcal{M}_{\mathcal{B}}(A)$ has a nice form, i.e., “is close to being diagonal”.

7 Polynomials and Triangulation of Matrices and Operators

Recall that the set of all polynomials $\mathbf{P}_n(\mathcal{K})$ with coefficients in $\mathcal{K}$ is a vector space with $\dim \mathbf{P}_n(\mathcal{K}) = n + 1$. Also, the set of all polynomials $\mathcal{K}[x]$ with coefficients in $\mathcal{K}$ is a vector space with $\dim \mathcal{K}[x] = \infty$.

Definition 5.7.1 Suppose $A \in \mathbf{M}_{m \times n n \times n}(\mathcal{K})$ and $f \in \mathcal{K}[t]$ with $f(t) = a_n t^n + a_{n-1} t^{n-1} + \cdots + a_1 t + a_0$ for some non-negative integer n and $a_i \in \mathcal{K}$ for $i = 0, 1, \dots, n$. We define

$$f(A) = a_n A^n + a_{n-1} A^{n-1} + \cdots + a_1 A + a_0 I_n,$$

and call it a **polynomial of matrix** A , where I_n is the identity matrix of size n . We can also define $f(A)$ for any operator A on a vector space V by the same formula where $A^k := A \circ A \circ \cdots \circ A$ (k times) and I is the identity operator on V .

Theorem 5.7.2 Suppose $f, g \in \mathcal{K}[t]$ and $A \in \mathbf{M}_{m \times n n \times n}(\mathcal{K})$. Then we have

- (1) for any $\alpha, \beta \in \mathcal{K}$, we have $(\alpha f + \beta g)(A) = \alpha f(A) + \beta g(A)$
- (2) $(fg)(A) = f(A)g(A)$

Theorem 5.7.3 Suppose $A \in \mathbf{M}_{m \times n n \times n}(\mathcal{K})$. Then there exists a nonzero polynomial $f \in \mathcal{K}[t]$ such that $f(A) = 0$.

Proof. Note $\dim \mathbf{M}_{m \times n n \times n}(\mathcal{K}) = n^2$. For any $N \geq n^2$, the set $\{I, A, A^2, \dots, A^N\}$ is linearly dependent. Thus there exist $\alpha_0, \alpha_1, \dots, \alpha_N \in \mathcal{K}$ such that $\alpha_0 I + \alpha_1 A + \dots + \alpha_N A^N = 0$. Let $f(t) = \alpha_N t^N + \dots + \alpha_1 t + \alpha_0$, then $f(A) = 0$. $\square$

Remark 5.7.4 The above theorems also hold for operators on a finite-dimensional vector space V over $\mathcal{K}$.

Definition 5.7.5 Suppose V is a finite-dimensional vector space over $\mathcal{K}$ with $\dim V \geq 1$ and A is an operator on V . A set of subspaces $\{V_1, V_2, \dots, V_n\}$ of V is called a **fan** of A in V if

- (1) $V_i \subseteq V_{i+1}$ for $i = 1, 2, \dots, n-1$,
- (2) $\dim V_i = i$ for $i = 1, 2, \dots, n$,
- (3) V_i is A -invariant for $i = 1, 2, \dots, n$, i.e., $Av \in V_i$ for any $v \in V_i$ and $i = 1, 2, \dots, n$.

Let $\{V_1, V_2, \dots, V_n\}$ be a fan of A in V . A basis $\{v_1, \dots, v_i\}$ of V is called a **fan basis** if $\{v_1, \dots, v_i\}$ is a basis for V_i for $i = 1, 2, \dots, n$.

Remark 5.7.6 Note for a given fan of A , a fan basis always exists and is not unique. However, a fan may not exist.

Theorem 5.7.7 Let $\mathcal{B} = \{v_1, \dots, v_n\}$ be a fan basis for a operator A . Then

$$\mathcal{M}_{\mathcal{B}}(A) = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ 0 & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{nn} \end{pmatrix}$$

is an upper-triangular matrix.

Proof. Since $V_i = \text{span}\{v_1, \dots, v_i\}$ is invariant under A for $i = 1, 2, \dots, n$, we have $Av_i \in V_i$ for $i = 1, 2, \dots, n$. Thus $Av_i = a_{1i}v_1 + a_{2i}v_2 + \dots + a_{ii}v_i$ for some $a_{1i}, a_{2i}, \dots, a_{ii} \in \mathcal{K}$ and thus $\mathcal{M}_{\mathcal{B}}(A)$ is an upper-triangular matrix. $\square$

Remark 5.7.8 If $\mathcal{C} = \{u_1, \dots, u_n\}$ is a basis of V such that $\mathcal{M}_{\mathcal{C}}(A)$ is an upper-triangular matrix, then $\mathcal{C}$ is a fan basis for A by letting $V_i = \text{span}\{u_1, \dots, u_i\}$ for $i = 1, 2, \dots, n$.

Definition 5.7.9 Suppose V is a finite-dimensional vector space over $\mathcal{K}$ with $\dim V \geq 1$ and A is an operator on V . If there exists a basis $\mathcal{B}$ for V such that $\mathcal{M}_{\mathcal{B}}(A)$ is an upper-triangular matrix, then we say A is **triangulable** and $\mathcal{B}$ is a **triangulation basis** for A .

Theorem 5.7.10

Appendix A Chinese-English glossary

中文	English
域	Field
子域	Subfield
加法单位元	Additive identity element
乘法单位元	Multiplicative identity element
加法逆元	Additive inverse
乘法逆元 / 倒数	Multiplicative inverse / Reciprocal
封闭性	Closure
结合律	Associative law
交换律	Commutative law
分配律	Distributive law
数 / 标量	Number / Scalar
向量空间	Vector space
零向量	Zero vector
向量	Vector
子空间	Subspace
线 (一维子空间)	Line (1-dimensional subspace)
平面 (二维子空间)	Plane (2-dimensional subspace)
线性组合	Linear combination
张成 / 生成子空间	Span
生成集 / 张成集	Spanning set / Set of generators
线性相关	Linearly dependent
线性无关	Linearly independent
极大线性无关组	Maximal linearly independent set / subset
基 / 基底	Basis
维数	Dimension
有限维向量空间	Finite-dimensional vector space
无限维向量空间	Infinite-dimensional vector space
分量 / 坐标	Components / Coordinates
坐标向量	Coordinate vector
多项式空间 $\mathcal{K}[x]$	Polynomial space $\mathcal{K}[x]$
多项式空间 $\mathbf{P}_n(\mathcal{K})$	Polynomial space $\mathbf{P}_n(\mathcal{K})$ (degree $\leq n$)
矩阵	Matrix
行向量	Row vector
列向量	Column vector
方阵	Square matrix
非方阵	Non-square matrix

中文	English
零矩阵	Zero matrix
矩阵加法	Matrix addition
矩阵数乘	Scalar multiplication of a matrix
矩阵空间 $\mathbf{M}_{m \times n}(\mathcal{K})$	Matrix space $\mathbf{M}_{m \times n}(\mathcal{K})$
标准基 (矩阵空间)	Standard basis of $\mathbf{M}_{m \times n}(\mathcal{K})$
单位矩阵 I_n	Identity matrix I_n
可逆矩阵 / 非奇异矩阵	Invertible (non-singular) matrix
奇异矩阵	Singular matrix
转置矩阵	Transpose of a matrix
对称矩阵	Symmetric matrix
对角矩阵	Diagonal matrix
线性方程组	Linear system
齐次线性方程组	Homogeneous linear system
非齐次线性方程组	Inhomogeneous / Nonhomogeneous linear system
解 (方程组)	Solution (of a system)
平凡解	Trivial solution
非平凡解	Nontrivial solution
线性映射 / 线性变换	Linear map / Linear transformation
平移	Translation (by a vector)
线性映射空间 $\mathcal{L}(V, W)$	Space of linear maps $\mathcal{L}(V, W)$
核 (线性映射的核)	Kernel of a linear map
像 (线性映射的像)	Image of a linear map
列空间 $\text{Col}(A)$	Column space $\text{Col}(A)$
零空间 $\text{Null}(A)$	Null space $\text{Null}(A)$
秩	Rank
零化度	Nullity
秩-零化度定理	Rank-Nullity Theorem
双射	Bijjective map
可逆映射	Invertible mapping
同构	Isomorphism
恒等映射 Id_V	Identity map Id_V
与矩阵对应的线性映射	Linear map associated with a matrix
线性映射的矩阵表示	Matrix representation of a linear map
对角化 (线性映射/矩阵)	Diagonalization (of a linear map / matrix)
可对角化线性映射 / 矩阵	Diagonalizable linear map / matrix
复合映射 $G \circ F$	Composition $G \circ F$ of linear maps
特征值	Eigenvalue
特征向量	Eigenvector
特征基	Eigenbasis
相似矩阵	Similar matrices
变换基矩阵	Change-of-basis matrix
标量积	Scalar product
非退化	Non-degenerate
正定	Positive-definite
范数	Norm
投影	Projection

中文	English
单位向量	Unit vector
归一化	Normalization
正交	Orthogonal
正交补	Orthogonal complement
正交基	Orthogonal basis
柯西-施瓦茨不等式	Cauchy-Schwarz inequality
三角不等式	Triangle inequality
格拉姆-施密特正交化过程	Gram-Schmidt process
贝塞尔不等式	Bessel's inequality