

1 Bera

Definition 1.1 (Lebesgue Integral). Let (X, \mathcal{M}, μ) be a measure space. Let $f : X \rightarrow \mathbb{C}$ be a $\mathcal{B}(\mathbb{C})/\mathcal{M}$ -measureable function. Define the following formulations of an integral:

- If f takes on only finitely many values in \mathbb{R}^+ , then give the standard representation of f as the simple function:

$$f(x) = \sum_i y_i \mathbb{1}_{\{f(x)=y_i\}}(x).$$

Then define the Lebesgue integral of f with respect to μ as:

$$\int_X f(x) d\mu(x) = \sum_i y_i \mu(\{f(x) = y_i\}).$$

- If f takes on values in \mathbb{R}^+ , then define the Lebesgue integral of f with respect to μ as:

$$\int_X f(x) d\mu(x) = \sup \left\{ \int_X g(x) d\mu(x) \mid g \leq f \text{ and } g \text{ is a simple function} \right\}.$$

This may or may not take on the “value” of ∞ .

- If f takes on values in \mathbb{R} , then give the functions $f^+ = \max\{f, 0\}$ and $f^- = \min\{f, 0\}$. Give the Lebesgue integral of f with respect to μ as:

$$\int_X f(x) d\mu(x) = \int_X f^+(x) d\mu(x) - \int_X f^- d\mu(x),$$

provided that both of the integrals on the right exist and at least one of them is finite.

- Lastly, if f takes on values in \mathbb{C} , then let u and v be the functions given by $u = \Re f$ and $v = \Im f$, i.e. $f(x) = u(x) + iv(x)$. Then define the Lebesgue integral of f relative to μ as:

$$\int_X f(x) d\mu(x) = \int_X u(x) d\mu(x) + i \int_X v(x) d\mu(x).$$

We usually write $\int f d\mu$ or just $\int f$ if the implied parts are easy to understand. Also we write $\int_A f$ to stand for $\int_X f \mathbb{1}_A$. In total, these parts define the **Lebesgue integral of f with respect to μ** .

Theorem 1.2 (Cauchy’s Integral Formula). Let U be an open subset of the \mathbb{C} and suppose that the disk

$$D(z_0, r) = \{z \in \mathbb{C} \mid |z - z_0| < r\}$$

is completely contained in U . Let $f : U \rightarrow \mathbb{C}$ be a holomorphic function and let γ be the circle oriented counterclockwise forming the boundary of $D(z_0, r)$. Then for every interior point a of $D(z_0, r)$,

$$f(a) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{za} dz.$$

Moreover, because $(z - a)^{-1}$ can be given by the power series:

$$\frac{1}{z-a} = \frac{1}{z} \sum_{n=0}^{\infty} \left(\frac{a}{z}\right)^n$$

it follows that f is analytic, infinitely differentiable, and

$$f^{(n)}(a) = \frac{n!}{2\pi i} \oint_{\gamma} \frac{f(z)}{(z-a)^{n+1}} dz.$$

Here are examples of the major fonts:

Regular ABCDEFGHIJKLMNOPQRSTUVWXYZ

abcdefghijklmnopqrstuvwxyz

1234567890

Slsshape ABCDEFGHIJKLMNOPQRSTUVWXYZ

abcdefghijklmnopqrstuvwxyz

1234567890

Bfseries ABCDEFGHIJKLMNOPQRSTUVWXYZ

abcdefghijklmnopqrstuvwxyz

1234567890

Slshape ABCDEFGHIJKLMNOPQRSTUVWXYZ

abcdefghijklmnopqrstuvwxyz

1234567890

Mathematics ABCDEFGHIJKLMNOPQRSTUVWXYZ

abcdefghijklmnopqrstuvwxyz

1234567890

Mathcal *A B C D E F G H I J K L M N O P Q R S T U V W X Y Z*

1/14

Mathbb ABCDEFGHIJKLMNOPQRSTUVWXYZ

פָּנָמִים וְעַמִּים

Mathds ABCDEFGHIJKLMNOPQRSTUVWXYZ

Ahk

1

Mathbbm ABCDEFGHIJKLMNOPQRSTUVWXYZ

a b c d e f g h i j k l m n o p q r s t u v w x y z

12

Mathfrak A B C D E F G H I J K L M N O P Q R S T U V W X Y Z

abcdefghijklmnopqrstuvwxyz

1234567890

Mathscr $\mathcal{A} \mathcal{B} \mathcal{C} \mathcal{D} \mathcal{E} \mathcal{F} \mathcal{G} \mathcal{H} \mathcal{I} \mathcal{J} \mathcal{K} \mathcal{L} \mathcal{M} \mathcal{N} \mathcal{O} \mathcal{P} \mathcal{Q} \mathcal{R} \mathcal{S} \mathcal{T} \mathcal{U} \mathcal{V} \mathcal{W} \mathcal{X} \mathcal{Y} \mathcal{Z}$

Greek αβγδεεζηθυικιλμνξπωρσστυφφχφψω

Greek (Capital) ΓΓΔΔΘΘΛΛΞΞΠΠΣΣΦΦΨΨΩΩ

Common symbols $\int \oint \iint \sum \prod \cup \cap \nabla \wedge \oplus \otimes$

Common symbols (displaystyle) $\int \oint \iiint \sum \prod \cup \cap \vee \wedge \oplus \otimes$

Inline Sup/Sub-scripts $\lim_{n \rightarrow \infty} \inf_{\alpha} \sup_{\substack{f: \mathbb{R} \rightarrow \mathbb{R} \\ f \text{ is } 1-\text{Lipschitz}}} \sum_{\alpha \in \mathcal{A}} \prod_{n=1}^{\infty} \int_a^b$

Inline Sup/Sub-scripts (displaystyle) $\lim_{n \rightarrow \infty} \inf_{\alpha} \sup_{\substack{f: \mathbb{R} \rightarrow \mathbb{R} \\ f \text{ is } 1-\text{Lipschitz}}} \sum_{\alpha \in \mathcal{A}} \prod_{n=1}^{\infty} \int_a^b$