


Higher 2000 - Paper 1 Solutions

- [SQA] 1. On the coordinate diagram shown, A is the point $(6, 8)$ and B is the point $(12, -5)$. Angle $AOC = p$ and angle $COB = q$.


Find the exact value of $\sin(p + q)$.



4

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
	4	C	NC	T9	$\frac{63}{65}$	2000 P1 Q1
•¹ ss: know to use trig expansion •² pd: process missing sides •³ ic: interpret data •⁴ pd: process				•¹ $\sin p \cos q + \cos p \sin q$ •² 10 and 13 •³ $\frac{8}{10} \cdot \frac{12}{13} + \frac{6}{10} \cdot \frac{5}{13}$ •⁴ $\frac{126}{130}$		


- [SQA] 2. A sketch of the graph of $y = f(x)$ where $f(x) = x^3 - 6x^2 + 9x$ is shown below.
The graph has a maximum at A and a minimum at B(3,0).



- (a) Find the coordinates of the turning point at A. 4
- (b) Hence sketch the graph of $y = g(x)$ where $g(x) = f(x + 2) + 4$.
Indicate the coordinates of the turning points. There is no need to calculate the coordinates of the points of intersection with the axes. 2
- (c) Write down the range of values of k for which $g(x) = k$ has 3 real roots. 1

Part	Marks	Level	Calc.	Content	Answer	U1 OC3
(a)	4	C	NC	C8	A(1,4)	2000 P1 Q2
(b)	2	C	NC	A3	sketch (translate 4 up, 2 left)	
(c)	1	A/B	NC	A2	$4 < k < 8$	
•¹ ss: know to differentiate •² pd: differentiate correctly •³ ss: know gradient = 0 •⁴ pd: process •⁵ ic: interpret transformation •⁶ ic: interpret transformation •⁷ ic: interpret sketch					•¹ $\frac{dy}{dx} = \dots$ •² $\frac{dy}{dx} = 3x^2 - 12x + 9$ •³ $3x^2 - 12x + 9 = 0$ •⁴ A = (1,4) translate $f(x)$ 4 units up, 2 units left •⁵ sketch with coord. of A'(-1,8) •⁶ sketch with coord. of B'(1,4) •⁷ $4 < k < 8$ (accept $4 \leq k \leq 8$)	

- [SQA] 3. Find the size of the angle a° that the line joining the points $A(0, -1)$ and $B(3\sqrt{3}, 2)$ makes with the positive direction of the x -axis.




3

Part	Marks	Level	Calc.	Content	Answer	U1 OC1
	3	C	NC	G2	30	2000 P1 Q3
¹ ss: know how to find gradient or equ. ² pd: process ³ ic: interpret exact value				¹ $\frac{2-(-1)}{3\sqrt{3}-0}$ ² $\tan a = \text{gradient}$ <i>stated or implied by</i> ³ $a = 30$		

- [SQA] 4. The diagram shows a sketch of the graphs of $y = 5x^2 - 15x - 8$ and $y = x^3 - 12x + 1$.

The two curves intersect at A and touch at B, i.e. at B the curves have a common tangent.



- (a) (i) Find the x -coordinates of the point of the curves where the gradients are equal. 4
- (ii) By considering the corresponding y -coordinates, or otherwise, distinguish geometrically between the two cases found in part (i). 1
- (b) The point A is $(-1, 12)$ and B is $(3, -8)$. 5
- Find the area enclosed between the two curves.

Part	Marks	Level	Calc.	Content	Answer	U2 OC2
(ai)	4	C	NC	C4	$x = \frac{1}{3}$ and $x = 3$	2000 P1 Q4
(aii)	1	C	NC	CGD	parallel and coincident	
(b)	5	C	NC	C17	$21\frac{1}{3}$	

•¹ ss: know to diff. and equate •² pd: differentiate •³ pd: form equation •⁴ ic: interpret solution •⁵ ic: interpret diagram •⁶ ss: know how to find area between curves •⁷ ic: interpret limits •⁸ pd: form integral •⁹ pd: process integration •¹⁰ pd: process limits	•¹ find derivatives and equate •² $3x^2 - 12$ and $10x - 15$ •³ $3x^2 - 10x + 3 = 0$ •⁴ $x = 3, x = \frac{1}{3}$ •⁵ tangents at $x = \frac{1}{3}$ are parallel, at $x = 3$ coincident •⁶ $\int(\text{cubic} - \text{parabola})$ or $\int(\text{cubic}) - \int(\text{parabola})$ •⁷ $\int_{-1}^3 \dots dx$ •⁸ $\int(x^3 - 5x^2 + 3x + 9)dx$ or equiv. •⁹ $[\frac{1}{4}x^4 - \frac{5}{3}x^3 + \frac{3}{2}x^2 + 9x]_{-1}^3$ or equiv. •¹⁰ $21\frac{1}{3}$
--	---

- [SQA] 5. Two sequences are generated by the recurrence relations $u_{n+1} = au_n + 10$ and $v_{n+1} = a^2v_n + 16$.

The two sequences approach the same limit as $n \rightarrow \infty$.

Determine the value of a and evaluate the limit.

5

Part	Marks	Level	Calc.	Content	Answer	U1 OC4
	4	C	NC	A13	$a = \frac{3}{5}, L = 25$	2000 P1 Q5
	1	A/B	NC	A12		
•¹ ss: know how to find limit •² pd: process •³ pd: process •⁴ ic: interpret coeff. of u_n •⁵ pd: process					•¹ $L = aL + 10$ or $L = a^2L + 16$ or $L = \frac{b}{1-a}$ •² $L = \frac{10}{1-a}$ or $L = \frac{16}{1-a^2}$ •³ $\frac{10}{1-a}$ or $\frac{16}{1-a^2}$ •⁴ $10a^2 - 16a + 6 = 0$ •⁵ $a = \frac{3}{5}$ and $L = 25$	

- [SQA] 6. For what range of values of k does the equation $x^2 + y^2 + 4kx - 2ky - k - 2 = 0$ represent a circle?

5

Part	Marks	Level	Calc.	Content	Answer	U2 OC4
	5	A	NC	G9, A17	for all k	2000 P1 Q6
•¹ ss: know to examine radius •² pd: process •³ pd: process •⁴ ic: interpret quadratic inequation •⁵ ic: interpret quadratic inequation					•¹ $g = 2k, f = -k, c = -k - 2$ <i>stated or implied by</i> •² •² $r^2 = 5k^2 + k + 2$ •³ (real $r \Rightarrow$) $5k^2 + k + 2 > 0$ (<i>accept</i> $\geq$) •⁴ use discr. or complete sq. or diff. •⁵ true for all k	


[SQA] 7. VABCD is a pyramid with a rectangular base ABCD.

Relative to some appropriate axes,

$$\overrightarrow{VA} \text{ represents } -7\mathbf{i} - 13\mathbf{j} - 11\mathbf{k}$$

$$\overrightarrow{AB} \text{ represents } 6\mathbf{i} + 6\mathbf{j} - 6\mathbf{k}$$

$$\overrightarrow{AD} \text{ represents } 8\mathbf{i} - 4\mathbf{j} + 4\mathbf{k}.$$



K divides BC in the ratio 1 : 3.

Find $\overrightarrow{VK}$ in component form.

3

Part	Marks	Level	Calc.	Content	Answer	U3 OC1
	3	C	CN	G25, G21, G20	$\begin{pmatrix} 1 \\ -8 \\ -16 \end{pmatrix}$	2000 P1 Q7
•¹ ss: recognise crucial aspect •² ic: interpret ratio •³ pd: process components				•¹ $\overrightarrow{VK} = \overrightarrow{VB} + \overrightarrow{BA} + \overrightarrow{AK}$ or $\overrightarrow{VK} = \overrightarrow{VB} + \overrightarrow{BK}$ •² $\overrightarrow{BK} = \frac{1}{4}\overrightarrow{BC}$ or $\frac{1}{4}\overrightarrow{AD}$ or $\begin{pmatrix} 2 \\ -1 \\ 1 \end{pmatrix}$ or $\begin{pmatrix} -1 \\ -7 \\ -17 \end{pmatrix}$ •³ $\overrightarrow{VK} = \begin{pmatrix} 1 \\ -8 \\ -16 \end{pmatrix}$		

[SQA] 8. The graph of $y = f(x)$ passes through the point $(\frac{\pi}{9}, 1)$.

If $f'(x) = \sin(3x)$ express y in terms of x .

4

Part	Marks	Level	Calc.	Content	Answer	U3 OC2
	4	A/B	NC	C18, C23	$y = -\frac{1}{3}\cos(3x) + \frac{7}{6}$	2000 P1 Q8
•¹ ss: know to integrate •² pd: integrate •³ ic: interpret $(\frac{\pi}{9}, 1)$ •⁴ pd: process				•¹ $y = \int \sin(3x) dx$ stated or implied by •² $-\frac{1}{3}\cos(3x)$ •³ $1 = -\frac{1}{3}\cos(\frac{3\pi}{9}) + c$ or equiv. •⁴ $c = \frac{7}{6}$		

[SQA] 9. Evaluate $\log_5 2 + \log_5 50 - \log_5 4$.

3

Part	Marks	Level	Calc.	Content	Answer	U3 OC3
	2	C	NC	A28	2	2000 P1 Q9
	1	A/B	NC	A28		
•¹ pd: use $\log_a x + \log_a y = \log_a xy$ •² pd: use $\log_a x - \log_a y = \log_a \frac{x}{y}$ •³ pd: use $\log_a a = 1$					•¹ $\log_5 100 - \log_5 4$ •² $\log_5 25$ •³ 2	

[SQA] 10. Find the maximum value of $\cos x - \sin x$ and the value of x for which it occurs in the interval $0 \leq x \leq 2\pi$.

6

Part	Marks	Level	Calc.	Content	Answer	U3 OC4
	6	A/B	CN	T14	max value $\sqrt{2}$ when $x = \frac{7\pi}{4}$	2000 P1 Q10
				•¹ ss: use e.g. $k \cos(x + a)$ •² ic: expand chosen rule •³ pd: compare coefficients •⁴ pd: process •⁵ pd: process •⁶ ic: interpret trig expression		
				•¹ e.g. use $k \cos(x + a)$ •² $k \cos x \cos a - k \sin x \sin a$ •³ $k \cos a = 1$ and $k \sin a = 1$ •⁴ $k = \sqrt{2}$ •⁵ $\tan a = 1, a = \frac{\pi}{4}$ (45° is bad form) •⁶ max. value = $\sqrt{2}$ when $x = \frac{7\pi}{4}$ (do not accept 45°)		


[END OF QUESTIONS]

Higher 2000 - Paper 2 Solutions

[SQA] 1. The diagram shows a sketch of the graph of $y = x^3 - 3x^2 + 2x$.

(a) Find the equation of the tangent to this curve at the point where $x = 1$.

(b) The tangent at the point $(2, 0)$ has equation $y = 2x - 4$. Find the coordinates of the point where this tangent meets the curve again.



5

5

Part	Marks	Level	Calc.	Content	Answer	U2 OC1
(a)	5	C	CN	C5	$x + y = 1$	2000 P2 Q1
(b)	5	C	CN	A23, A22, A21	$(-1, -6)$	

•¹ ss: know to differentiate •² pd: differentiate correctly •³ ss: know that gradient = $f'(1)$ •⁴ ss: know that y-coord = $f(1)$ •⁵ ic: state equ. of line •⁶ ss: equate equations •⁷ pd: arrange in standard form •⁸ ss: know how to solve cubic •⁹ pd: process •¹⁰ ic: interpret	•¹ $y' = \dots$ •² $3x^2 - 6x + 2$ •³ $y'(1) = -1$ •⁴ $y(1) = 0$ •⁵ $y - 0 = -1(x - 1)$ •⁶ $2x - 4 = x^3 - 3x^2 + 2x$ •⁷ $x^3 - 3x^2 + 4 = 0$ •⁸ $\begin{array}{r rrrr} \dots & 1 & -3 & 0 & 4 \\ & & \dots & \dots & \dots \\ & \dots & \dots & \dots & \dots \end{array}$ •⁹ identify $x = -1$ from working •¹⁰ $(-1, -6)$
---	--

[SQA] 2. (a) Find the equation of AB, the perpendicular bisector of the line joining the points $P(-3,1)$ and $Q(1,9)$.


(b) C is the centre of a circle passing through P and Q. Given that QC is parallel to the y -axis, determine the equation of the circle.

(c) The tangents at P and Q intersect at T.

Write down

(i) the equation of the tangent at Q

(ii) the coordinates of T.



Part	Marks	Level	Calc.	Content	Answer	U2 OC4
(a)	4	C	CN	G7	$x + 2y = 9$	2000 P2 Q2
(b)	3	C	CN	G10	$(x - 1)^2 + (y - 4)^2 = 25$	
(c)	2	C	CN	G11, G8	(i) $y = 9$, (ii) $T(-9, 9)$	

- ¹ ss: know to use midpoint
- ² pd: process gradient of PQ
- ³ ss: know how to find perp. gradient
- ⁴ ic: state equ. of line
- ⁵ ic: interpret "parallel to y -axis"
- ⁶ pd: process radius
- ⁷ ic: state equ. of circle
- ⁸ ic: interpret diagram
- ⁹ ss: know to use equ. of AB

- ¹ midpoint = $(-1, 5)$
- ² $m_{PQ} = \frac{9-1}{1-(-1)}$
- ³ $m_{\perp} = -\frac{1}{2}$
- ⁴ $y - 5 = -\frac{1}{2}(x - (-1))$
- ⁵ $y_C = 4$ stated or implied by •⁷
- ⁶ radius = 5 or equiv. stated or implied by •⁷
- ⁷ $(x - 1)^2 + (y - 4)^2 = 25$
- ⁸ $y = 9$
- ⁹ $T = (-9, 9)$

[SQA] 3. $f(x) = 3 - x$ and $g(x) = \frac{3}{x}, x \neq 0$.

(a) Find $p(x)$ where $p(x) = f(g(x))$.

2

(b) If $q(x) = \frac{3}{3-x}, x \neq 3$, find $p(q(x))$ in its simplest form.

3

Part	Marks	Level	Calc.	Content	Answer	U1 OC2
(a)	2	C	CN	A4	$3 - \frac{3}{x}$	2000 P2 Q3
(b)	2	C	CN	A4	x	
(b)	1	A/B	CN	A4		

•¹ ic: interpret composite func.

•² pd: process

•³ ic: interpret composite func.

•⁴ pd: process

•⁵ pd: process

•¹ $f\left(\frac{3}{x}\right)$ stated or implied by •²

•² $3 - \frac{3}{x}$

•³ $p\left(\frac{3}{3-x}\right)$ stated or implied by •⁴

•⁴ $3 - \frac{3}{\frac{3}{3-x}}$

•⁵ x


[SQA] 4. The parabola shown crosses the x -axis at $(0,0)$ and $(4,0)$, and has a maximum at $(2,4)$.

The shaded area is bounded by the parabola, the x -axis and the lines $x = 2$ and $x = k$.

(a) Find the equation of the parabola.

(b) Hence show that the shaded area, A , is given by

$$A = -\frac{1}{3}k^3 + 2k^2 - \frac{16}{3}.$$



2

Part	Marks	Level	Calc.	Content	Answer	U2 OC2
(a)	2	C	CN	A19	$y = 4x - x^2$	2000 P2 Q4
(b)	3	C	CN	C16	proof	

•¹ ic: state standard form

•² pd: process for x^2 coeff.

•³ ss: know to integrate

•⁴ pd: integrate correctly

•⁵ pd: process limits and complete proof

•¹ $ax(x - 4)$

•² $a = -1$

•³ $\int_2^k$ (function from (a))

•⁴ $-\frac{1}{3}x^3 + 2x^2$

•⁵ $-\frac{1}{3}k^3 + 2k^2 - \left(-\frac{8}{3} + 8\right)$

[SQA] 5. Solve the equation $3 \cos 2x^\circ + \cos x^\circ = -1$ in the interval $0 \leq x \leq 360$.

5

Part	Marks	Level	Calc.	Content	Answer	U2 OC3
	5	A/B	CR	T10	60, 131.8, 228.2, 300	2000 P2 Q5
				•¹ ss: know to use $\cos 2x = 2 \cos^2 x - 1$ •² pd: process •³ ss: know to/and factorise quadratic •⁴ pd: process •⁵ pd: process	•¹ $3(2 \cos^2 x^\circ - 1)$ •² $6 \cos^2 x^\circ + \cos x^\circ - 2 = 0$ •³ $(2 \cos x^\circ - 1)(3 \cos x^\circ + 2)$ •⁴ $\cos x^\circ = \frac{1}{2}, x = 60, 30$ •⁵ $\cos x^\circ = -\frac{2}{3}, x = 132, 228$	

[SQA] 6. A goldsmith has built up a solid which consists of a triangular prism of fixed volume with a regular tetrahedron at each end.

The surface area, A , of the solid is given by

$$A(x) = \frac{3\sqrt{3}}{2} \left(x^2 + \frac{16}{x} \right)$$

where x is the length of each edge of the tetrahedron.Find the value of x which the goldsmith should use to minimise the amount of gold plating required to cover the solid.

6

Part	Marks	Level	Calc.	Content	Answer	U1 OC3								
	6	A/B	CN	C11	$x = 2$	2000 P2 Q6								
				•¹ ss: know to differentiate •² pd: process •³ ss: know to set $f'(x) = 0$ •⁴ pd: deal with x^{-2} •⁵ pd: process •⁶ ic: check for minimum	•¹ $A'(x) = \dots$ •² $\frac{3\sqrt{3}}{2}(2x - 16x^{-2})$ or $3\sqrt{3}x - 24\sqrt{3}x^{-2}$ •³ $A'(x) = 0$ •⁴ $-\frac{16}{x^2}$ or $-\frac{24\sqrt{3}}{x^2}$ •⁵ $x = 2$ •⁶ <table><tr><td>x</td><td>2^-</td><td>2</td><td>2^+</td></tr><tr><td>$A'(x)$</td><td>$-ve$</td><td>0</td><td>$+ve$</td></tr></table> so $x = 2$ is min.		x	2^-	2	2^+	$A'(x)$	$-ve$	0	$+ve$
x	2^-	2	2^+											
$A'(x)$	$-ve$	0	$+ve$											

- [SQA] 7. For what value of t are the vectors $\mathbf{u} = \begin{pmatrix} t \\ -2 \\ 3 \end{pmatrix}$ and $\mathbf{v} = \begin{pmatrix} 2 \\ 10 \\ t \end{pmatrix}$ perpendicular? 2


Part	Marks	Level	Calc.	Content	Answer	U3 OC1
	2	C	CN	G27	$t = 4$	2000 P2 Q7
•¹ ss: know to use scalar product •² ic: interpret scalar product				•¹ $\mathbf{u} \cdot \mathbf{v} = 2t - 20 + 3t$ •² $\mathbf{u} \cdot \mathbf{v} = 0 \Rightarrow t = 4$		

- [SQA] 8. Given that $f(x) = (5x - 4)^{\frac{1}{2}}$, evaluate $f'(4)$. 3

Part	Marks	Level	Calc.	Content	Answer	U3 OC2
	1	C	CN	C21	$\frac{5}{8}$	2000 P2 Q8
	2	A/B	CN	C21		
•¹ pd: differentiate power •² pd: differentiate 2nd function •³ pd: evaluate $f'(x)$					•¹ $\frac{1}{2}(5x - 4)^{-\frac{1}{2}}$ •² $\times 5$ •³ $f'(4) = \frac{5}{8}$	

- [SQA] 9. A cuboid measuring 11 cm by 5 cm by 7 cm is placed centrally on top of another cuboid measuring 17 cm by 9 cm by 8 cm.

Coordinates axes are taken as shown.



- (a) The point A has coordinates $(0, 9, 8)$ and C has coordinates $(17, 0, 8)$.

Write down the coordinates of B.

1

- (b) Calculate the size of angle ABC.

6

Part	Marks	Level	Calc.	Content	Answer	U3 OC1
(a)	1	C	CN	G22	$B(3, 2, 15)$	2000 P2 Q9
(b)	6	C	CR	G28	92.5°	

- ¹ ic: interpret 3-d representation
- ² ss: know to use scalar product
- ³ pd: process vectors
- ⁴ pd: process vectors
- ⁵ pd: process lengths
- ⁶ pd: process scalar product
- ⁷ pd: evaluate scalar product

- ¹ $B = (3, 2, 15)$ treat $\begin{pmatrix} 3 \\ 2 \\ 15 \end{pmatrix}$ as bad form
- ² $\cos \widehat{ABC} = \frac{\vec{BA} \cdot \vec{BC}}{|\vec{BA}| |\vec{BC}|}$
- ³ $\vec{BA} = \begin{pmatrix} -3 \\ 7 \\ -7 \end{pmatrix}$
- ⁴ $\vec{BC} = \begin{pmatrix} 14 \\ -2 \\ -7 \end{pmatrix}$
- ⁵ $|\vec{BA}| = \sqrt{107}, |\vec{BC}| = \sqrt{249}$
- ⁶ $\vec{BA} \cdot \vec{BC} = -7$
- ⁷ $\widehat{ABC} = 92.5^\circ$


[SQA] 10. Find $\int \frac{1}{(7-3x)^2} dx$.

2

Part	Marks	Level	Calc.	Content	Answer	U3 OC2
	2	A/B	CN	C22, C14	$\frac{1}{3(7-3x)} + c$	2000 P2 Q10
•¹ pd: integrate function •² pd: deal with function of function					•¹ $\frac{1}{-1}(7-3x)^{-1}$ •² $\times \frac{1}{-3}$	

[SQA] 11. The results of an experiment give rise to the graph shown.

- (a) Write down the equation of the line in terms of P and Q .



2

It is given that $P = \log_e p$ and $Q = \log_e q$.

- (b) Show that p and q satisfy a relationship of the form $p = aq^b$, stating the values of a and b .

4

Part	Marks	Level	Calc.	Content	Answer	U3 OC3
(a)	2	A/B	CR	G3	$P = 0.6Q + 1.8$	2000 P2 Q11
(b)	4	A/B	CR	A33	$a = 6.05, b = 0.6$	

- ¹ ic: interpret gradient
- ² ic: state equ. of line
- ³ ic: interpret straight line
- ⁴ ss: know how to deal with x of $x \log y$
- ⁵ ss: know how to express number as log
- ⁶ ic: interpret sum of two logs

- ¹ $m = \frac{1.8}{3} = 0.6$
- ² $P = 0.6Q + 1.8$

Method 1

- ³ $\log_e p = 0.6 \log_e q + 1.8$
- ⁴ $\log_e q^{0.6}$
- ⁵ $\log_e 6.05$
- ⁶ $p = 6.05q^{0.6}$

Method 2

$$\ln p = \ln aq^b$$

- ³ $\ln p = \ln a + b \ln q$
- ⁴ $\ln p = 0.6 \ln q + 1.8$ *stated or implied by •⁵ or •⁶*
- ⁵ $\ln a = 1.8$
- ⁶ $a = 6.05, b = 0.6$

[END OF QUESTIONS]