

# 2014 Mathematics Paper 2 National 5

Finalised Marking Instructions

#### © Scottish Qualifications Authority 2014

The information in this publication may be reproduced to support SQA qualifications only on a non-commercial basis. If it is to be used for any other purposes written permission must be obtained from SQA's NQ Assessment team.

Where the publication includes materials from sources other than SQA (secondary copyright), this material should only be reproduced for the purposes of examination or assessment. If it needs to be reproduced for any other purpose it is the centre's responsibility to obtain the necessary copyright clearance. SQA's NQ Assessment team may be able to direct you to the secondary sources.

These Marking Instructions have been prepared by Examination Teams for use by SQA Appointed Markers when marking External Course Assessments. This publication must not be reproduced for commercial or trade purposes.



#### General Marking Principles for National 5 Mathematics

This information is provided to help you understand the general principles you must apply when marking candidate responses to questions in this Paper. These principles must be read in conjunction with the detailed marking instructions, which identify the key features required in candidate responses.

- (a) Marks for each candidate response must always be assigned in line with these General Marking Principles and the Detailed Marking Instructions for this assessment.
- (b) Marking should always be positive. This means that, for each candidate response, marks are accumulated for the demonstration of relevant skills, knowledge and understanding: they are not deducted from a maximum on the basis of errors or omissions.
- (c) Credit must be assigned in accordance with the specific assessment guidelines.
- (d) Candidates may use any mathematically correct method to answer questions except in cases where a particular method is specified or excluded.
- (e) Working subsequent to an error must be followed through, with possible credit for the subsequent working, provided that the level of difficulty involved is approximately similar. Where, subsequent to an error, the working is easier, candidates lose the opportunity to gain credit.
- (f) Where transcription errors occur, candidates would normally lose the opportunity to gain a processing mark.
- (g) Scored out working which has not been replaced should be marked where still legible. However, if the scored out working has been replaced, only the work which has not been scored out should be marked.
- (h) Where a candidate has made multiple attempts, mark all attempts and award the lowest mark.
- (i) Unless specifically mentioned in the specific assessment guidelines, do not penalise:
  - Working subsequent to a correct answer
  - Correct working in the wrong part of a question
  - Legitimate variations in solutions
  - Bad form
  - Repeated error within a question

# **Detailed Marking Instructions for each question**

Question		Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •	
1.		<ul> <li>Ans: 590</li> <li>1 know how to decrease by 15%</li> <li>2 know how to calculate roll</li> <li>3 carry out calculations correctly within a valid strategy and round to the nearest ten</li> </ul>	3	•¹ ×0·85 •² 964×0·85³ •³ 590	
Notes:			l		
1. For an	answ	er of 590 without working		award 3/3	<b>///</b>
2. For an	ansv	ver of 592 or 592·0165 without wo	rking	award 2/3	√√x
		ncorrect percentage has been used possibility of awarding 2/3	d, the work	ring must be followed through	x√√
4. For an	<b>4.</b> For an answer of 2460 $(964 \times 0.85 \times 3)$ with working, award 1/3				
5. For an answer of 530 $(964-964\times0.15\times3)$ with working, award 1/3				√xx	
6. For an	<b>6.</b> For an answer of 430 $(964 \times 0.15 \times 3)$			award 0/3	xxx

Question		Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
2.		Ans: B (8, 4, 10), C (4, 0, 10)	2	
		• 1 state coordinates of B		• 1 (8, 4, 10)
		• 2 state coordinates of C		• <sup>2</sup> (4, 0, 10)

# Notes:

award 1/2 ×√

- For eg B(8, 4, 9) leading to C(4, 0, 9)
   The maximum mark available is 1/2 where
  - (a) brackets are omitted
  - (b) answers are given in component form

Que	Question		Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
3.	(a)		Ans: $5a + 3c = 158.25$ • 1 construct equation	1	$\bullet^1 5a + 3c = 158.25$
Note	es:	I		l .	
,	<b>1.</b> Ac	cept	variables other than $a$ and $c$ .		
	(b)		Ans: $3a + 2c = 98$	1	
			•¹ construct equation		$\bullet^1 \ 3a + 2c = 98$
Note	es:				
	(c)		Ans: Adult ticket costs £22.50 Child ticket costs £15.25	4	
			• 1 evidence of scaling		
			$ullet^2$ follow a valid strategy through to produce values for $a$ and $c$		$ullet^2$ values for $a$ and $c$
			$ullet^3$ calculate correct values for $a$ and $c$		• $^{3}$ $a = 22.5$ and $c = 15.25$
			• 4 communicate answers in money		• <sup>4</sup> Adult £22·50 Child £15·25

1. The fourth mark may only be awarded when **all** of the following are given in the final answer: the words "adult" and "child", the £ signs and **both** amounts written with two decimal figures.

Question			Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
4.	(a)	(i)	Ans: $\overline{x} = 56.5$	1	
			•¹ calculate mean		$\bullet^1 \ x = 56.5$

1. Do not accept 56.5 rounded to 57.

(ii)	Ans: $s = 2.4$	3	
	• 1 calculate $(x-\bar{x})^2$		•¹ 0·25, 0·25, 2·25, 2·25, 12·25, 12·25
	•² substitute into formula		$\bullet^2 \sqrt{\frac{29 \cdot 5}{5}}$
	• 3 calculate standard deviation		•³ 2·4(2)

#### Notes:

1. For use of alternative formula, award marks as follows:

• 1 calculate  $\sum x$  and  $\sum x^2$ 

•¹ 339, 19183

• <sup>2</sup> substitute into formula

 $\int_{0}^{2} \sqrt{\frac{19183 - \frac{339^{2}}{6}}{5}}$ 

• 3 calculate standard deviation

• <sup>3</sup> 2·4(2....)

2. For correct answer without working

award 0/3

(b)	Ans: No, standard deviation is greater OR No, times are more spread out	1	
	• <sup>1</sup> no, with valid explanation		• <sup>1</sup> e.g. No, standard deviation is greater

#### Notes:

- 1. Answer must be consistent with answer to part (a)(ii).
- **2.** Accept "No, as 3.2 > 2.4"
- 3. Only award the mark if it is clear that the reason is based on standard deviation only.
- 4. Do not accept "No, times are less consistent" without further explanation.

Question		Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
5.		Ans: 3072 cm <sup>3</sup>	3	
		• ¹ state linear scale factor		• $\frac{24}{15}$ or 1.6
		•² state volume scale factor		
		• 3 calculate volume (calculation must involve a power of the scale factor) and state correct units		• <sup>3</sup> 3072cm <sup>3</sup>

Some common answers

(b) 1920cm<sup>3</sup> (
$$\left(\frac{24}{15}\right)^2 \times 750$$
) award 2/3  $\checkmark \times \checkmark$ 

(c)1200cm<sup>3</sup> (
$$\left(\frac{24}{15}\right) \times 750$$
)

(d)675000000cm<sup>3</sup> (
$$\left(\frac{24}{15}\right) \times 750^3$$
) award 1/3  $\checkmark \times \times$   
(e)183cm<sup>3</sup> ( $\left(\frac{15}{24}\right)^3 \times 750$ ) award 2/3  $\times \checkmark \checkmark$   
(f) 933cm<sup>3</sup> ( $\left(\frac{15}{24}\right)^3 \times 750 + 750$ ) award 2/3  $\times \checkmark \checkmark$ 

(e)183cm<sup>3</sup> 
$$\left(\left(\frac{15}{24}\right)^3 \times 750\right)$$

(f) 933cm<sup>3</sup> 
$$\left(\left(\frac{15}{24}\right)^3 \times 750 + 750\right)$$

- The third mark is not available where premature rounding leads to an incorrect answer. eg  $4.1 \times 750 = 3075 \text{cm}^3$ award 2/3 ✓√×
- 3. Alternative Method
  - 1 know how to find radius of smaller cylinder

$$\bullet^1 \quad \sqrt{\frac{750}{15\pi}}$$

• know how to find radius of larger cylinder

$$\bullet^2 \left(\frac{24}{15}\right) \times \sqrt{\frac{750}{15\pi}}$$

• 3 calculate volume and state correct units

• 3072cm 3

Question	Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
6.	Ans: no, with valid reason.	4	
	• 1 valid strategy		• 1 use Converse of Pythagoras' Theorem eg 110 <sup>2</sup> and 85 <sup>2</sup> + 75 <sup>2</sup>
	•² evaluation		•² 12 100 and 12 850
	• <sup>3</sup> comparison		• 3 e.g. $110^2 \neq 75^2 + 85^2$
	• 4 valid conclusion		• 4 No, since not right angled

1. For alternative methods, award marks as follows:

(i) • 1 valid strategy

• 1 use Pythagoras' Theorem  $eg 85^2 + 75^2$ 

• <sup>2</sup> evaluation

 $\bullet^2 h = 113.36$ 

• 3 comparison

•  $^{3}$  e.g.  $113 \cdot 36 > 110$ 

• 4 valid conclusion

• 4 No, since not right angled

(ii) •¹ valid strategy

• 1 substitute correctly into cosine

$$eg \ \frac{85^2 + 75^2 - 110^2}{2 \times 85 \times 75}$$

• <sup>2</sup> evaluation

• 2 86·6°

• 3 comparison

 $\bullet^3 86.6^{\circ} < 90^{\circ}$ 

• 4 valid conclusion

• 4 No, since not right angled

There must be an explicit comparison for the award of the third mark. 2.

eg 
$$\sqrt{85^2 + 75^2} = 113.36$$
.

No, since not right angled.

award 3/4 ✓✓×✓

3. Conclusion must involve reference to "not a right angle".

eg 
$$110^2 = 85^2 + 75^2 \rightarrow 12100 \neq 12850$$
.

No, Hightown is not due north of Lowtown. award  $3/4 \checkmark \checkmark \checkmark \times$ 

- The final mark is not available where the candidate's only conclusion is an invalid statement involving the word bearing.
  - eg "No, Hightown is on a bearing of 87° from Lowtown, not 90°"

Que	stion	Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
7.		Ans: 150 cm <sup>3</sup>	5	awaranig a mark ac caen
		• ¹ substitute correctly into formula for volume of cone		$\bullet^{1} \frac{1}{3} \times \pi \times 4^{2} \times 15 \ (=251 \cdot 32)$
		• <sup>2</sup> substitute correctly into formula for volume of sphere <b>or</b> hemisphere		$ \begin{array}{cccccccccccccccccccccccccccccccccccc$
		• 3 know to subtract volume of hemisphere from volume of cone		•³ evidence
		<ul> <li>4 carry out all calculations correctly (must involve difference or sum of two volume calculations)</li> </ul>		• <sup>4</sup> 145·24
		• <sup>5</sup> round <b>final answer</b> to 2 significant figures		• <sup>5</sup> 150 (cm <sup>3</sup> )

- **1.** Accept variations in  $\pi$ .
- 2. Some common answers (working must be shown):

(i) 
$$39 \left(\frac{1}{3} \times \pi \times 4^2 \times 15 - \frac{4}{3} \times \pi \times 3 \cdot 7^3\right) \qquad \text{award } 4/5 \quad \checkmark \checkmark \checkmark \checkmark$$

(ii) 120 
$$(\frac{1}{3} \times \pi \times 4^2 \times 15 - \frac{1}{2} \times \frac{4}{3} \times \pi \times 4^3)$$
 award 4/5  $\checkmark \times \checkmark \checkmark$ 

(iii) 110 
$$(\frac{1}{3} \times \pi \times 3 \cdot 7^2 \times 15 - \frac{1}{2} \times \frac{4}{3} \times \pi \times 3 \cdot 7^3)$$
 award 4/5  $\times \checkmark \checkmark \checkmark \checkmark$ 

(iv) 160 
$$(\frac{1}{3} \times \pi \times 8^2 \times 15 - \frac{1}{2} \times \frac{4}{3} \times \pi \times 7 \cdot 4^3)$$
 award 4/5  $\times \checkmark \checkmark \checkmark$ 

(v) 360 
$$(\frac{1}{3} \times \pi \times 4^2 \times 15 + \frac{1}{2} \times \frac{4}{3} \times \pi \times 3 \cdot 7^3)$$
 award 4/5  $\checkmark \checkmark \checkmark \checkmark \checkmark$ 

(vi) 460 
$$(\frac{1}{3} \times \pi \times 4^2 \times 15 + \frac{4}{3} \times \pi \times 3 \cdot 7^3)$$
 award 4/5  $\checkmark \checkmark \checkmark \checkmark \checkmark$ 

(vii) 80 
$$(\frac{1}{3} \times \pi \times 3.7^2 \times 15 - \frac{1}{2} \times \frac{4}{3} \times \pi \times 4^3)$$
 award 3/5  $\times \times \checkmark \checkmark \checkmark$ 

(viii) 250 
$$(\frac{1}{3} \times \pi \times 4^2 \times 15)$$
 award 2/5  $\checkmark \times \times \times \checkmark$ 

3. The final mark is only available where answers to all intermediate steps involve at least three significant figures.

eg 
$$251 \cdot 32 - 106 \cdot 08 = 250 - 110 = 140$$
 award  $4/5 \checkmark \checkmark \checkmark \checkmark \checkmark$ 

Question		Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
8.		Ans: $5n^4$	3	
		•¹ simplify powers in numerator		•¹ 10 <i>n</i> <sup>6</sup>
		• <sup>2</sup> cancel constants		$\bullet^2 \frac{5n^6}{n^2}$
		• $^3$ eliminate $n$ from denominator		$\bullet^3 5n^4$

1. For 
$$5n^4$$
 without working award 3/3

2. For a final answer of 
$$\frac{5n^4}{1}$$
 award 2/3  $\checkmark \times \checkmark$ 

3. For an answer of 
$$5n^3$$

(a) (i) 
$$\frac{10n^5}{2n^2} = 5n^3$$
 award 2/3 ×  $\checkmark$ 

(ii) 
$$\frac{10n^6}{2n^2} = 5n^3$$
 award 2/3  $\checkmark\checkmark$ ×

(b) (i) 
$$\frac{n^4 \times 10}{2n} = \frac{n^4 \times 10}{2} = 5n^3$$
 award 1/3  $\checkmark \times \checkmark$ 

(ii)  $5n^3$  without working award 1/3

Question	Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
9.	Ans: $\frac{4x-15}{x(x+5)}$ • 1 correct common denominator (or correct numerator)  • 2 consistent numerator (or denominator)  • 3 simplify	3	• $^{1}x(x+5)$ or $7x-3(x+5)$ • $^{2}\frac{7x-3(x+5)}{x(x+5)}$ • $^{3}\frac{4x-15}{x(x+5)}$

#### Notos

1. Correct answer without working

award 3/3

2. For 
$$\frac{7x}{x(x+5)} - \frac{3(x+5)}{x(x+5)}$$

3. For subsequent incorrect working, the final mark is not available.

Que	stion	Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
10.	(a)	• 1 substitute correctly into cosine rule  • 2 calculate cos B correctly  • 3 calculate angle ABC correctly	3	$ cos B = \frac{8^2 + 11^2 - 13^2}{2 \times 8 \times 11} $ • cos B = 0.09 • 85 or 84.8
N				

- For 1.48 (uses RAD) or 94.2 (uses GRAD), with working award 3/3
- The  $2^{nd}$  mark can be awarded for  $\cos^{-1}\left(\frac{16}{176}\right)$

(b)	Ans: 155·2°	2	
	• 1 know how to calculate the angle		•1 360 – 120 – [answer to (a)] or equivalent
	•² correctly calculate the angle within a valid strategy		• <sup>2</sup> 155·2
Notes:			

Question	Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
11.	$Ans: a = \frac{2(s - ut)}{t^2}$	3	
	• ¹ subtract <i>ut</i>		$\bullet^1 s - ut = \frac{1}{2}at^2$
	• <sup>2</sup> multiply by 2		$ \bullet^2 \ 2(s-ut) = at^2 $
	$\bullet$ <sup>3</sup> divide by $t^2$		$\bullet^3 a = \frac{2(s-ut)}{t^2}$
N		1	

# Notes:

Correct answer without working 1.

award 3/3

For subsequent incorrect working, the final mark is not available. 2.

$$a = \frac{s - ut}{\frac{1}{2}t^2}$$

award 2/3

Question	Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
12.	Ans: $x^{\circ} = 63^{\circ}$ , 297°	3	_
	• 1 rearrange equation		$\bullet^1 \cos x = \frac{5}{11}$
	• $^2$ find one value of $x$		• $^{2}$ $x = 63$
	• $^3$ find another value of $x$		$\bullet^3 x = 297$
Notes			

- 1.
- The  $2^{nd}$  angle must be consistent with the first angle. For x = 1.1,358.9 (uses RAD), award 3/3 (with working), award 2/3 (without working) 2.
- For x = 70, 290 (uses GRAD), award 3/3 (with working), award 2/3 (without working) 3.

Question	Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
13.	Ans: 151·3 m²  •1 know how to find area of segment	5	•¹ evidence of e.g. major sector + triangle or circle — minor sector + triangle
	•² know to express sector as a fraction of a circle		• $\frac{310}{360}$ or $\frac{50}{360}$
	•3 know how to find area of sector		•3 $\frac{310}{360} \times \pi \times 7^2$ (= 132·56) or $\frac{50}{360} \times \pi \times 7^2$ (= 21·38)
	• 4 know how to calculate area of triangle		$\bullet^4  \frac{1}{2} \times 7 \times 7 \times \sin 50  (=18.77)$
	• 5 carry out all calculations correctly within a valid strategy		• <sup>5</sup> 151·3 m <sup>2</sup>

Notes for question 13 are on next page.

Question	Expected Answer(s) Give one mark for each •	Max Mark	Illustrations of evidence for awarding a mark at each •
Notes:			

- 1. Accept variations in  $\pi$ ; disregard premature or incorrect rounding of  $\frac{310}{360}$  or  $\frac{50}{360}$
- 2. Use of RAD or GRAD (working must be shown)
  - (a) For 149.9 [uses GRAD]

award 5/5

(b) Where the use of RAD leads to an answer of  $126 \cdot 1(-6 \cdot 43 + 132 \cdot 56)$ or 139·0(6·43+132·56)

award 4/5

3. Some common answers (working must be shown):

$$56.6 \left(\frac{310}{360} \times \pi \times 14 + \frac{1}{2} \times 7 \times 7 \times \sin 50^{\circ}\right) \qquad \text{award } 4/5 \quad \checkmark \checkmark \checkmark \checkmark$$

$$40.1 \left(\frac{50}{360} \times \pi \times 7^{2} + \frac{1}{2} \times 7 \times 7 \times \sin 50^{\circ}\right) \qquad \text{award } 4/5 \quad \times \checkmark \checkmark \checkmark$$

$$2.6 \left(\frac{50}{360} \times \pi \times 7^{2} - \frac{1}{2} \times 7 \times 7 \times \sin 50^{\circ}\right) \qquad \text{award } 4/5 \quad \times \checkmark \checkmark \checkmark$$

$$24.9 \left(\frac{50}{360} \times \pi \times 14 + \frac{1}{2} \times 7 \times 7 \times \sin 50^{\circ}\right) \qquad \text{award } 3/5 \quad \times \checkmark \times \checkmark \checkmark$$

$$132.6 \left(\frac{310}{360} \times \pi \times 7^{2}\right) \qquad \text{award } 2/5 \quad \times \checkmark \checkmark \times$$

$$21.4 \left(\frac{50}{360} \times \pi \times 7^{2}\right) \qquad \text{award } 2/5 \quad \times \checkmark \checkmark \times$$

$$18.8 \left(\frac{1}{2} \times 7 \times 7 \times \sin 50^{\circ}\right) \qquad \text{award } 1/5 \quad \times \times \times \checkmark \times$$

$$153.9 \left(\pi \times 7^{2}\right) \qquad \text{award } 0/5$$

4. The fifth mark is only available when the area of triangle MON is calculated using trigonometry.

## [END OF MARKING INSTRUCTIONS]

award 0/5