


21. Newman-Casey PA, Stem M, Talwar N, et al. Risk factors associated with developing branch retinal vein occlusion among enrollees in a united states managed care plan. *Ophthalmology*. 2014;121:1939–1948.
22. Sperduto RD, Hiller R, Chew E, et al. Risk factors for hemiretinal vein occlusion: comparison with risk factors for central and branch retinal vein occlusion: the eye disease case-control study. *Ophthalmology*. 1998;105:765–771.
23. Prisco D, Marcucci R. Retinal vein thrombosis: risk factors, pathogenesis and therapeutic approach. *Pathophysiol Haemost Thromb*. 2002;32:308–311.
24. Lim DH, Shin KY, Han K, et al. Differential effect of the metabolic syndrome on the incidence of retinal vein occlusion in the Korean population: a nationwide cohort study. *Transl Vis Sci Technol*. 2020;9:15.
25. Hernán MA, Robins JM. *Causal Inference: What If*. CRC Press; 2020.
26. Zhou JQ, Xu L, Wang S, et al. The 10-year incidence and risk factors of retinal vein occlusion: the Beijing eye study. *Ophthalmology*. 2013;120:803–808.
27. Cugati S, Wang JJ, Rochtchina E, Mitchell P. Ten-year incidence of retinal vein occlusion in an older population: the Blue Mountains Eye Study. *Arch Ophthalmol*. 2006;124:726–732.
28. Arakawa S, Yasuda M, Nagata M, et al. Nine-year incidence and risk factors for retinal vein occlusion in a general Japanese population: the Hisayama Study. *Invest Ophthalmol Vis Sci*. 2011;52:5905–5909.
29. Devers MC, Campbell S, Simmons D. Influence of age on the prevalence and components of the metabolic syndrome and the association with cardiovascular disease. *BMJ Open Diabetes Res Care*. 2016;4:e000195.

Pictures & Perspectives



X-Ray Vision during Iodine-125 Plaque Radiation Treatment of a Choroidal Melanoma

Iodine-125 brachytherapy was initiated in a 53-year-old man with a choroidal melanoma. The patient's background as a Hazmat radiation exposure instructor allowed him to measure the exposure capabilities of his radioactive plaque. Based on his understanding of radioactive decay and waste gamma radiation, he used the inverse square law to determine a convergent hotspot anterior to the orbit, just lateral to the nose (A). Using the PD-10i dosimeter (SAIC), he determined that the exposure needed for a hand plate x-ray scan would be 10 to 15 millirem, achievable with 10 minutes of unprotected exposure. He performed a hand x-ray scan using these parameters (B). Magnified version of Fig A-B is available online at www.opthalmologyretina.org.

KEVIN R. CARD, BS
DANTE J. PIERAMICI, MD

California Retina Consultants and Research Foundation, Santa Barbara, California