

Ophthalmic Images

Amelanotic and Pigmented Uveal Melanoma

Andreia Margarida Mendes Silva, MD; Maria Cristina Ferrão Fonseca, MD; Rui Daniel Proença, MD, PhD




Figure. Fundus photograph of the left eye at presentation showing a bilobed, elevated lesion with pigmented and nonpigmented portions.

A 48-year-old white woman presented with a sudden, painless decrease in the vision of her left eye. Her visual acuity was 20/100. Results of an anterior segment evaluation were normal. Fundus examination revealed a bilobed, elevated lesion nasal to the optic disc with an amelanotic portion and a darkly pigmented one (**Figure**). Ultrasonography revealed a mass consistent with choroidal melanoma with a basal diameter of 17.5 mm and thick-

ness of 7.97 mm. A systemic workup revealed no evidence of metastatic disease. She underwent episcleral brachytherapy with an iodine 125 plaque. Brachytherapy was followed by tumor regression documented by ultrasonography 3 months after the treatment. After the first year, she maintained the visual acuity noted before treatment, with regression more readily apparent within the amelanotic portion of the lesion.

ARTICLE INFORMATION

Author Affiliations: Department of Ophthalmology, Centro de Responsabilidade Integrado em Oftalmologia do Centro Hospitalar e Universitário de Coimbra, Coimbra, Portugal (Silva, Fonseca, Proença); Faculty of Medicine, University of Coimbra, Coimbra, Portugal (Fonseca, Proença).

Corresponding Author: Andreia Margarida Mendes Silva, MD, Department of Ophthalmology, Centro Hospitalar e Universitário de Coimbra, Praceta Prof Mota Pinto, 3000-075 Coimbra, Portugal (rproenca@chuc.min-saude.pt).

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

Additional Contributions: We thank the patient for granting permission to publish this information.