

S.G.: Consultant — Genentech, Ohr, Regeneron, SciFluor, Ophthotech, Allergan

HUMAN SUBJECTS: Human subjects were included in this study. No animal subjects were used in this study. The study was approved by a local institutional review board (IRBco). The study was performed in accordance with the tenets of the Declaration of Helsinki.

Author Contributions:

Conception and design: Adrean, Chaili, Ramkumar, Pirouz, Grant

Analysis and interpretation: Adrean, Chaili, Ramkumar, Pirouz, Grant

Data collection: Adrean, Chaili

Obtained funding: None

Overall responsibility: Adrean, Chaili, Ramkumar, Pirouz, Grant

Abbreviations and Acronyms:


ANCHOR = Anti-VEGF Antibody for the Treatment of Predominantly Classic Choroidal Neovascularization in Age-Related Macular Degeneration; **AMD** = age-related macular degeneration; **CATT** = Comparison of

Age-Related Macular Degeneration Treatments Trial; **CNV** = choroidal neovascularization; **ETDRS** = Early Treatment Diabetic Retinopathy Study; **GA** = geographic atrophy; **HORIZON** = Open-Label Extension Trial of Ranibizumab for Choroidal Neovascularization Secondary to Age-Related Macular Degeneration; **HIPAA** = Health Insurance Portability and Accountability Act; **MARINA** = Minimally Classic/Occult Trial of the Anti-VEGF Antibody Ranibizumab In the Treatment of Neovascular AMD; **nAMD** = neovascular age-related macular degeneration; **PRN** = pro re nata; **RCT** = randomized controlled trial; **RPE** = retinal pigment epithelium; **TES** = treat-extend-stop; **VA** = visual acuity; **VEGF** = vascular endothelial growth factor.

Correspondence:

Sean D. Adrean, MD, Retina Consultants of Orange County, 301 West Bastanchury Avenue, No. 285, Fullerton, CA 92835. E-mail: seadrean@yahoo.com.

Pictures & Perspectives



Choroidal Melanoma with Overlying Lipofuscin Sediment

A 56-year-old woman had a pigmented choroidal lesion with overlying orange pigment sedimentation (Fig 1A) resembling the pseudohypopyon seen in Best vitelliform macular dystrophy. Corresponding hyperautofluorescence on fundus autofluorescence was seen (Fig 1B) and B-scan ultrasonography documented a 2.2-mm thick, dome-shaped hollow lesion (Fig 1C). Cytology (original magnification, 1000×) direct smear with Papanicolaou stain showed a moderately cellular aspirate of small cells with hyperchromic nuclei (Fig 1D). The patient elected to proceed with plaque brachytherapy and fine-needle aspiration biopsy (FNAB) revealed a class 1A gene-expression profile. To the best of our knowledge, this is the second published case of a pseudohypopyon overlying a choroidal melanoma. (Magnified version of Fig 1A-D is available online at www.aaojournal.org).

KELSEY ROELOFS, MD¹

LAKSHIMA PUTTAGUNTA, MD²

EZEKIEL WEIS, MD, MPH^{1,3}

¹Department of Ophthalmology and Visual Sciences, University of Alberta, Edmonton, Alberta, Canada; ²Department of Laboratory Medicine and Pathology, University of Alberta, Edmonton, Alberta, Canada; ³Department of Surgery, University of Calgary, Edmonton, Alberta, Canada