

Supported by the National Eye Institute, National Institutes of Health, Bethesda, Maryland (cooperative agreement nos.: U10 EY017823, U10 EY017825, U10 EY017826, and U10 EY017828). The funding organization participated in the design and conduct of the study, data analysis and interpretation, and review of the manuscript.

Author Contributions:

Conception and design: Maguire, Martin, Ying, Jaffe, Daniel, Toth, Grunwald, Ferris, Fine

Analysis and interpretation: Maguire, Martin, Ying, Jaffe, Daniel, Toth, Grunwald, Ferris, Fine

Data collection: Maguire, Martin, Ying, Jaffe, Daniel, Toth, Grunwald, Ferris, Fine

Obtained funding: Maguire, Martin

Overall responsibility: Maguire, Martin, Ying, Jaffe, Daniel, Toth, Grunwald, Ferris, Fine


Abbreviations and Acronyms:

AMD = age-related macular degeneration; **CATT** = Comparison of Age-Related Macular Degeneration Treatments Trials; **OCT** = optical coherence tomography; **PRN** = pro re nata; **VA** = visual acuity; **VEGF** = vascular endothelial growth factor.

Correspondence:

Maureen G. Maguire, PhD, Department of Ophthalmology, University of Pennsylvania, 3535 Market Street, Suite 700, Philadelphia, PA 19104. E-mail: maguirem@mail.med.upenn.edu.

Pictures & Perspectives



Scleral Necrosis Simulating Recurrent Uveal Melanoma after Plaque Brachytherapy

A 59-year-old man presented with a pigmented subconjunctival lesion and multifocal scleral thinning (Fig 1A) overlying a choroidal melanoma 7 years after successful brachytherapy (85 Gray). B-scan ultrasonography showed regression of the intraocular tumor, with apparent extension through sclera (Fig 1B, arrow, S = sclera; C = cornea; I = iris). Biopsy revealed sclera infiltrated by pigmented cells (Fig 1C, star, H&E 10 $\times$), which proved to be melanophages (Fig 1D, star, CD68 10 $\times$) and chronic nongranulomatous inflammation. Melanoma cocktail stain was negative for viable tumor. Scleral necrosis is a rare late complication of plaque brachytherapy. Extraocular material may represent benign cells, not necessarily connoting tumor reactivation.

CHRISTOPHER K.H. BURRIS, MD

MARIA E. RODRIGUEZ, MD

HEATHER D. POTTER, MD

Department of Ophthalmology and Visual Sciences, University of Wisconsin — Madison, Madison, Wisconsin