




Conclusion:

Inter-rater and intra-rater agreements for target volumes were high, though substantial variability was observed for OAR delineation. No clear advantage of 7T MR versus 3T MR was found in terms of inter-rater agreement. However, treatment volumes delineated using 7T MR were generally smaller. Although 7T MR provides higher resolution, it did not show an improvement in delineation consistency.

Keywords: intra-rater variability, inter-rater variability

References:

(1) Sebastian Regnery, Benjamin R. Knowles, Daniel Paech, Nicolas Behl, Jan-Eric Meissner, Paul Windisch, Semi Ben Harrabi, Denise Bernhardt, Heinz-Peter Schlemmer, Mark E. Ladd, Stefan Rieken, Jürgen Debus, Sebastian Adeberg, High-resolution FLAIR MRI at 7 Tesla for treatment planning in glioblastoma patients, Radiotherapy and Oncology, Volume 130, 2019, Pages 180-184, ISSN 0167-8140, <https://doi.org/10.1016/j.radonc.2018.08.002>.

3191

Mini-Oral

Stereotactic radiosurgery for uveal melanoma – long-term outcomes and volumetric analysis

Jakub Cvek¹, Lukas Knybel¹, Andrea Soltysova¹, Eva Skacelikova¹, Jan Nemcansky²

¹Oncology, University Hospital Ostrava and Faculty of Medicine, Ostrava, Czech Republic. ²Ophtalmology, University Hospital Ostrava and Faculty of Medicine, Ostrava, Czech Republic

Purpose/Objective:

Our retrospective observational study aims to analyze tumor volume reduction, intraocular pressure, and visual acuity in long-term follow-up after stereotactic radiosurgery for uveal melanomas.

Material/Methods:

From 2010 to 2022, 192 uveal melanomas were treated with stereotactic radiosurgery in retrobulbar anesthesia and akinesia. The gross tumor volume (GTV), ciliary body, lens, and optic pathway were delineated after precise image registrations of CT and both T1 (with intravenous contrast enhancement) and T2 MRI datasets. No additional margins were added and a sequential optimization algorithm for one fraction was calculated. On-line image guidance based on the base of the skull and two orthogonal radiographs were provided. Prophylaxis was based on