

Metastatic Risk for Distinct Patterns of Postirradiation Local Recurrence of Posterior Uveal Melanoma

J. William Harbour, MD,^{1,2} Devron H. Char, MD,^{1,3} Stew Kroll, MS,¹ Jeanne M. Quivey, MD,³ Joseph Castro, MD³

Objective: The purpose of the study is to compare the prognostic significance of horizontal/marginal versus vertical/diffuse patterns of postirradiation local recurrence of posterior uveal melanoma.

Design: The study design was a nonrandomized, retrospective clinical study. Semi-parametric and nonparametric statistical techniques were used.

Participants: Seven hundred sixty-six posterior uveal melanoma patients were studied.

Intervention: Either iodine-125 plaque or helium ion radiation therapy was performed.

Main Outcome Measures: Local tumor recurrence and systemic metastasis were measured.

Results: Local tumor recurrence was detected in 66 (8.6%) of 766 irradiated tumors. The 5-year actuarial rate of local recurrence was 10%. The recurrence pattern was horizontal/marginal in 27 patients (41%) and vertical/diffuse in 39 patients (59%). Systemic metastasis was detected in 5 patients (19%) with horizontal/marginal recurrence and in 19 patients (49%) with vertical/diffuse recurrence. After known metastatic risk factors were controlled, the relative risk for metastasis was 2.2 for horizontal/marginal recurrence and 5.1 for vertical/diffuse recurrence ($P = 0.05$). The actuarial rate of systemic metastasis was 2.9% per year for all patients, 6.3% per year for patients with horizontal/marginal recurrence, and 15.5% per year for patients with vertical/diffuse recurrence.

Conclusions: Postirradiation local recurrence of posterior uveal melanoma is a risk factor for systemic metastasis. Vertical/diffuse recurrences may be associated more strongly with metastatic disease than horizontal/marginal recurrences.

Ophthalmology 1997; 104:1785–1793

Originally received: October 3, 1996.

Revision accepted: June 6, 1997.

¹ Department of Ophthalmology, Ocular Oncology Unit, University of California, San Francisco, California.

² Department of Ophthalmology and Visual Sciences, Washington University School of Medicine, and the Barnes Retina Service, St. Louis, Missouri.

³ Department of Radiation Oncology, University of California, San Francisco, California.

Presented in part at the American Academy of Ophthalmology Annual Meeting, Chicago, Illinois, October 1996.

Supported in part by a Heed-Knapp Ophthalmic Fellowship (JWH).

The authors have no proprietary interest in this study.

Address correspondence to J. William Harbour, MD, Washington University School of Medicine, Campus Box 8096, 660 South Euclid Ave, St Louis, MO 63110-1093.

Postirradiation local recurrence of posterior uveal melanoma is an occasional complication of both episcleral plaque brachytherapy and charged particle teletherapy. Local failure has been reported in 4% to 18% of patients treated with radioactive plaques¹⁻⁷ and 2% to 5% of patients treated with charged particle therapy (helium ions or protons).^{8,9} Several studies have reported an increased risk for systemic metastasis and melanoma-related mortality in patients with local recurrence.^{4,8,10} However, none of these reports distinguished between different morphologic patterns of local melanoma recurrence.

Local recurrences may present predominantly as a thin, horizontal extension along a tumor margin (horizontal/marginal recurrence) or as an area of focally or diffusely increased tumor thickness (vertical or diffuse recurrence).