


# Fixpoint of Function $f: D \rightarrow D$

- Let  $D = (S, \sqsubseteq)$ , and let  $f: D \rightarrow D$  be a function on this domain.
- A value  $x \in S$  is called a fixpoint of  $f$  if  $f(x) = x$ .


# Fixpoint of Function $f: D \rightarrow D$

- Let  $D = (S, \sqsubseteq)$ , and let  $f: D \rightarrow D$  be a function on this domain.
- A value  $x \in S$  is called a fixpoint of  $f$  if  $f(x) = x$ .
- For the powerset domain:




# Fixpoint of Function $f: D \rightarrow D$

- Let  $D = (S, \sqsubseteq)$ , and let  $f: D \rightarrow D$  be a function on this domain.
- A value  $x \in S$  is called a fixpoint of  $f$  if  $f(x) = x$ .
- For the powerset domain:
  - Every element of the domain is a fixpoint for the identity function  $I = \lambda x. x$ .




# Fixpoint of Function $f: D \rightarrow D$

- Let  $D = (S, \sqsubseteq)$ , and let  $f: D \rightarrow D$  be a function on this domain.
- A value  $x \in S$  is called a fixpoint of  $f$  if  $f(x) = x$ .
- For the powerset domain:
  - Every element of the domain is a fixpoint for the identity function  $I = \lambda x. x$ .
  - For the function  $G = \lambda x. x \cup \{a\}$ , the elements  $\{a\}$ ,  $\{a, b\}$ ,  $\{a, c\}$ , and  $\{a, b, c\}$  are fixpoints.



# Fixpoint of Function $f: D \rightarrow D$

- Let  $D = (S, \sqsubseteq)$ , and let  $f: D \rightarrow D$  be a function on this domain.
- A value  $x \in S$  is called a fixpoint of  $f$  if  $f(x) = x$ .
- For the powerset domain:
  - Every element of the domain is a fixpoint for the identity function  $I = \lambda x. x$ .
  - For the function  $G = \lambda x. x \cup \{a\}$ , the elements  $\{a\}$ ,  $\{a, b\}$ ,  $\{a, c\}$ , and  $\{a, b, c\}$  are fixpoints.
  - The function  $H = \lambda x. \{a\} \setminus x$  has no fixpoints.




# Fixpoint Theorem #1

- If  $D = (S, \sqsubseteq)$  is a domain,  $\perp$  is its least element, and  $f: D \rightarrow D$  is monotonic, then  $f$  has a **least fixpoint** that is the largest element in the sequence (chain)  $\left[ \perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots \right]$ .


# Fixpoint Theorem #1

- If  $D = (S, \sqsubseteq)$  is a domain,  $\perp$  is its least element, and  $f: D \rightarrow D$  is monotonic, then  $f$  has a least fixpoint that is the largest element in the sequence (chain)  $\left[ \perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots \right]$ .
- Example
  - For the powerset domain,  $\perp = \emptyset$ .



# Fixpoint Theorem #1


- If  $D = (S, \sqsubseteq)$  is a domain,  $\perp$  is its least element, and  $f: D \rightarrow D$  is monotonic, then  $f$  has a least fixpoint that is the largest element in the sequence (chain)  $\left[ \perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots \right]$ .
- Example
  - For the powerset domain,  $\perp = \emptyset$ .
  - For the identity function  $I$ , the sequence is  $[\emptyset, \emptyset, \emptyset, \emptyset, \dots]$ , so the least fixpoint is  $\emptyset$ , which is correct.





# Fixpoint Theorem #1

- If  $D = (S, \sqsubseteq)$  is a domain,  $\perp$  is its least element, and  $f: D \rightarrow D$  is monotonic, then  $f$  has a least fixpoint that is the largest element in the sequence (chain)  $\left[ \perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots \right]$ .
- Example
  - For the powerset domain,  $\perp = \emptyset$ .
  - For the identity function  $I$ , the sequence is  $[\emptyset, \emptyset, \emptyset, \emptyset, \dots]$ , so the least fixpoint is  $\emptyset$ , which is correct.
  - For the function  $G = \lambda x. x \cup \{a\}$ , the sequence is  $[\emptyset, \{a\}, \{a\}, \{a\}, \dots]$ , so the least fixpoint is  $\{a\}$ , which is correct.



# Proof of Fixpoint Theorem #1

- The largest element of the sequence is a fixpoint.
  - $\perp \sqsubseteq f(\perp)$  (by definition of  $\perp$ ).
  - $f(\perp) \sqsubseteq f(f(\perp))$  (from previous fact and monotonicity of  $f$ ).
  - $f(f(\perp)) \sqsubseteq f(f(f(\perp)))$  (same argument).
  - Since the set  $D$  is finite, the chain  $[\perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots]$  cannot grow arbitrarily, so it has some largest element that  $f$  maps to itself. Therefore, we have constructed a fixpoint of  $f$ .

# Proof of Fixpoint Theorem #1


- The largest element of the sequence is a fixpoint.
  - $\perp \sqsubseteq f(\perp)$  (by definition of  $\perp$ ).
  - $f(\perp) \sqsubseteq f(f(\perp))$  (from previous fact and monotonicity of  $f$ ).
  - $f(f(\perp)) \sqsubseteq f(f(f(\perp)))$  (same argument).
  - Since the set  $D$  is finite, the chain  $[\perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots]$  cannot grow arbitrarily, so it has some largest element that  $f$  maps to itself. Therefore, we have constructed a fixpoint of  $f$ .
- This is the least fixpoint.
  - Let  $p$  be any other fixpoint of  $f$ .
  - $\perp \sqsubseteq p$  (from definition of  $\perp$ ).
  - So  $f(\perp) \sqsubseteq f(p) = p$  (by monotonicity of  $f$ ).
  - Similarly,  $f(f(\perp)) \sqsubseteq p$ , etc.
  - Therefore, all elements of the chain are  $\sqsubseteq p$ , so largest element of chain must be  $\sqsubseteq p$ .
  - Therefore, the largest element of chain is the least fixpoint of  $f$ .

# Solving A Fixpoint Equation

- If  $D$  is a domain and  $f: D \rightarrow D$  is monotonic, then the equation  $x = f(x)$  has a least solution given by the largest element in the sequence  $[\perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots]$ .
  - Proof: follows trivially from fixpoint theorem #1.


# Solving A Fixpoint Equation

- If  $D$  is a domain and  $f: D \rightarrow D$  is monotonic, then the equation  $x = f(x)$  has a least solution given by the largest element in the sequence  $[\perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots]$ .
  - Proof: follows trivially from fixpoint theorem #1.
- Generalization #1: The proof goes through even if  $D$  is not a finite set but only has finite height.
  - All we need is that there be no infinite chains.



# Solving A Fixpoint Equation

- If  $D$  is a domain and  $f: D \rightarrow D$  is monotonic, then the equation  $x = f(x)$  has a least solution given by the largest element in the sequence  $[\perp, f(\perp), f(f(\perp)), f(f(f(\perp))), \dots]$ .
  - Proof: follows trivially from fixpoint theorem #1.
- Generalization #1: The proof goes through even if  $D$  is not a finite set but only has finite height.
  - All we need is that there be no infinite chains.



- Generalization #2: If  $D$  is a domain with a *greatest* element  $\top$  and  $f: D \rightarrow D$  is monotonic, then the equation  $x = f(x)$  has a *greatest* solution given by the **smallest** element in the descending sequence  $[\top, f(\top), f(f(\top)), f(f(f(\top))), \dots]$ .

# Functions with Multiple Arguments

- If  $D$  is a domain, a function  $f: D \times D \rightarrow D$  that takes two arguments is said to be **monotonic** if it is monotonic in each argument when the other argument is held constant.
  - $\forall x_0, x_1, y \in D: x_0 \sqsubseteq x_1 \implies f(x_0, y) \sqsubseteq f(x_1, y).$
  - $\forall x, y_0, y_1 \in D: y_0 \sqsubseteq y_1 \implies f(x, y_0) \sqsubseteq f(x, y_1).$

# Functions with Multiple Arguments

- If  $D$  is a domain, a function  $f: D \times D \rightarrow D$  that takes two arguments is said to be monotonic if it is monotonic in each argument when the other argument is held constant.
  - $\forall x_0, x_1, y \in D: x_0 \sqsubseteq x_1 \implies f(x_0, y) \sqsubseteq f(x_1, y).$
  - $\forall x, y_0, y_1 \in D: y_0 \sqsubseteq y_1 \implies f(x, y_0) \sqsubseteq f(x, y_1).$
- Fixpoint Theorem #2
  - If  $D$  is a domain and functions  $f, g: D \times D \rightarrow D$  are monotonic, the following system of simultaneous equations has a least solution computed in the obvious way.
$$\begin{aligned}x &= f(x, y) \\ y &= g(x, y)\end{aligned}$$



# Functions with Multiple Arguments

- If  $D$  is a domain, a function  $f: D \times D \rightarrow D$  that takes two arguments is said to be monotonic if it is monotonic in each argument when the other argument is held constant.
  - $\forall x_0, x_1, y \in D: x_0 \sqsubseteq x_1 \implies f(x_0, y) \sqsubseteq f(x_1, y).$
  - $\forall x, y_0, y_1 \in D: y_0 \sqsubseteq y_1 \implies f(x, y_0) \sqsubseteq f(x, y_1).$
- Fixpoint Theorem #2
  - If  $D$  is a domain and functions  $f, g: D \times D \rightarrow D$  are monotonic, the following system of simultaneous equations has a least solution computed in the obvious way.
$$\begin{aligned}x &= f(x, y) \\ y &= g(x, y)\end{aligned}$$
- This theorem generalizes to more than two equations and to the case when  $D$  has a greatest element  $\top$ .

## Formalization: Product Constructor

- Suppose  $D_1 = (S_1, \sqsubseteq_1)$  and  $D_2 = (S_2, \sqsubseteq_2)$  are domains.

# Formalization: Product Constructor

- Suppose  $D_1 = (S_1, \sqsubseteq_1)$  and  $D_2 = (S_2, \sqsubseteq_2)$  are domains.
- Define  $D = D_1 \times D_2 = (S, \sqsubseteq)$  to be the following domain:
  - Set  $S = S_1 \times S_2$ , i.e., elements are ordered pairs in which the first member is from  $S_1$  and the second is from  $S_2$ .
  - The ordering relation  $\sqsubseteq$  is defined by
$$\langle d_1, d_2 \rangle \sqsubseteq \langle d_3, d_4 \rangle \equiv d_1 \sqsubseteq_1 d_3 \wedge d_2 \sqsubseteq_2 d_4.$$

# Formalization: Product Constructor

- Suppose  $D_1 = (S_1, \sqsubseteq_1)$  and  $D_2 = (S_2, \sqsubseteq_2)$  are domains.
- Define  $D = D_1 \times D_2 = (S, \sqsubseteq)$  to be the following domain:
  - Set  $S = S_1 \times S_2$ , i.e., elements are ordered pairs in which the first member is from  $S_1$  and the second is from  $S_2$ .
  - The ordering relation  $\sqsubseteq$  is defined by
$$\langle d_1, d_2 \rangle \sqsubseteq \langle d_3, d_4 \rangle \equiv d_1 \sqsubseteq_1 d_3 \wedge d_2 \sqsubseteq_2 d_4.$$

- Replace the system of equations

$$x = f(x, y)$$

$$y = g(x, y)$$

with one equation  $\langle x, y \rangle = h(\langle x, y \rangle) = \langle f(x, y), g(x, y) \rangle$ .

# Formalization: Product Constructor

- Suppose  $D_1 = (S_1, \sqsubseteq_1)$  and  $D_2 = (S_2, \sqsubseteq_2)$  are domains.
- Define  $D = D_1 \times D_2 = (S, \sqsubseteq)$  to be the following domain:
  - Set  $S = S_1 \times S_2$ , i.e., elements are ordered pairs in which the first member is from  $S_1$  and the second is from  $S_2$ .
  - The ordering relation  $\sqsubseteq$  is defined by
$$\langle d_1, d_2 \rangle \sqsubseteq \langle d_3, d_4 \rangle \equiv d_1 \sqsubseteq_1 d_3 \wedge d_2 \sqsubseteq_2 d_4.$$
- Replace the system of equations
$$\begin{aligned}x &= f(x, y) \\ y &= g(x, y)\end{aligned}$$
with one equation  $\langle x, y \rangle = h(\langle x, y \rangle) = \langle f(x, y), g(x, y) \rangle$ .
- Check that  $D$  is a domain and that  $h$  is a monotonic function.