

复旦大学大数据学院
School of Data Science, Fudan University

魏忠钰

Bayes' Nets: Independence

May 23rd, 2018

Bayes' Nets

- A Bayes' net is an efficient encoding of a probabilistic model of a domain
- Questions we can ask:
 - Inference: given a fixed BN, what is $P(X \mid e)$?
 - Representation: given a BN graph, what kinds of distributions can it encode?
 - Modeling: what BN is most appropriate for a given domain?



Conditional Independence

- X and Y are **independent** if

$$\forall x, y \quad P(x, y) = P(x)P(y) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y$$


- X and Y are **conditionally independent** given Z

$$\forall x, y, z \quad P(x, y|z) = P(x|z)P(y|z) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y|Z$$

- (Conditional) independence is a property of a distribution

- Example:

$$\textit{Alarm} \perp\!\!\!\perp \textit{Fire} | \textit{Smoke}$$




Bayes Nets: Assumptions


- Assumptions we are required to make to define the Bayes net when given the graph:

$$P(x_i | x_1 \cdots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

- Beyond above “chain rule \rightarrow Bayes net” conditional independence assumptions
 - Often additional conditional independences
 - They can be read off the graph
- Important for modeling: understand assumptions made when choosing a Bayes net graph




Example



- Conditional independence assumptions directly from simplifications in chain rule:
- Additional implied conditional independence assumptions?


Independence in a BN

- Important question about a BN:
 - Are two nodes independent given certain evidence?
 - If yes, can prove using algebra (tedious in general)
 - If no, can prove with a counter example
 - Example:



- Question: are X and Z necessarily independent?
 - Answer: no. Example: low pressure causes rain, which causes traffic.
 - X can influence Z, Z can influence X (via Y)
 - They *could* be independent: how?

D-separation: Outline



- Study independence properties for triples
- Analyze complex cases in terms of member triples
- D-separation: a condition / algorithm for answering such queries

Causal Chains

- This configuration is a "causal chain" ■ Guaranteed X independent of Z ? *No!*



X: Low pressure

Y: Rain

Z: Traffic

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:

- Low pressure causes rain causes traffic, high pressure causes no rain causes no traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- In numbers:

$$P(+y \mid +x) = 1, P(-y \mid -x) = 1, \\ P(+z \mid +y) = 1, P(-z \mid -y) = 1$$

Causal Chains

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

- Guaranteed Given Y, X independent of Z?

$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\ &= P(z|y) \end{aligned}$$


Yes!

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Evidence along the chain “blocks” the influence

Common Cause

- Guaranteed X independent of Z ? *No!*
 - One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.
- Example:
 - Project due causes both Email box busy and library full
 - In numbers
$$P(+x \mid +y) = 1, P(-x \mid -y) = 1,$$
$$P(+z \mid +y) = 1, P(-z \mid -y) = 1$$



$$P(x, y, z) = P(y)P(x|y)P(z|y)$$


Common Cause

- Guaranteed X and Z independent given Y ?

$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{P(y)P(x|y)P(z|y)}{P(y)P(x|y)} \\ &= P(z|y) \end{aligned}$$

Yes!


Observing the cause blocks influence between effects.




$$P(x, y, z) = P(y)P(x|y)P(z|y)$$

Common Effect

- Are X and Y independent?
 - Yes: the ballgame and the rain cause traffic, but they are not correlated
 - Still need to prove they must be
- Are X and Y independent given Z?
 - No: seeing traffic puts the rain and the ballgame in competition as explanation.
- This is backwards from the other cases
 - Observing an effect **activates** influence between possible causes.




The General Case





The General Case

- General question: in a given BN, are two variables independent (given evidence)?
- Solution: analyze the graph
- Any complex example can be broken into repetitions of the three canonical cases



Reachability


- Recipe: shade evidence nodes, look for paths in the resulting graph
- Attempt 1: if two nodes are connected by an undirected path not blocked by a shaded node, they are conditionally independent
- Almost works, but not quite
 - Where does it break?
 - Answer: the v-structure at T doesn't count as a link in a path unless "active"




Active / Inactive Paths

- Question: Are X and Y conditionally independent given evidence variables {Z}?
 - Yes, if X and Y “d-separated” by Z
 - Consider all (undirected) paths from X to Y
 - No active paths = independence!
- A path is active if every triple is active:
 - Causal chain $A \rightarrow B \rightarrow C$ where B is unobserved (either direction)
 - Common cause $A \leftarrow B \rightarrow C$ where B is unobserved
 - Common effect (aka v-structure)
 $A \rightarrow B \leftarrow C$ where B or one of its descendants is observed
- All it takes to block a path is a single inactive segment

Inactive Triples



Active Triples



D-Separation

■ Query: $X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$?

■ Check all (undirected!) paths between X_i and X_j

■ If one or more active, then **independence not guaranteed**

$$X_i \not\perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$

■ Otherwise (i.e. if all paths are inactive),
then independence is guaranteed

$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$


Example

$$R \perp\!\!\!\perp B$$

Yes

$$R \perp\!\!\!\perp B | T$$

$$R \perp\!\!\!\perp B | T'$$



Example


$L \perp\!\!\!\perp T' | T$ *Yes*

$L \perp\!\!\!\perp B$ *Yes*

$L \perp\!\!\!\perp B | T$

$L \perp\!\!\!\perp B | T'$

$L \perp\!\!\!\perp B | T, R$ *Yes*



Example

- Variables:

- R: Raining
- T: Traffic
- D: Roof drips
- S: I'm sad


- Questions:

$$T \perp\!\!\!\perp D$$

$$T \perp\!\!\!\perp D | R$$

$$T \perp\!\!\!\perp D | R, S$$

Yes





- Given a Bayes net structure, can run d-separation algorithm to build a complete list of conditional independences that are necessarily true of the form

$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$

- This list determines the set of probability distributions that can be represented

Computing All Independences


COMPUTE ALL THE
INDEPENDENCES!





Topology Limits Distributions

- Given some graph topology G , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution


$$\{X \perp\!\!\!\perp Y, X \perp\!\!\!\perp Z, Y \perp\!\!\!\perp Z, \\ X \perp\!\!\!\perp Z \mid Y, X \perp\!\!\!\perp Y \mid Z, Y \perp\!\!\!\perp Z \mid X\}$$



$$\{X \perp\!\!\!\perp Z \mid Y\}$$



$$\{\}$$



- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone
- A Bayes' net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

- Representation
- Conditional Independences
- Probabilistic Inference
 - Enumeration (exact, exponential complexity)
 - Variable elimination (exact, worst-case exponential complexity, often better)
 - Probabilistic inference is NP-complete
 - Sampling (approximate)
- Learning Bayes' Nets from Data